

FURTHER BOUNDS FOR THE ESTIMATION ERROR VARIANCE OF A CONTINUOUS STREAM WITH STATIONARY VARIOGRAM

S. S. DRAGOMIR, N. S. BARNETT, AND I. S. GOMM

ABSTRACT. In this paper we establish an upper bound for the estimation error variance of a continuous stream with a stationary variogram V which is assumed to be of the r -Hölder type (Lipschitzian) on $[-d, d]$. Functional properties for the mapping $\xi(t) := E[(\bar{X} - X(t))^2]$, $t \in [0, d]$, are also given.

1. INTRODUCTION

In [1], the authors considered $X(t)$ as defining the quality of a product at time t where $X(t)$ is a continuous time stochastic process which may be non-stationary. Typically, $X(t)$ represents a continuous stream industrial process such as is common in many areas of the chemical industry.

The paper [1] was concerned with issues related to sampling the stream with a view to estimating the mean quality characteristic of the flow, \bar{X} , over the interval $[0, d]$. Specifically, focus was on obtaining the sampling location, said to be optimal, which minimizes the estimation error variance, $E[(\bar{X} - X(t))^2]$, $0 \leq t \leq d$.

Given that t is as specified, the problem is to find the value of t (the sampling location) that minimizes $E[(\bar{X} - X(t))^2]$. It is shown that for constant stream flows, the optimal sampling point is the midpoint of $[0, d]$ for situations where the process variogram,

$$V(u) = \frac{1}{2} E[(\bar{X} - X(t))^2],$$

where

$$V(0) = 0, \quad V(-u) = V(u)$$

is stationary (note that variogram stationarity is not equivalent to process stationarity).

The paper [1] continues to consider optimal sampling locations for situations where the stream flow rate varies. The optimal sampling location is seen to depend on both the flow rate function and the form of the process variogram - some examples are given.

In [2], rather than focussing on the optimal sampling point, the authors have focussed on the actual value of the estimation error variance itself. They obtained the following result by employing an inequality of the Ostrowski type for double integrals.

1991 *Mathematics Subject Classification*. Primary 62 X xx; Secondary 26 D 15.

Key words and phrases. Error variance, Continuous stream with stationary variogram.

Theorem 1. *Let $V : (-d, d) \rightarrow \mathbb{R}$ be a twice differentiable variogram having the second derivative $V'' : (-d, d) \rightarrow \mathbb{R}$ which is bounded. If $\|V''\|_\infty := \sup_{t \in (-d, d)} |V''(t)| < \infty$, then*

$$(1.1) \quad E \left[(\bar{X} - X(t))^2 \right] \leq \left[\frac{1}{4} + \frac{\left(t - \frac{d}{2}\right)^2}{d^2} \right]^2 d^2 \|V''\|_\infty$$

for all $t \in [0, d]$.

Note that the best inequality we can get from (1.1) is that one for which $t = t_0 = \frac{d}{2}$ giving the bound

$$E \left[(\bar{X} - X(t_0))^2 \right] \leq \frac{d^2}{16} \|V''\|_\infty.$$

It should be noted that the above result requires double differentiability of V in $(-d, d)$ and that this condition does not hold for the case of a linear variogram. That is,

$$V(u) = a|u|, \quad u \in \mathbb{R}.$$

For other results on Ostrowski's inequality we refer to the recent papers [3]- [7] and the book [8].

In this note we point out another bound for the estimation error variance which does not require the differentiability of V . Some functional properties are also given.

2. THE RESULTS

Firstly, let us recall the concept of the r -Hölder type mappings.

Definition 1. *The mapping $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ is said to be of the r -Hölder type with $r \in (0, 1]$ if*

$$(2.1) \quad |f(x) - f(y)| \leq H|x - y|^r$$

for all $x, y \in [a, b]$ with a certain $H > 0$.

If $r = 1$, we get the classical concept of Lipschitzian mappings.

Example 1. *If $r \in (0, 1]$, then the mapping $f(x) = x^r$ satisfies the condition*

$$(2.2) \quad |f(x) - f(y)| = |x^r - y^r| \leq |x - y|^r \text{ for all } x, y \in [0, \infty),$$

which shows that f is of the r -Hölder type with the constant $H = 1$, on every closed interval $[a, b]$.

Example 2. *Any differentiable mapping $f : [a, b] \rightarrow \mathbb{R}$ having the derivative bounded in (a, b) is Lipschitzian on (a, b) .*

The following result holds.

Theorem 2. *Assume that the variogram $V : [-d, d] \rightarrow \mathbb{R}$ is of the r -Hölder type on $[-d, d]$ with the constant $H > 0$. Then we have the inequality*

$$(2.3) \quad E \left[(\bar{X} - X(t))^2 \right] \leq \frac{2H}{d} \left[\frac{t^{r+1} + (d-t)^{r+1}}{r+1} \right] \leq \frac{2Hd}{r+1}$$

for all $t \in [0, d]$.

Proof. From [1], using an identity given in [9], it can be shown that

$$\begin{aligned} 0 &\leq E \left[(\bar{X} - X(t))^2 \right] \\ &= \frac{2}{d} \left\{ \int_0^t V(u) du + \int_0^{d-t} V(u) du \right\} - \frac{1}{d^2} \int_0^d \int_0^d V(v-u) dudv. \end{aligned}$$

Also, observe that (see [1])

$$\int_0^d V(v-t) dv = \int_0^t V(u) du + \int_0^{d-t} V(u) du$$

and

$$\int_0^d V(t-u) du = \int_0^t V(u) du + \int_0^{d-t} V(u) du$$

and then we get the identity

$$\begin{aligned} (2.4) \quad 0 &\leq E \left[(\bar{X} - X(t))^2 \right] \\ &= \frac{1}{d} \int_0^d V(v-t) dv + \frac{1}{d} \int_0^d V(t-u) du \\ &= \frac{1}{d^2} \left[d \int_0^d V(v-t) dv + d \int_0^d V(t-u) du \right. \\ &\quad \left. - \int_0^d \int_0^d V(v-u) dvdu \right] \\ &= \frac{1}{d^2} \int_0^d \int_0^d [V(v-t) + V(t-u) - V(v-u)] dvdu. \end{aligned}$$

Using the fact that V is of the r -Hölder type, we can write that

$$\begin{aligned} (2.5) \quad |V(v-t) - V(v-u)| &\leq H |v-t-v+u|^r \\ &= H |u-t|^r \end{aligned}$$

for all $u, v, t \in [0, d]$ and

$$(2.6) \quad |V(t-u)| = |V(t-u) - V(0)| \leq H |t-u|^r$$

for all $t, x \in [0, d]$.

Now, using (2.4) – (2.6), we get

$$\begin{aligned}
E \left[(\bar{X} - X(t))^2 \right] &= \left| \frac{1}{d^2} \int_0^d \int_0^d [V(v-t) + V(t-u) - V(v-u)] dvdu \right| \\
&\leq \frac{1}{d^2} \int_0^d \int_0^d |V(v-t) - V(v-u) + V(t-u)| dvdu \\
&\leq \frac{1}{d^2} \int_0^d \int_0^d |V(v-t) - V(v-u)| + |V(t-u)| dvdu \\
&\leq \frac{1}{d^2} \int_0^d \int_0^d [H|t-u|^r + H|t-u|^r] dvdu \\
&= \frac{2H}{d} \int_0^d |t-u|^r du \\
&= \frac{2H}{d} \left[\int_0^t (t-u)^r du + \int_t^d (u-t)^r du \right] \\
&= \frac{2H}{d} \left[\frac{t^{r+1} + (d-t)^{r+1}}{r+1} \right]
\end{aligned}$$

and the first inequality in (2.3) is proved. The second part is obvious. ■

Corollary 1. *If V is Lipschitzian with the constant $L > 0$, then we have the inequality:*

$$(2.7) \quad E \left[(\bar{X} - X(t))^2 \right] \leq \left[\frac{1}{4} + \frac{(t - \frac{d}{2})^2}{d^2} \right] Ld.$$

Proof. Choose $r = 1$ to get in the right hand side of the inequality (2.3) :

$$\frac{t^2 + (d-t)^2}{2} = \left[\frac{1}{4} + \frac{(t - \frac{d}{2})^2}{d^2} \right] d^2.$$

Then, by (2.3), we deduce (2.7). ■

Remark 1. *It is easy to see that the mapping $g : [0, d] \rightarrow \mathbb{R}$, $g(t) := t^{r+1} + (d-t)^{r+1}$ has the properties*

$$\inf_{t \in [0, d]} g(t) = g\left(\frac{d}{2}\right) = \frac{d^{r+1}}{2^r}$$

and

$$\sup_{t \in [0, d]} g(t) = g(0) = g(d) = d^{r+1}$$

which shows that the best inequality we can get from (2.3) is that one for which $t = t_0 = \frac{d}{2}$, getting

$$(2.8) \quad E \left[(\bar{X} - X(t_0))^2 \right] \leq \frac{2^{1-r} H d^r}{r+1}.$$

For the Lipschitzian case, we get

$$(2.9) \quad E \left[(\bar{X} - X(t_0))^2 \right] \leq \frac{1}{2} Ld.$$

Define the mapping $\xi : [0, d] \rightarrow \mathbb{R}$ given by

$$\xi(t) = E \left[(\bar{X} - X(t))^2 \right].$$

The following property of continuity for ξ holds.

Theorem 3. *If V is of the r -Hölder type with the constant $H > 0$ on the interval $[0, d]$, then ξ is of the r -Hölder type with the constant $2H$.*

Proof. Let $t_1, t_2 \in [0, d]$. Then we have

$$\begin{aligned} & |\xi(t_2) - \xi(t_1)| \\ &= \left| \frac{1}{d^2} \int_0^d \int_0^d [V(v-t_2) + V(t_2-u) - V(v-u)] dudv \right. \\ &\quad \left. - \frac{1}{d^2} \int_0^d \int_0^d [V(v-t_1) + V(t_1-u) - V(v-u)] dudv \right| \\ &= \left| \frac{1}{d^2} \int_0^d \int_0^d [(V(v-t_2) - V(v-t_1)) + (V(t_2-u) - V(t_1-u))] dudv \right| \\ &\leq \frac{1}{d^2} \int_0^d \int_0^d [|V(v-t_2) - V(v-t_1)| + |V(t_2-u) - V(t_1-u)|] dudv \\ &\leq \frac{1}{d^2} \int_0^d \int_0^d [H|t_2-t_1|^r + H|t_2-t_1|^r] dudv \\ &= \frac{2H|t_2-t_1|^r d^2}{d^2} \\ &= 2H|t_2-t_1|^r \end{aligned}$$

and the theorem thus proved. ■

Corollary 2. *If V is L -Lipschitzian on $[0, d]$, then ξ is $2L$ -Lipschitzian on $[0, d]$.*

The following result concerning the convexity property of the mapping ξ defined above on $[0, d]$ holds.

Theorem 4. *If the variogram $V : [-d, d]$ is monotonic nondecreasing on the interval $[0, d]$, then $\xi(\cdot)$ is convex on $[0, d]$.*

Proof. We know that for all $t \in [0, d]$

$$\xi(t) = \frac{2}{d} \left\{ \int_0^t V(u) du + \int_0^{d-t} V(u) du \right\} - \frac{1}{d^2} \int_0^d \int_0^d V(v-u) dudv.$$

Then

$$\xi'(t) = \frac{2}{d} [V(t) - V(d-t)].$$

Now, let $t_1, t_2 \in [0, d]$ with $t_2 > t_1$. Then

$$\begin{aligned} & \xi(t_2) - \xi(t_1) - (t_2 - t_1)\xi'(t_1) \\ &= \frac{2}{d} \left\{ \int_0^{t_2} V(u) du + \int_0^{d-t_2} V(u) du \right\} - \frac{2}{d} \left\{ \int_0^{t_1} V(u) du + \int_0^{d-t_1} V(u) du \right\} \\ & \quad - \frac{2}{d} [V(t_1) - V(d-t_1)](t_2 - t_1) \\ &= \frac{2}{d} \left\{ \int_{t_1}^{t_2} V(u) du - \int_{d-t_2}^{d-t_1} V(u) du - (t_2 - t_1)V(t_1) + (t_2 - t_1)V(d-t_1) \right\}. \end{aligned}$$

As V is nondecreasing on $[0, d]$, then

$$\int_{t_1}^{t_2} V(u) du \geq (t_2 - t_1)V(t_1)$$

and

$$\int_{d-t_2}^{d-t_1} V(u) du \leq (t_2 - t_1)V(d-t_1)$$

which implies that

$$\xi(t_2) - \xi(t_1) \geq (t_2 - t_1)\xi'(t_1)$$

for all $t_2 > t_1 \in [0, d]$, which shows that the mapping $\xi(\cdot)$ is convex on $[0, d]$. ■

REFERENCES

- [1] N.S. Barnett, I.S. Gomm, and L. Armour: Location of the optimal sampling point for the quality assessment of continuous streams, *Australian J. Statistics*, **37**(2), 1995, 145-152.
- [2] N.S. Barnett and S.S. Dragomir, A note on bounds for the estimation error variance of a continuous stream with stationary variogram, *J. KSIAM*, Vol. **2** (2)(1998), 49-56.
- [3] N.S. Barnett and S.S. Dragomir: An Ostrowski's type inequality for double integrals and applications to cubature formulae, *submitted*.
- [4] S.S. Dragomir and S. Wang, A new inequality of Ostrowski's type in L_1 norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. of Math.*, **28** (1997), 239-244.
- [5] S.S. Dragomir and S. Wang, An inequality of Ostrowski-Grüss' type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules, *Computers Math. Applic.*, **33**(1997), 15-20.
- [6] S.S. Dragomir and S. Wang, Applications of Ostrowski's inequality to the estimation of error bounds for some special means and some numerical quadrature rules, *Appl. Math. Lett.*, **11** (1998), 105-109.
- [7] S.S. Dragomir and S. Wang, A new inequality of Ostrowski's type in L_p norm and applications to some special means and to some numerical quadrature rules, *submitted*.
- [8] D.S. MITRINOVIĆ, J.E. PEČARIĆ and A.M. FINK: *Inequalities for Functions and Their Integrals and Derivatives*, Kluwer Academic Publishers, 1994.
- [9] I.W. Saunders, G.K. Robinson, T. Lwin and R.J. Holmes, A simplified variogram method for the estimation error variance in sampling from continuous stream, *Internat. J. Mineral Processing*, **25**(1989), 175-198.

School of Communications and Informatics
Victoria University of Technology,
PO Box 14428, MC Melbourne City,
8001 Victoria, Australia.
email{neil, sever, isg}@matilda.vu.edu.au
<http://rgmia.vu.edu.au>