

## ON PARTITION CONGRUENCES FOR OVERCUBIC PARTITION PAIRS

BYUNGCHAN KIM

ABSTRACT. In this note, we investigate partition congruences for a certain type of partition function, which is named as the overcubic partition pairs in light of the literature. Let  $\overline{cp}(n)$  be the number of overcubic partition pairs. Then we will prove the following congruences:

$$\overline{cp}(8n + 7) \equiv 0 \pmod{64} \quad \text{and} \quad \overline{cp}(9m + 3) \equiv 0 \pmod{3}.$$

### 1. Introduction

In a paper [3], H.-C. Chan initiated the study of the cubic partitions by showing a close relation between a certain type of partition function and Ramanujan's cubic continued fraction. Cubic partition function  $c(n)$  is defined by

$$\sum_{n=0}^{\infty} c(n)q^n = \frac{1}{(q; q)_{\infty} (q^2; q^2)_{\infty}}.$$

Here and in the sequel, we will use the following standard  $q$ -series notation:

$$(a; q)_{\infty} := \prod_{n=1}^{\infty} (1 - aq^{n-1}), \quad |q| < 1.$$

Motivated by his works ([3, 4, 5]), many partition congruences for analogous partition functions have been investigated. In particular, the author studied its overpartition analog [7] in which the overcubic partition function  $\overline{c}(n)$  was defined by

$$(1.1) \quad \sum_{n=0}^{\infty} \overline{c}(n)q^n = \frac{(-q; q)_{\infty} (-q^2; q^2)_{\infty}}{(q; q)_{\infty} (q^2; q^2)_{\infty}}.$$

---

Received April 26, 2011.

2010 *Mathematics Subject Classification.* 11P83.

*Key words and phrases.* partition, cubic partition, overcubic partition pair.

More recently, H. Zhao and Z. Zhong [13] investigated congruences for the following partition function:

$$(1.2) \quad \sum_{n=0}^{\infty} cp(n)q^n = \frac{1}{(q; q)_{\infty}^2 (q^2; q^2)_{\infty}^2}.$$

Since  $cp(n)$  counts a pair of cubic partitions, we will call  $cp(n)$  the number of cubic partition pairs. We can interpret  $cp(n)$  as the number of 4-color partitions of  $n$  with colors  $r$ ,  $y$ ,  $o$ , and  $b$  subject to the restriction that the colors  $o$  and  $b$  appear only in even parts.

In this article, we will investigate congruence properties of the following partition function  $\overline{cp}(n)$ , which is named as the number of overcubic partition pairs in light of (1.1) and (1.2),

$$(1.3) \quad \sum_{n=0}^{\infty} \overline{cp}(n)q^n = \frac{(-q; q)_{\infty}^2 (-q^2; q^2)_{\infty}^2}{(q; q)_{\infty}^2 (q^2; q^2)_{\infty}^2}.$$

We can interpret  $\overline{cp}(n)$  as the number of 4-color partitions of  $n$  with colors  $r$ ,  $y$ ,  $o$ , and  $b$  subject to the restriction that the colors  $o$  and  $b$  appear only in even parts in which we may overline the first occurrence parts.

First, we will investigate congruences for  $\overline{cp}(n)$  modulo 64 by using arithmetic properties of quadratic forms.

**Theorem 1.** *For all nonnegative integers  $n$ ,*

$$\overline{cp}(8n + 7) \equiv 0 \pmod{64}.$$

Next, we find the following congruence for  $\overline{cp}(n)$  via the aid of modular forms.

**Theorem 2.** *For all nonnegative integers  $n$ ,*

$$\overline{cp}(9n + 3) \equiv 0 \pmod{3}.$$

Combining the above two congruences, we can obtain the following congruences.

**Corollary 3.** *For all nonnegative integer  $n$ ,*

$$\overline{cp}(72n + 39) \equiv 0 \pmod{192}.$$

This paper is organized as follows. In Section 2, we review a 2-adic generating function of the overpartition function and basic properties of modular forms. In Section 3, we prove our results.

## 2. Preliminaries

Let  $\overline{P}(q)$  be

$$\overline{P}(q) = \frac{(-q; q)_{\infty}}{(q; q)_{\infty}},$$

and  $\varphi(q)$  be

$$\varphi(q) = \sum_{n=1}^{\infty} q^{n^2}.$$

Then, the coefficients of  $\varphi(q)^k = \sum_{n \geq 1} c_k(n)q^n$  are the number of representations of  $n = n_1^2 + \dots + n_k^2$  where each  $n_i$  is a positive integer. From Mahlburg’s paper [10], we know that

$$(2.1) \quad \overline{P}(q) = 1 + \sum_{k=1}^{\infty} (-2)^k \varphi(-q)^k.$$

By reducing (2.1) modulo 64, we arrive at (2.2)

$$\begin{aligned} \overline{CP}(q) &= \sum_{n=0}^{\infty} \overline{cp}(n)q^n \\ &= \overline{P}(q)^2 \overline{P}(q^2)^2 \\ &= (1 - 2\varphi(-q) + 2^2\varphi(-q)^2 - \dots)^2 (1 - 2\varphi(-q^2) + 2^2\varphi(-q^2)^2 - \dots)^2 \\ &= 1 - 4\varphi(-q) - 4\varphi(-q^2) + 12\varphi(-q)^2 + 12\varphi(-q^2)^2 + 16\varphi(-q)\varphi(-q^2) \\ &\quad + 16\varphi(-q)^4 + 16\varphi(-q^2)^4 - 48\varphi(-q)^2\varphi(-q^2) - 48\varphi(-q)\varphi(-q^2)^2 \\ &\quad - 48\varphi(-q)^2\varphi(-q^2)^2 + 144\varphi(-q)^2\varphi(-q^2)^2 - 32\varphi(-q)^3 \\ &\quad - 32\varphi(-q^2)^3 \pmod{64}. \end{aligned}$$

Now we give some basic properties of modular functions. For more details on this subject, consult [12]. Define

$$\Gamma = SL_2(\mathbb{Z}), \Gamma_0(N) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma : c \equiv 0 \pmod{N} \right\}.$$

For a meromorphic function  $f$  on the complex upper half plane  $\mathcal{H}$ , define the slash operator by

$$f| \begin{pmatrix} a & b \\ c & d \end{pmatrix} := f\left(\frac{az + b}{cz + d}\right).$$

We say a meromorphic function  $f$  is a modular function on  $\Gamma_0(N)$  if  $f$  is invariant under the slash operator. Let  $\mathcal{M}_0(\Gamma_0(N))$  denote the vector space of modular functions on  $\Gamma_0(N)$ . For a prime  $p$ , we define the  $U_p$ -operator as follows. If  $f(q)$  has a Fourier expansion  $f(q) = \sum a(n)q^n$ , then

$$U_p f(z) := \sum a(pn)q^n.$$

It is well known that  $U_p f(z) \in \mathcal{M}_0(\Gamma_0(Np))$  provided  $f(z) \in \mathcal{M}_0(\Gamma_0(Np^2))$ . Let  $\eta(z) = q^{1/24}(q; q)_{\infty}$  be Dedekind eta function, where  $q = \exp(2\pi iz)$  and  $z$  is in the upper half complex plane  $\mathcal{H}$ . For a fixed  $N$  and integers  $r_i$ ’s, a

function of the form

$$(2.3) \quad f(z) := \prod_{\substack{n|N \\ n>0}} \eta(nz)^{r_n}$$

is called an  $\eta$ -quotient. By the famous theorem of Newman [11], we can determine when an  $\eta$ -quotient becomes a modular function. We call the orbits of  $\mathbb{Q} \cup \{i\infty\} \pmod{\Gamma_0(N)}$  the cusps of  $\Gamma_0(N)$ . Moreover, by the work of Ligozat [9], we can calculate the order of the  $\eta$ -quotient  $f$  at the cusps  $c/d$  of  $\Gamma_0(N)$  provided  $f \in \mathcal{M}_0(\Gamma_0(N))$ . Recall that if  $p|N$  and  $f \in \mathcal{M}_0(\Gamma_0(pN))$ , then  $U_p f \in \mathcal{M}_0(\Gamma_0(N))$ . The following theorem gives bounds on the order of  $U_p f$  at cusps of  $\Gamma_0(N)$  in terms of the order of  $f$  at cusps of  $\Gamma_0(pN)$ .

**Theorem 4** (Theorem 4 of [6]). *Let  $p$  be a prime and  $\pi(n)$  be the highest power of  $p$  dividing  $n$ . Suppose that  $f \in \mathcal{M}_0(\Gamma_0(pN))$ , where  $p|N$  and  $\alpha = c/d$  is a cusp of  $\Gamma_0(N)$ . Then,*

$$\text{ord}_\alpha U_p f \geq \begin{cases} \frac{1}{p} \text{ord}_{\alpha/p} f, & \text{if } \pi(d) \geq \frac{1}{2} \pi(N), \\ \text{ord}_{\alpha/p} f, & \text{if } 0 < \pi(d) < \frac{1}{2} \pi(N), \\ \min_{0 \leq \beta \leq p-1} \text{ord}_{(\alpha+\beta)/p} f, & \text{if } \pi(d) = 0. \end{cases}$$

### 3. Proofs of theorems

We now start the proof of Theorem 1.

*Proof of Theorem 1.* For a quadratic form  $Q = \sum_{i=1}^k a_i x_i^2$ , let  $R(n, Q)$  be the number of representations of  $n$  by  $n = a_1 n_1^2 + a_2 n_2^2 + \dots + a_k n_k^2$ , where  $a_i$ 's and  $n_i$ 's are positive integers. By (2.2), it suffices to show the following congruences:

$$\begin{aligned} R(8n + 7, x_1^2) &\equiv R(8n + 7, 2x_1^2) \\ &\equiv R(8n + 7, x_1^2 + x_2^2) \\ &\equiv R(8n + 7, 2x_1^2 + 2x_2^2) \equiv 0 \pmod{16}, \\ R(8n + 7, x_1^2 + 2x_2^2) &\equiv R(8n + 7, x_1^2 + x_2^2 + 2x_3^2) \\ &\equiv R(8n + 7, x_1^2 + 2x_2^2 + 2x_3^2) \\ &\equiv R(8n + 7, x_1^2 + x_2^2 + x_3^2 + x_4^2) \\ &\equiv R(8n + 7, 2x_1^2 + 2x_2^2 + 2x_3^2 + 2x_4^2) \\ &\equiv R(8n + 7, x_1^2 + x_2^2 + 2x_3^2 + 2x_4^2) \equiv 0 \pmod{4}, \\ R(8n + 7, x_1^2 + x_2^2 + x_3^2) &\equiv R(8n + 7, 2x_1^2 + 2x_2^2 + 2x_3^2) \equiv 0 \pmod{2}. \end{aligned}$$

Note that  $x^2 \equiv 0, 1, \text{ or } 4 \pmod{8}$ . By using this, we can see that there is no integer solution for  $8n + 7 = x_1^2 + x_2^2 + 2x_3^2$  and we also see that if  $8n + 7 = x_1^2 + x_2^2 + x_3^2 + x_4^2$ , then at least one of  $x_i$  is different from others. By using a similar argument, we can verify the above congruences hold.  $\square$

*Remark.* By proceeding as in [8], one can prove that  $\{n \in \mathbb{N} : \overline{cp}(n) \equiv 0 \pmod{2^k}\}$  has the arithmetic density 1 for any fixed positive integer  $k$ . As we focus on explicit congruences in this article, we will omit the proof of this fact.

Now we turn to the proof of Theorem 2.

*Proof of Theorem 2.* Let define  $G(z)$  as

$$G(z) = \frac{\eta^2(4z)\eta^4(9z)\eta^2(18z)}{\eta^4(z)\eta^2(2z)\eta^2(36z)}.$$

Then, by applying Newman’s theorem, we see that  $G(z) \in \mathcal{M}(\Gamma_0(36))$ . By employing  $U_3$  operator, we see that  $U_3G(z) \in \mathcal{M}(\Gamma_0(12))$  and arrive at

$$U_3G(z) = \left( \sum_{n=0}^{\infty} \overline{cp}(3n)q^n \right) \frac{(q^3; q^3)_{\infty}^4 (q^9; q^9)_{\infty}^2}{(q^{12}; q^{12})_{\infty}^2}.$$

By applying Theorem 4, we can find lower bounds for the orders of  $U_3G(z)$  at the cusps as follows:

d	1	2	3	4	6	12
$\text{ord}_{c/d}U_3G(z) \geq$	-6	-6	2	-6	1	0

After multiplying  $\Delta(z)^2 = \eta^{48}(z)$  to  $U_3G(z)$ , we observe that  $U_3G(z)\Delta(z)^2$  is in  $\mathcal{M}_{24}(\Gamma_0(12))$  and  $U_3G(z)\Delta(z)^2$  is a holomorphic modular form and so is  $U_3(U_3G(z)\Delta(z)^2)$ . Moreover,

$$\begin{aligned} U_3(U_3G(z)\Delta(z)^2) &\equiv U_3(U_3G(z)\eta^{16}(3z)) \pmod{3} \\ &\equiv U_3 \left( \sum_{n=0}^{\infty} \overline{cp}(3n)q^{n+2} \right) \frac{(q; q)_{\infty}^{20} (q^3; q^3)_{\infty}^2}{(q^4; q^4)_{\infty}^2} \pmod{3} \\ &\equiv \left( \sum_{n=1}^{\infty} \overline{cp}(9n - 6)q^n \right) \frac{(q; q)_{\infty}^{20} (q^3; q^3)_{\infty}^2}{(q^4; q^4)_{\infty}^2} \pmod{3}. \end{aligned}$$

Since the dimension of holomorphic modular form of weight 24 with level 12 is 49, by Sturm’s theorem, we can verify that  $\overline{cp}(9n - 6) \equiv 0 \pmod{3}$  by checking up to  $n = 50$ . □

*Remark.* There is no single eta-quotient in  $\mathcal{M}_0(\Gamma_0(12))$  which coincides with  $U_3G(z)$ .

### References

- [1] G. E. Andrews and F. G. Garvan, *Dyson’s crank of a partition*, Bull. Amer. Math. Soc. (N.S.) **18** (1988), no. 2, 167–171.
- [2] B. C. Berndt, *Number Theory in the Spirit of Ramanujan*, American Mathematical Society, Providence, RI, 2006.
- [3] H.-C. Chan, *Ramanujan’s cubic continued fraction and an analog of his “most beautiful identity”*, Int. J. Number Theory **6** (2010), no. 3, 673–680.
- [4] ———, *Ramanujan’s cubic continued fraction and Ramanujan type congruences for a certain partition function*, Int. J. Number Theory **6** (2010), no. 4, 819–834.

- [5] ———, *Distribution of a certain partition function modulo powers of primes*, Acta Math. Sin. (Engl. Ser.) **27** (2011), 625–634.
- [6] B. Gordon and K. Hughes, *Ramanujan congruences for  $q(n)$* , Analytic number theory (Philadelphia, Pa., 1980), 333–359, Lecture Notes in Math., 899, Springer, Berlin-New York, 1981.
- [7] B. Kim, *The overcubic partition function mod 3*, Proceedings of Ramanujan Rediscovered 2009: A Conference in Memory of K. Venkatachaliengar on the Centenary of His Birth, Lecture Note Series of the Ramanujan Mathematical Society **14** (2010), 157–163.
- [8] ———, *Overpartition pairs modulo powers of 2*, Discrete Math. **311** (2011), no. 10–11, 835–840.
- [9] G. Ligozat, *Courbes modulaires de genre 1*, Bull. Soc. Math. France, Mem. 43. Supplement au Bull. Soc. Math. France Tome 103, no. 3. Societe Mathematique de France, Paris, 1975. 80 pp.
- [10] K. Mahlburg, *The overpartition function modulo small powers of 2*, Discrete Math. **286** (2004), no. 3, 263–267.
- [11] M. Newman, *Construction and application of a class of modular functions II*, Proc. London Math. Soc. (3) **9** (1959), 373–387.
- [12] K. Ono, *Web of Modularity: arithmetic of the coefficients of modular forms and  $q$ -series*, CBMS Regional Conference Series in Mathematics, 102. Published for the Conference Board of the Mathematical Sciences, Washington, DC, by the American Mathematical Society, Providence, RI, 2004.
- [13] H. Zhao and Z. Zhong, *Ramanujan type congruences for a partition function*, The Electronic Journal of Combinatorics **18** (2011), P. 58.

SCHOOL OF LIBERAL ARTS  
SEOUL NATIONAL UNIVERSITY OF SCIENCE AND TECHNOLOGY  
SEOUL 139-743, KOREA  
*E-mail address:* bkim4@seoultech.ac.kr