RESEARCH ARTICLE

Evolution of the Mir-155 Family and Possible Targets in Cancers and the Immune System

Guang-Bing Xie¹, Wei-Jia Liu¹, Zhi-Jun Pan¹, Tian-Yin Cheng^{2*}, Chao Luo^{1*}

Abstract

The mir-155 family is not only involved in a diversity of cancers, but also as a regulator of the immune system. However, the evolutionary history of this family is still unclear. The present study indicates that mir-155 evolved independently with lineage-specific gain of miRNAs. In addition, arm switching has occurred in the mir-155 family, and alternative splicing could produce two different lengths of ancestral sequences, implying the alternative splicing can also drive evolution for intragenic miRNAs. Here we screened validated target genes and immunity-related proteins, followed by analysis of the mir-155 family function by high-throughput methods like the gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis. The high-throughput analysis showed that the CCND1 and EGFR genes were outstanding in being significantly enriched, and the target genes cebpb and VCAM1 and the protein SMAD2 were also vital in mir-155-related immune response activities. Therefore, we conclude that the mir-155 family is highly conserved in evolution, and CCND1 and EGFR genes might be potential targets of mir-155 with regard to progress of cancers, while the cebpb and VCAM1 genes and the protein SMAD2 might be key factors in the mir-155 regulated immune activities.

Keywords: mir-155 family - evolution - function - interaction - cancer - immunity

Asian Pac J Cancer Prev, **15** (18), 7547-7552

Introduction

MicroRNAs (miRNAs) are approximately 22nt, endogenous and non-coding RNAs that play important roles in the post-transcriptional repression of organisms (Bartel, 2004). Generally, the miRNAs are conducted by the RNA polymerase II, forming the primary transcripts (pri-miRNAs). Then the pri-miRNAs are cleaved by RNase III enzyme Drosha with the help of DGCR8 (Landthaler et al., 2004; Price et al., 2011), followed by the release of a hairpin-shaped RNA molecule called miRNA precursor (pre-miRNA). The transport of pre-miRNAs into the cytoplasm by the Exportin 5 will then trigger the alternative splicing by Dicer enzyme and form an imperfect symmetry (miRNA: miRNA*duplex), consisted of one mature (3' arm) and one immature (5' arm) sequences (Yi et al., 2003; Zhang et al., 2004; MacRae et al., 2006). After that, the mature one enters the RNA-induced silencing complex (RISC) while the other will be degraded or retained in some species (Schwarz et al., 2003; Bartel, 2004). The sequence analysis of pre-miRNA and miRNA: miRNA*duplex in different species can help us to understand the distribution and evolution of a miRNA family. Besides, it has been hot topic to identify the regulation function of miRNAs in wide ranging

signaling cascades and modulators of different cell-type and context dependent activities (Farooqi et al., 2014).

It is commonly believed that microRNAs are evolutionarily conserved among different organisms (Xu et al., 2012). However, a considerable amount of studies have stated that most microRNAs follows dynamic evolution, especially the novel microRNAs. For example, miR-941 was found to be a rapid regulatory evolution microRNA in the human lineage, which leads to a diversity of evolution patterns (Hu et al., 2012). Then many current studies either focus on the investigation of cluster evolution or the local duplication evolutionary patterns of a microRNA family, for example, the mir-2 family, the let-7 family, the mir-17 family, the mir-395 family, the mir-548 family (Guddeti et al., 2005; Concepcion et al., 2012; Hertel et al., 2012; Liang et al., 2012; Marco et al., 2012a). However, it has been seldom studied how the independent and single-member microRNA family evolves up to date. Moreover, the current studies have usually emphasized on the construction of protein-protein interaction (PPI) network, but pay rare attention on the miRNA regulatory network.

Mir-155 (mir-155) is an important regulatory factor participating in many physiological and pathological processes of organism (O'Connell et al., 2012; Elton et al.,

¹College of Orient Science and Technology, ²College of Veterinary Medicine, Hunan Agriculture University, Changsha, China *For correspondence: superlluo@163.com

2013), definitely such as malignant growth (Babar et al., 2012; Mattiske et al., 2012), viral infections (Vargova et al., 2011; Wang et al., 2011) and cardiovascular diseases (Corsten et al., 2012). Further, the mir-155 family is an independent and single-member's family with unclear evolutionary patterns. Therefore, we extracted all the messages about the mir-155 family in different species, and performed multiple sequence alignment, as well as comparative genomics in the current study. We expect the study can supply new insights in the evolutionary patterns of independent and single-member miRNA family and the miRNA regulatory network.

Materials and Methods

Data source

First, the Taxonomy Tools of the NCBI (<http://www.ncbi.nlm.nih.gov/taxonomy/>) was used to determine the distribution of the mir-155 family. Then all the members of mir-155 family, as well as their corresponding species, annotations, deep sequencing data, miRNA/miRNA* complex and pre-miRNA sequences from different animal species were extracted from the miRase database (Release 20 <http://mirbase.org/index.shtml>) (Kozomara and Griffiths-Jones, 2011).

The construction of Phylogenetic trees

To identify the ancestral sequence, Phylogenetic trees of mir-155 family were reconstructed with MEGA 4.0 (Tamura et al., 2007) by 1,000 bootstrap resampling using Neighbor-Joining (NJ) method. It has been reported that the mir-100 is the only microRNA which is shared by all bilateria (Grimson et al., 2008; Griffiths-Jones et al., 2011). Therefore, we also analyzed the relationship between the mir-100 and mir-155 using Neighbor-Joining (NJ) method. After that, we screened the species containing the ancestral sequences of mir-155 from the mir-100 family. Then we compared the sequences mir-155 and mir-100 of the same species to determine the origin of the mir-155 family.

Relative Arm Usage

To quantify the relative amount of mature miRNA products from each arm of the same miRNA hairpin precursor, we performed the "relative arm usage" measure as reported (Marco et al., 2010). The quantity of relative arm usage was calculated as follows: $\text{Relative arm usage} = \log^2(N_5'/N_3')$; where N_5' is the number of reads mapped to the 5' arm of the hairpin precursor and N_3' the number of reads from the 3' arm. Positive values indicate a bias toward 5' arm usage while the negative values a bias toward 3' arm. Zero means that mature sequences are produced at equal levels from both arms. Besides, miRDB is an online database system used for microRNA target prediction and functional annotation (Wang, 2008). In the current study, we used the miRDB database to predict the target sites of mir-155.

Prediction of the target genes and the target mRNAs

BioMart is a freely available, open source, federated database system that provides a unified access to disparate,

geographically distributed data sources (Zhang et al., 2011). The Ensembl and Ensembl Genomes BioMarts can supply us a point of access to the high-quality gene annotation, variation data, functional and regulatory annotation and evolutionary relationships from genomes spanning the taxonomic space (Kinsella et al., 2011). To predict the expressed target mRNAs of mir-155, we fetched the messages of 3'UTR of the whole-genome from ENSEMBEL (Release 72) via Biomart. Here we only chose the canonical seed to predict the 3'UTRs (Bartel, 2009), and the target sites were performed with miRecords (Xiao et al., 2009) and miRTarbase (Hsu et al., 2011).

Functional and pathway enrichment analysis

Molecule Annotation System (MAS 3.0, <http://bioinfo.capitalbio.com/mas3/>) is convenient analysis management software for the analysis of significantly enriched Gene Ontology (GO), pathway terms with user-friendly web interface and stable and fast analysis procedure. The selected target genes were queried using MAS to obtain richness of the functions and Kyoto Encyclopedia of Genes & Genomes (KEGG) pathway. Based on the hypergeometric distribution, the pathways or GO terms with $p < 0.01$ and $q < 0.01$ were chose for the next analysis.

Construction of an miRNA-target gene- cancer regulatory network

Cytoscape is a popular bioinformatics package for biological network visualization and data integration (Smoot et al., 2011). We selected the cancer-related pathways from the enriched KEGG pathways, and then constructed a regulatory network of mir-155-pathway-target gene with the help of Cytoscape 2.8.

Construction of an mir-155-target gene-protein regulatory network

Human Protein Reference Database (HPRD) an object-oriented database that integrates a great deal of information relevant to the function of human proteins in health and disease (Prasad et al., 2009). In the present study, we chose the genes which were regulated by mir-155 and enriched in the immune related functions after functional enrichment analysis. Then we analyzed and constructed the mir-155-target gene-protein regulatory network using Cytoscape 2.8. All the protein interaction data used here should be verified by experiment before.

Results

Mir-155 family all distributes in vertebrates

The annotation in the miRBase database (Release 20) showed that mir-155 family totally had 20 members. Based on the results of taxonomy browser of NCBI, we found that the members of mir-155 family all distributed in vertebrates (Figure 1).

Mir-155 is a conserved microRNA family in vertebrates

According to the multiple alignment of all precursor sequences (Figure 2a), we found that the 5' arm of the family precursor was highly conserved, and 3' arm has some changes the nucleotides. The highly conserved

Figure 3. The Phylogenic Tree of mir-155 Family. The phylogenic tree was reconstructed by MEGE 4.0. by 1,000 bootstrap resampling (MarcoMacPherson et al., 2012b)

[illegible]

Figure 4. The Relationship Between the mir-155 and mir-100. (B) The ancestral sequences of mir-155 family. (C) The multiple sequence alignment of mature sequences of mir-155 family. We can find the seed sequences of the 5' arm are 100% conserved. B and C alignment were performed by miRBase (Release 20, <http://mirbase.org/index.shtml>) (KinsellaKahari et al., 2011)

between the two ancestral sequences was that the sequence from *D. rerio* was longer than the sequence from *I. punctatus* (Figure 4c). Date have shown the mir-100 is the only microRNA shared by all bilateria (GrimsonSrivastava et al., 2008; Griffiths-JonesHui et al., 2011). Based on the comparison between the mir-100 and mir-155 family in the same species, we found that the original time of mir-155 family was just the time when the same sequences generated in mir-100 family. The original time of the mir-100 and mir-155 families was parallel, and mir-155 family came from the lineage-specific gain of miRNAs, but not from the original miRNAs (Figure 4a).

The mir-155 family is highly conserved in function

To verify whether the arm switching happened in mir-155 or not, we further calculated the “Relative Arm Usage” (Table 1). The 5' arm was highly expressed compared with the 3' arm, which suggested the dominantly mature microRNA was generated from the 5' arm. Based on the

The relative arm usage is defined: $\log_2(N5'/N3')$. Positive values indicate a bias toward 5' arm usage and negative values a bias toward 3' arm. Zero means that mature sequences are produced at equal levels from both arms (Marco et al., 2010). Targets were predicted by miRDB (Wienholds et al., 2005), and "-" indicate there was no data available.

The target gene- pathway-cancer regulatory network of mir-155

There were totally 738 target genes regulated by mir-155 were screened in the current study. All the targets were queried for the KEGG pathway and GO enrichments by using MAS. The KEGG pathway analysis showed mir-155 was involved in a diversity of pathways, especially the various cancers. There were totally 88 pathways, and of them, 11 pathways were presented relation with various cancers (Table 2), such as Colorectal cancer, Pancreatic cancer. Therefore, we constructed the target gene-pathway-cancer regulatory network of mir-155 (Figure 5) to uncover the relationship between mir-155 and cancers. The CCND1, EGFR genes were involved in most of the cancer-related pathways.

The construction of mir-155-target gene-protein regulatory network

Based on the GO analysis, the target genes were enriched in 366 terms of cellular molecular function, 533 of biological processes and 172 of cellular components. With the help of HPRD, there were totally 379 proteins with interaction data with mir-155 target genes were enriched in immune-related GO terms, such as CD31, CRKL and Fgr. The mir-155-target gene-protein regulatory network was shown in Figure 6. The target genes *cebpb* and *VCAM1* and the protein *SMAD2* might be the key factors in the mir-155 regulated immune activities.

Mir-155 has been reported to play important roles in inhibition of malignant growth, viral infections and even in attenuating the progression of cardiovascular

Pathway	Count	<i>p</i> -Value	<i>q</i> -Value	Genes
Colorectal cancer	15	6.31E-14	1.44E-12	APC; MLH1; MSH2; MSH6; CTNNB1; KRAS; SMAD2; JUN; RAC1; SMAD4; EGFR; SMAD3; CCND1; MET; GSK3B
Pancreatic cancer	12	5.73E-11	4.06E-10	KRAS; SMAD2; RAC1; SMAD4; CDKN2A; EGFR; SMAD3; CCND1; STAT3; CDK4; NFKB1; E2F2
Endometrial cancer	10	4.59E-10	2.46E-09	APC; MLH1; FOXO3; CTNNB1; KRAS; CTNNA1; PDPK1; EGFR; CCND1; GSK3B
Melanoma	11	7.05E-10	3.47E-09	FGF7; KRAS; MITF; FGF2; FGF13; CDKN2A; EGFR; CCND1; MET; CDK4; E2F2
Bladder cancer	9	1.24E-09	5.48E-09	KRAS; CDKN2A; EGFR; THBS1; CCND1; IL8; CDK4; MMP1; E2F2
Non-small cell lung cancer	8	2.20E-07	4.92E-07	FOXO3; KRAS; PDPK1; CDKN2A; EGFR; CCND1; CDK4; E2F2
Prostate cancer	9	9.84E-07	1.69E-06	CTNNB1; KRAS; PDPK1; EGFR; CCND1; CDK2; GSK3B; NFKB1; E2F2
Renal cell carcinoma	8	1.89E-06	3.03E-06	HIF1A; KRAS; ETS1; JUN; RAP1B; RAC1; PAK2; MET
Small cell lung cancer	8	8.07E-06	1.17E-05	NOS3; COL4A2; APAF1; CCND1; CDK2; CDK4; NFKB1; E2F2
Basal cell carcinoma	4	3.97 E-03	2.23E-03	APC; CTNNB1; WNT5A; GSK3B
Thyroid cancer	3	4.67 E-03	2.56E-03	CTNNB1; KRAS; CCND1

diseases (WangToomey et al., 2011; Zhi-Yong et al., 2011; CorstenPapageorgiou et al., 2012; MattiskeSuetani et al., 2012). Moreover, mir-155 has been regarded as a critical regulator of immune cell development, function, and disease (O'Connell et al., 2009), however, it is still unclear about the evolution of the mir-155 family. In the current study, we found that the mir-155 family was highly conserved in 3' arm unlike the mir-2 family and there was no gene copy coming out of this family, which made the nucleotide divergence, only emerged among species. The KEGG pathway enrichment analysis showed that the CCND1, EGFR genes might be the potential targets of mir-155 in the progress of cancers. Moreover, the target genes cebpb and VCAM1 and the protein SMAD2 might be the key factors in the mir-155 regulated immune activities.

The evolutionary process of microRNA is slower even than the most conserved genes in the metazoan genome (GrimsonSrivastava et al., 2008; Wheeler et al., 2009). Therefore, it is possible for microRNA families to be used in tracking the evolutionary origin of the most ancient animal phylogeny. In addition, the miRNAs have experienced various evolutionary patterns during the evolutionary history. The accepted evolutionary patterns of microRNA have been shown as duplications (including local duplication and non-local duplication), losses, rearrangements, arm switching, new hairpin, hairpin shifting, seed shifting, editing (Blow et al., 2006; Maher et al., 2006; de Wit et al., 2009; MarcoHui et al., 2010; Griffiths-JonesHui et al., 2011; Marco et al., 2012b; Marco et al., 2013). Of them, duplication is the leading evolutionary pattern, which usually results in the clustered changes in most of lineages, for instance, the mir-2 and mir-17 family. However, the mir-155 family is very different from them in these aspects. The duplications are rarely happened in mir-155 family, as well as the family cluster. Further, the mir-155 family was highly conserved in 3' arm, while the 5' arm was highly expressed because of the arm switching. Therefore, the dominant mature microRNA was produced from the 5' arm, which was consistent with the predicted results of miRDB. The mature microRNAs are composed of non-canonical and canonical mature miRNA. In the current study, we found that the seed sequences of the non-canonical mature miRNA, 2–8 nucleotides of the mature microRNAs which play a crucial role in transcript targeting (LewisShih et al., 2003; NahviShoemaker et al., 2009), have undergone slight changes. Therefore, we speculate that the mir-155 family is suffering slow evolution, and the non-canonical mature miRNA might have the potential to change the mir-155 family.

MiRNA is well-preserved in a range of specimen types and is much sensitivity than proteins, therefore, miRNAs has the potential of biomarkers for various molecular diagnostic applications (Pritchard et al., 2012). Moreover, the mir-155 family is a vital small regulatory RNA in organism, and the function is widely involved in immune adjustment (Vigorito et al., 2013) and a variety of cancers (MattiskeSuetani et al., 2012). Therefore, we constructed the target gene-pathway-cancer regulatory network of mir-155, and found CCND1, EGFR genes might be the key targets of mir-155 in cancer-related pathways. In our study,

CCND1 was enriched in the Colorectal cancer, Pancreatic cancer, Endometrial cancer, Melanoma, Bladder cancer, Non-small cell lung cancer, Prostate cancer, Small cell lung cancer and Thyroid cancer pathways. Moreover, it has been reported CCND1 amplification also occurs in breast, esophageal, hepatocellular, and head and neck cancer (Croce, 2008). The CCND1 gene controls the cell cycle by the regulation on cyclin-dependent kinase 4 (CDK4) or CDK6. The mutations, amplification and overexpression of this gene will result in the alteration of cell cycle progression, and then further disturb normal mitosis, even contribute to tumorigenesis. EGFR gene is related to many epithelial cancers. Therefore, we speculate that mir-155 may have regulatory effect on many cancers which mainly performs through the control of the cell cycle by the CCND1 gene.

Furthermore, as shown in Figure 4, SMAD2 plays a very important role in the regulatory network of mir-155 target genes and immune-related proteins. As known, SMAD2 proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways, and is responsible for the transmission of extracellular signals from ligands of the Transforming Growth Factor beta (TGF- β) superfamily of growth factors into the cell nucleus. TGF- β is a pleiotropic cytokine taking part in the processes of fibrosis, angiogenesis, apoptosis and immunosuppression (Louafi et al., 2010; Das et al., 2014). The TGF- β receptor-Smad2/3 pathway is essential for the cell apoptosis and proliferation (Zhi-Yong1 et al., 2011; DasXu et al., 2014). Moreover, it has been indicated that miR-155 antisense oligonucleotide can induce cell apoptosis and inhibit cell proliferation, and therefore, may be a therapeutic target of breast carcinoma (Zheng et al., 2013). This report also indicated that SMAD2 was a predicted target for mir-155. Therefore, the regulatory effect of mir-155 on immune function mainly performs through the SMAD protein.

In conclusion, the mir-155 family is a vertebrate-specific and conserved family which is suffering slow evolution. In addition, mir-155 family is special for arm usage which the 5' arm acts as dominant function. The CCND1 gene and the SMAD protein might be the key targets of mir-155 in diverse cancer pathways and immune system.

Acknowledgements

This work is supported by College of Orient Science and Technology, College of Veterinary Medicine, College of Animal Science And Technology and Horticulture and Landscape College, Hunan Agriculture University. Conflict of interest: None

References

- Babar IA, Cheng CJ, Booth CJ, et al (2012). Nanoparticle-based therapy in an *in vivo* microRNA-155 (miR-155)-dependent mouse model of lymphoma. *Proc Natl Acad Sci USA*, **109**, 1695-704.
- Bartel DP (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*, **116**, 281-97.
- Bartel DP (2009). MicroRNAs: target recognition and regulatory

- functions. *Cell*, **136**, 215-33.
- Blow MJ, Grocock RJ, van Dongen S, et al (2006). RNA editing of human microRNAs. *Genome biology*, **7**, 27.
- Concepcion CP, Bonetti C, Ventura A (2012). The microRNA-17-92 family of microRNA clusters in development and disease. *Cancer J*, **18**, 262-7.
- Corsten MF, Papageorgiou A, Verhesen W, et al (2012). MicroRNA profiling identifies microRNA-155 as an adverse mediator of cardiac injury and dysfunction during acute viral myocarditis. *Circulation Res*, **111**, 415-25.
- Croce CM (2008). Oncogenes and cancer. *N Engl J Med*, **358**, 502-11.
- Darty K, Denise A, Ponty Y (2009). VARNA: Interactive drawing and editing of the RNA secondary structure. *Bioinformatics*, **25**, 1974.
- Das R, Xu S, Quan X, et al (2014). Upregulation of mitochondrial Nox4 mediates TGF- β -induced apoptosis in cultured mouse podocytes. *Am J Physiol Renal Physiol*, **306**, 155-67.
- de Wit E, Linsen SE, Cuppen E, Berezikov E (2009). Repertoire and evolution of miRNA genes in four divergent nematode species. *Genome Res*, **19**, 2064-74.
- Elton TS, Selemmon H, Elton SM, Parinandi NL (2013). Regulation of the MIR155 host gene in physiological and pathological processes. *Gene*, **532**, 1-12.
- Farooqi AA, Qureshi MZ, Coskunpinar E, et al (2014). miR-421, miR-155 and miR-650: emerging trends of regulation of cancer and apoptosis. *Asian Pac J Cancer Prev*, **15**, 1909-12.
- Griffiths-Jones S, Hui JH, Marco A, Ronshaugen M (2011). MicroRNA evolution by arm switching. *EMBO Rep*, **12**, 172-7.
- Grimson A, Srivastava M, Fahey B, et al (2008). Early origins and evolution of microRNAs and Piwi-interacting RNAs in animals. *Nature*, **455**, 1193-7.
- Guddeti S, De Chun ZHANG ALL, LESEBERG C H, et al (2005). Molecular evolution of the rice miR395 gene family. *Cell Res*, **15**, 631-8.
- Hertel J, Bartschat S, Wintsche A, et al (2012). Evolution of the let-7 microRNA Family. *RNA biology*, **9**, 231-41.
- Hsu SD, Lin FM, Wu WY, et al (2011). miRTarBase: a database curates experimentally validated microRNA-target interactions. *Nucleic Acids Res*, **39**, 163-9.
- Hu H Y, He L, Fominykh K, et al (2012). Evolution of the human-specific microRNA miR-941. *Nature Communications*, **3**, 1145.
- Kinsella RJ, Kähäri A, Haider S, et al (2011). Ensembl BioMarts: a hub for data retrieval across taxonomic space. *Database*, **2011**, 30.
- Kozomara A, Griffiths-Jones S (2011). miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Res*, **39**, 152-7.
- Landthaler M, Yalcin A, Tuschl T (2004). The human DiGeorge syndrome critical region gene 8 and its *D. melanogaster* homolog are required for miRNA biogenesis. *Curr Biol*, **14**, 2162-7.
- Larkin M, Blackshields G, Brown N, et al (2007). Clustal W and Clustal X version 2.0. *Bioinformatics*, **23**, 2947-8.
- Lewis BP, Shih IH, Jones-Rhoades MW, et al (2003). Prediction of mammalian microRNA targets. *Cell*, **115**, 787-98.
- Liang T, Guo L, Liu C (2012). Genome-wide analysis of mir-548 gene family reveals evolutionary and functional implications. *J Biomed Biotechnol*, **2012**, 679563.
- Louafi F, Martinez-Nunez RT, Sanchez-Elsner T (2010). MicroRNA-155 targets SMAD2 and modulates the response of macrophages to transforming growth factor- β . *J Biol Chem*, **285**, 41328-36.
- MacRae IJ, Zhou K, Li F, et al (2006). Structural basis for double-stranded RNA processing by Dicer. *Science*, **311**, 195-8.
- Maher C, Stein L, Ware D (2006). Evolution of Arabidopsis microRNA families through duplication events. *Genome research*, **16**, 510-9.
- Marco A, Hooks K, Griffiths-Jones S (2012a). Evolution and function of the extended miR-2 microRNA family. *RNA biology*, **9**, 242-8.
- Marco A, Hui JH, Ronshaugen M, Griffiths-Jones S (2010). Functional shifts in insect microRNA evolution. *Genome Biol Evol*, **2**, 686.
- Marco A, MacPherson JJ, Ronshaugen M, Griffiths-Jones S (2012b). MicroRNAs from the same precursor have different targeting properties. *Silence*, **3**, 8-.
- Marco A, Ninova M, Ronshaugen M, Griffiths-Jones S (2013). Clusters of microRNAs emerge by new hairpins in existing transcripts. *Nucleic Acids Res*, **41**, 7745-52.
- Mattiske S, Suetani RJ, Neilsen PM, Callen DF (2012). The oncogenic role of miR-155 in breast cancer. *Cancer Epidemiol Biomarkers Prev*, **21**, 1236-43.
- Nahvi A, Shoemaker CJ, Green R (2009). An expanded seed sequence definition accounts for full regulation of the hid 3' UTR by bantam miRNA. *RNA*, **15**, 814-22.
- O'Connell RM, Chaudhuri AA, Rao DS, Baltimore D (2009). Inositol phosphatase SHIP1 is a primary target of miR-155. *Proc Natl Acad Sci USA*, **106**, 7113-8.
- O'Connell RM, Rao DS, Baltimore D (2012). microRNA regulation of inflammatory responses. *Annu Rev Immuno*, **30**, 295-312.
- Prasad TK, Goel R, Kandasamy K, et al (2009). Human protein reference database-2009 update. *Nucleic Acids res*, **37**, 767-72.
- Price N, Cartwright RA, Sabath N, et al (2011). Neutral evolution of robustness in Drosophila microRNA precursors. *Mol Biol Evol*, **28**, 2115-23.
- Pritchard CC, Cheng HH, Tewari M (2012). MicroRNA profiling: approaches and considerations. *Nat Rev Gene*, **13**, 358-69.
- Schwarz D S, Hutvagner G, Du T, et al (2003). Asymmetry in the assembly of the RNAi enzyme complex. *Cell*, **115**, 199-208.
- Smoot ME, Ono K, Ruscheinski J, et al (2011). Cytoscape 2.8: new features for data integration and network visualization. *Bioinformatics*, **27**, 431-2.
- Tamura K, Dudley J, Nei M, Kumar S (2007). MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol*, **24**, 1596-9.
- Vargova K, Curik N, Burda P, et al (2011). MYB transcriptionally regulates the miR-155 host gene in chronic lymphocytic leukemia. *Blood*, **117**, 3816-25.
- Vigorito E, Kohlhaas S, Lu D, Leyland R (2013). miR-155: an ancient regulator of the immune system. *Immunological reviews*, **253**, 146-57.
- Wang L, Toomey N L, Diaz L A, et al (2011). Oncogenic IRFs provide a survival advantage for Epstein-Barr virus-or human T-cell leukemia virus type 1-transformed cells through induction of BIC expression. *J Virol*, **85**, 8328-37.
- Wang X (2008). miRDB: a microRNA target prediction and functional annotation database with a wiki interface. *Rna*, **14**, 1012-7.
- Wheeler BM, Heimberg AM, Moy VN, et al (2009). The deep evolution of metazoan microRNAs. *Evol Dev*, **11**, 50-68.
- Wienholds E, Kloosterman WP, Miska E, et al (2005). MicroRNA expression in zebrafish embryonic development. *Science*, **309**, 310-1.
- Xiao F, Zuo Z, Cai G, et al (2009). miRecords: an integrated resource for microRNA-target interactions. *Nucleic Acids Res*, **37**, 105-10.
- Xu L, Dai WQ, Xu XF, et al (2012). Effects of multiple-target anti-microRNA antisense oligodeoxynucleotides on proliferation and migration of gastric cancer cells. *Asian Pac J Cancer Prev*, **13**, 3203-7.
- Yi R, Qin Y, Macara I G, Cullen B R (2003). Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev*, **17**, 3011-6.
- Zhang H, Kolb F A, Jaskiewicz L, et al (2004). Single processing center models for human Dicer and bacterial RNase III. *Cell*, **118**, 57-68.
- Zhang J, Haider S, Baran J, et al (2011). BioMart: a data federation framework for large collaborative projects. *Database*, **2011**, 38.
- Zheng SR, Guo GL, Zhai Q, et al (2013). Effects of miR-155 antisense oligonucleotide on breast carcinoma cell line MDA-MB-157 and implanted tumors. *Asian Pac J Cancer Prev*, **14**, 2361-6.
- Liu ZY, Zhang GL, Wang MM, Xiong YN, Cui HQ (2011). MicroRNA-663 targets TGFB1 and regulates lung cancer proliferation. *Asian Pac J Cancer Pre*, **12**, 2819-23.
- Zuker M (2003). Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res*, **31**, 3406-15.