Study of Apparent Diffusion Coefficient Changes According to Spinal Disease in MR Diffusion-weighted Image

Yeong-Cheol Heo¹ and Jae-Hwan Cho^{2*}

¹Department of Radiological Science, College of Health Sciences Eulji University, Seongnam 13135, Korea

²Department of Radiological Technology, Ansan University, Ansan 15518, Korea

(Received 1 February 2017, Received in final form 16 February 2017, Accepted 16 February 2017)

In this study, we compared the standardized value of each signal intensity, the apparent diffusion coefficient (ADC) that digitizes the diffusion of water molecules, and the signal to noise ratio (SNR) using b value 0 400, 1400 (s/mm²). From March 2013 to December 2013, patients with suspicion of simple compound fracture and metastatic spine cancer were included in the MR readout. We used a 1.5 Tesla Achieva MRI system and a Syn-Spine Coil. Sequence is a DWI SE-EPI sagittal (diffusion weighted imaging spin echo-echo planar imaging sagittal) image with b-factor (s/mm²) 0, 400, 1400 were used. Data analysis showed ROI (Region of Interest) in diseased area with high SI (signal intensity) in diffusion-weighted image b value 0 (s/mm²) Using the MRicro program, each SI was calculated with images of b-value 0, 400, and 1400 (s/mm²), ADC map was obtained using Matlab Software with each image of b-value, The ADC is obtained by applying the ROI to the same position. The standardized values (SI₄₀₀/SI₀, SI₁₄₀₀/SI₀) of simple compression fractures were 0.47 ± 0.04 and 0.23 ± 0.03 and the standardized values (SI₄₀₀/SI₀, SI₁₄₀₀/SI₀) of the metastatic spine were 0.57 ± 0.07 and 0.32 ± 0.08 And the standardized values of the two diseases were statistically significant ($p < 0.05$). The ADC (mm²/s) for b value 400 (s/mm²) and 1400 (s/mm²) of the simple compression fracture disease site were 1.70 ± 0.16 and 0.93 ± 0.28 and 1.24 ± 0.21 and 0.80 ± 0.15 for the metastatic spine. The ADC (mm²/s) for b value 400 (s/mm²) was statistically significant ($p < 0.05$) but the ADC (mm²/s) for b value 1400 ($p > 0.05$). In conclusion, multi - b value recognition of signal changes in diffusion - weighted imaging is very important for the diagnosis of various spinal diseases.

Keywords: Apparent Diffusion Coefficient (ADC), B value, Diffusion weighted imaging (DWI)

1. Introduction

Magnetic resonance imaging (MRI) has been reported to be sensitive for the diagnosis of osteoporotic complications such as compression fracture or insufficiency fracture of the spine [1]. Diffusion weighted images (DWI) were used to visualize the movement of water in the intracellular space. The skeletal muscle damage was detected early in the signal intensity changes of the muscles. Especially, relatively weak damage [2] it has been reported to be useful in the differential diagnosis of benign and malignant spinal compression fractures [3-6]. In order to obtain a diffusion-weighted image, a very strong pair of oblique magnetic fields, that is, a diffusion gradient oblique magnetic field, is used in addition to the oblique magnetic

field, which is normally used for obtaining an image. The integral of the intensity and time of the diffusion gradient gradient magnetic field is called a gradient factor or b value, and the diffusion-emphasized image can be obtained as the number of inclinations increases [7]. The apparent diffusion coefficient (ADC) is the numerical value of the degree of diffusion of water [8]. Diffusion-weighted images show early changes in muscle signal intensity when skeletal muscle damage is present and have been reported to be useful in the differential diagnosis of benign and malignant spinal compression fractures. However, is not reported much. In order to investigate the changes of simple compression fracture and metastatic spine cancer according to b value, the normalized value of each signal intensity and the diffusion degree of water molecule were measured using b value 0, 400, 1400 (s/mm²) and the signal-to-noise ratio (SNR).

©The Korean Magnetism Society. All rights reserved.

*Corresponding author: Tel: +82-31-400-6941

Fax: +82-31-363-7703, e-mail: 8452404@hanmail.net

Fig. 1. (Color online) Diffusion-weighted images of each b-value and ADC map were used to plot the ROI on the diseased area and background.

2. Subjects and Method

From March 2013 to December 2013, 22 patients were enrolled in the medical records of patients suspected of having a simple compound fracture and metastatic spine cancer. The mean age of the patients was 11 71.2, 56.6. The test equipment was a 1.5 Tesla Achieva MRI system (Philips Healthcare, Netherlands) and a Syn-Spine Coil (Philips Healthcare, Netherlands). Sequence is a DWI SE-EPI sagittal (diffusion weighted imaging spin echo-echo planar imaging sagittal) image with TR (ms) 1988, TE (ms) 68, FOV (mm) 128×128 , thickness (mm) 4, gap (mm) 0.4, Slice 13, b-factor (s/mm^2) 0, 400, 1400 were used. Data analysis showed ROI (Region of Interest) in diseased area with high SI (signal intensity) in diffusion-weighted image b value 0 (s/mm^2). Using the MRIcro program, each SI was calculated with images of b-value 0, 400, and 1400 (s/mm^2), ADC map was obtained using Matlab Software with each image of b-value, and the ADC was obtained by applying the ROI to the same position (Fig. 1). Data analysis was performed by standardizing SI of 400 and 1400 (s/mm^2) according to b value 0 for the first two diseases ($\text{SI}_{400}/\text{SI}_0$, $\text{SI}_{1400}/\text{SI}_0$) Respectively. The equation for obtaining the signal to noise ratio applied to this experiment is as follows.

$$\text{SNR} = \frac{\text{SI region, surrounding tissues}}{\text{SDN}}$$

SDN : The standard deviation of noise in background

SI : signal intensity

The difference of the mean value of the SNR and the mean diffusion coefficient obtained by each b value and the mean value of the signal - to - noise ratio and the diffusivity coefficient by disease were measured by the independent T test (SPSS win 18.0). ($P < 0.05$)

3. Result

The standardized values ($\text{SI}_{400}/\text{SI}_0$, $\text{SI}_{1400}/\text{SI}_0$) of simple compression fractures were 0.47 ± 0.04 and 0.23 ± 0.03 and the standardized values of $\text{SI}_{400}/\text{SI}_0$ and $\text{SI}_{1400}/\text{SI}_0$ of the metastatic spine were 0.57 ± 0.07 and 0.32 ± 0.08 . The mean value of standardized values was statistically significant ($p < 0.05$) (Table 1, 2). The ADC (mm^2/s) for b value 400 (s/mm^2) and 1400 (s/mm^2) of the simple compression fracture disease site were 1.70 ± 0.16 and 0.93 ± 0.28 and 1.24 ± 0.21 and 0.80 ± 0.15 for the metastatic spine.

The ADC (mm^2/s) for b value 400 (mm^2/s) was statistically significant ($p < 0.05$) but the ADC (mm^2/s) for b value 1400 ($p > 0.05$). The SNR according to the b value (0/400/1400 (mm^2/s)) of simple compression fracture and metastatic spine cancer disease was $65.61 \pm 2.61 / 36.32 \pm 2.92 / 20.22 \pm 2.16$, $48.40 \pm 2.72 / 39.27 \pm 7.51 / 23.02 \pm 6.25$ (Table 3, 4) (Fig. 2).

The SNR measurement values according to b value (0/400/1400 (s/mm^2)) are shown in Table 5.

B value of 400 s/mm^2 was 36.32 ± 2.92 for simple compression fractures and 39.27 ± 7.51 for metastatic spine cancers. B value of 1400 s/mm^2 was 20.00 ± 2.16 for simple

Table 1. Standardized signal intensity of disease site in patients with simple compression fracture.

Signal Intensity b value (s/mm^2)	SI/SI ₀ normalization value											Average
	1	2	3	4	5	6	7	8	9	10	11	
SI ₄₀₀ /SI ₀	0.47	0.5	0.52	0.53	0.46	0.44	0.53	0.44	0.42	0.4	0.46	0.47 ± 0.04
SI ₁₄₀₀ /SI ₀	0.23	0.26	0.25	0.26	0.22	0.21	0.26	0.19	0.22	0.18	0.23	0.23 ± 0.03

Table 2. Standardized signal intensity of disease site in patients with metastatic spine cancer.

SI/SI ₀ normalization value												
Signal Intensity b value (s/mm ²)	1	2	3	4	5	6	7	8	9	10	11	Average
SI ₄₀₀ /SI ₀	0.53	0.52	0.59	0.5	0.55	0.6	0.48	0.71	0.54	0.6	0.7	0.57 ± 0.07
SI ₁₄₀₀ /SI ₀	0.26	0.26	0.3	0.27	0.28	0.35	0.23	0.49	0.24	0.35	0.43	0.32 ± 0.08

Table 3. Apparent diffusion coefficient of the normal and diseased parts of simple compression fracture patients.

ADC (mm ² /s)													
b value (s/mm ²)	Part	1	2	3	4	5	6	7	8	9	10	11	Average
400	Normal part	0.68	0.73	0.65	0.88	0.42	0.38	0.39	0.59	0.55	0.68	0.45	0.58 ± 0.15
1400		0.34	0.38	0.31	0.55	0.24	0.23	0.29	0.3	0.44	0.34	0.22	0.33 ± 0.09
400	Diseased part	1.67	1.53	1.42	1.56	1.72	1.98	1.57	1.78	1.91	1.81	1.78	1.70 ± 0.16
1400		0.96	0.95	0.86	0.96	1.13	1.14	0.96	1.13	1.07	0.99	0.1	0.93 ± 0.28

Table 4. Apparent diffusion coefficient of the normal and diseased parts of metastatic spine cancer.

ADC (mm ² /s)													
b value (s/mm ²)		1	2	3	4	5	6	7	8	9	10	11	Average
400	Normal part	0.43	0.35	0.84	0.6	0.7	0.55	0.8	0.84	0.95	0.48	0.63	0.65 ± 0.18
1400		0.2	0.32	0.44	0.31	0.3	0.35	0.5	0.3	0.5	0.24	0.27	0.34 ± 0.1
400	Diseased part	1.5	1.34	1.5	1.24	0.91	1.04	1.4	1.1	1.49	1.11	0.97	1.24 ± 0.21
1400		1	0.73	0.98	0.75	0.67	0.7	1	0.7	0.95	0.7	0.57	0.80 ± 0.15

Fig. 2. In patients with simple fractures, ADC map images of b values of 400 s/mm² (a) and 1400 s/mm² (b), and ADC map images of b values 400 s/mm² (c) and 1400 s/mm² (d) in metastatic spine cancer patients.

compression fracture and 23.02 ± 6.25 for metastatic spine. In conclusion, the SNR values according to disease-specific b values were higher in metastatic spinal cancers ($p < 0.05$) (Fig. 3).

4. Discussion

In order to obtain a diffusion-weighted image, a very strong pair of oblique magnetic fields, that is, a diffusion gradient oblique magnetic field, is used in addition to the oblique magnetic field, which is normally used for obtaining an image. The integral of intensity and time of the diffusion gradient magnetic field is called a gradient factor or b value, and a diffusion-emphasized image can be obtained as the number of inclinations increases. In other words, the larger the diffusion coefficient of the tissue, and the larger the number of inclinometers used, the greater the signal degradation due to diffusion [9-13]. Generally, a b value of about 400 s/mm² can be imaged during spinal examination to image the signal loss due to diffusion. Rho [14] *et al.* measured the signal intensity in diffusion-weighted images by using b value of 150 s/mm² in patients with vertebral fractures classified into four categories such as acute osteoporosis, metastasis, chronic osteoporosis and spondylitis. On MRI, metastatic compression fractures and spondylitis have high signal intensity,

Table 5. Signal to noise ratio of simple compression fracture patients and metastatic spine cancer patients.

b value (s/mm ²)	Disease	SNR											Average
		1	2	3	4	5	6	7	8	9	10	11	
0	Simple compression fracture	68.02	69.52	62.45	67.9	68.54	64.19	62.89	66.58	65.02	65.46	61.17	65.61 ± 2.61
400		33.61	34.89	33.43	36.73	40.85	33.58	42.67	37.88	35.91	35.73	34.27	36.32 ± 2.92
1400		16.77	20.72	18.48	20.44	21.79	17.72	24.47	18.18	19.57	22.4	19.43	20.00 ± 2.16
0	Metastatic spine cancer	50.95	47.56	48.65	48.03	47.98	46.57	45.04	52.34	47.7	45.77	51.76	48.40 ± 2.27
400		34.1	49.72	40.37	33.14	49.89	35.37	29.16	42.67	28.1	41.51	47.89	39.27 ± 7.51
1400		16.68	26.12	21.49	20.86	28.36	21.69	15.31	31.29	11.92	30.57	28.92	23.02 ± 6.25

Fig. 3. In patients with simple fractures, DW images of b values of 0 s/mm² (a), 400 s/mm² (b) and 1400 s/mm² (c), and DW images of b values 0 s/mm² (d), 400 s/mm² (e) and 1400 s/mm² (f) in metastatic spine cancer patients.

acute osteoporotic compression fractures have low signal intensity, and chronic osteoporotic compression fractures show various signal intensities. However, they did not mention the difference in signal intensity in diffusion-weighted images due to changes in b Value. In this study, patients with high signal intensity among patients with chronic osteoporotic fracture based on b value of 400 s/mm² were studied. The difference of signal-to-noise ratio according to the change of b value and the signal-to- The differences were studied. DeLano *et al.* [9] reported that signal intensity decreased in white matter and granule as b value increased in normal brain. In this study, b value of 0, 400, 1400 (s/mm²) was used and there was a clear difference between the standardized value and the present diffusivity at b value 400 (s/mm²). In conclusion, it is suggested that the use of diffusely weighted diffusivity in

various spine diseases will be helpful for accurate diagnosis in addition to general spinal MRI. We could not apply various b values as a limitation of this study. We did not establish sex, age, stage and stage of each disease, and the first onset of metastatic cancer. The conclusions drawn from the data where high signal intensities are derived should be supplemented in the future.

5. Conclusion

The use of Multi - b value for the detection of signal changes in diffusion - weighted imaging is very important for the diagnosis of various spinal diseases.

References

- [1] J. S. Moon and K. C. Won, Yeungnam Univ. J. of Med. **25**, 19 (2008).
- [2] K. J. Kim, J. Korean Radiol. Soc. **42**, 175 (2000).
- [3] A. Baur, A. Stäbler, R. Brüning, R. Bartl, A. Krödel, M. Reiser, and M. Deimling, Radiology. **207**, 349 (1998).
- [4] R. Abanoz, B. Hakyemez, and M. Parlak, Tani Girisim Radyol. **9**, 176 (2003).
- [5] A. Biffar, A. Baur-Melnyk, G. P. Schmidt, M. F. Reiser, and O. Dietrich, Invest Radiol. **46**, 601 (2011).
- [6] A. Baur, A. Stabler, A. Huber, and M. Reiser, Semin Musculoskelet Radiol. **5**, 35 (2001).
- [7] J. H. Lee, J. Korean Radiol. Soc. **49**, 455 (2003).
- [8] D. Chien, K. K. Kwong, D. R. Gress, F. S. Buonanno, R. B. Buxton, and B. R. Rosen, AJNR Am. J. Neuroradiol. **13**, 1097 (1992).
- [9] M. C. DeLano, T. G. Cooper, J. E. Siebert, M. J. Potchen, and K. Kuppusamy, AJNR Am. J. Neuroradiol. **21**, 1830 (2000).
- [10] J. H. Burdette, D. D. Durden, A. D. Elster, and Y. F. Yen, J. Comput. Assist. Tomogr. **25**, 515 (2001).
- [11] Y. C. Heo, H. K. Lee, and J. H. Cho, J. Magn. **18**, 422 (2013).
- [12] Y. J. Lee, M. H. Choi, H. J. Goh, and D. K. Han, J. Magn. **21**, 281 (2013).
- [13] A. R. Cho, H. K. Lee, H. J. Yoo, and C. S. Park, J. Magn. **20**, 381 (2015).
- [14] B. H. Rho, J. Korean Radiol. Soc. **43**, 349 (2000).