ARTICLE

Detection of Antibiotic Resistance and Resistance Genes in Enterococci Isolated from Sucuk, a Traditional Turkish Dry-Fermented Sausage

Furkan Demirgöl^{1,2} and Yasin Tuncer^{1*}

¹Süleyman Demirel University, Faculty of Engineering, Department of Food Engineering, 32260, Isparta, Turkey

²Istanbul Kavram Vocational School, Department of Hotel, Restaurant and Catering Services, 34680, İstanbul, Turkey

OPEN ACCESS

Received March 8, 2017
Revised August 16, 2017
Accepted August 23, 2017

*Corresponding author

Yasin Tuncer

Süleyman Demirel University, Faculty of Engineering, Department of Food Engineering, Isparta 32260, Turkey
Tel: +90-246-211-17-13
Fax: +90-246-237-04-37
E-mail: yasintuncer@sdu.edu.tr

Copyright © Korean Society for Food Science of Animal Resources

This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

The aim of this study was to isolate enterococci in Sucuk, a traditional Turkish dry-fermented sausage and to analyze isolates for their biodiversity, antibiotic resistance patterns and the presence of some antibiotic resistance genes. A total of 60 enterococci strains were isolated from 20 sucuk samples manufactured without using a starter culture and they were identified as *E. faecium* (73.3%), *E. faecalis* (11.7%), *E. hirae* (8.3%), *E. durans* (3.3%), *E. mundtii* (1.7%) and *E. thailandicus* (1.7%). Most of the strains were found resistant to rifampin (51.67%) followed by ciprofloxacin (38.33%), nitrofurantoin (33.33%) and erythromycin (21.67%). All strains were found susceptible to ampicillin. Only *E. faecium* FYE4 and FYE60 strains displayed susceptibility to all antibiotics. Other strains showed different resistance patterns to antibiotics. *E. faecalis* was found more resistant to antibiotics than other species. Most of the strains (61.7%) displayed resistance from between two and eight antibiotics. The *ermB*, *ermC*, *gyrA*, *tetM*, *tetL* and *vanA* genes were detected in some strains. A lack of correlation between genotypic and phenotypic analysis for some strains was detected. The results of this study indicated that Sucuk manufactured without using a starter culture is a reservoir of multiple antibiotic resistant enterococci. Consequently, Sucuk is a potential reservoir for the transmission of antibiotic resistance genes from animals to humans.

Keywords *Enterococcus*, sucuk, antibiotic resistance, antibiotic resistance gene, PCR

Introduction

Fermentation is one of the oldest known food preservation techniques and has been used since ancient times to produce various fermented meat products to protect the meat. Fermented meat products are important sources of valuable nutrients such as protein, fat, essential amino acids, minerals and vitamins (Ojha *et al.*, 2015). Sucuk is a traditional Turkish dry-fermented sausage that is one of the most popular meat products produced in Turkey. In addition, it is popular in many Middle Asians, Middle Eastern, Southeastern European and Northern European countries (Ercoskun and Özkal, 2011). Sucuk is produced from a mixture of beef, sheep and/or water buffalo meats, beef fat and/or sheep tail fat, salt, sugar, nitrite/nitrate and various spices. In the traditional Sucuk production process, the prepared Sucuk

batter is stuffed into special casings, air-dried in a bovine small intestine and then fermented and ripened to develop its typical sensory characteristics (Ercoşkun and Özkal, 2011; Kaban, 2013).

Enterococcus is a large genus of lactic acid bacteria (LAB) that is important both in food and clinical microbiology (Yogurtcu and Tuncer, 2013). Some species of this genus, especially *E. faecalis* and *E. faecium*, are the relevant components of the bacterial population of some traditional cheeses (Cariolato *et al.*, 2008; Yogurtcu and Tuncer, 2013) and sausages (Landeta *et al.*, 2013; Yüceer and Özden Tuncer, 2015) produced in different European countries. On the other hand, some enterococci are considered as opportunistic human pathogens that often cause hospital-acquired infections such as endocarditis, bacteremia and urinary tract infection. Enterococcal infections are predominantly associated with *E. faecalis* and *E. faecium* (Ogier and Serror, 2008). Although antibiotic resistance is not in itself a virulence factor, multiple antibiotic resistance of enterococci is a contributing factor to their pathogenesis. Enterococci have intrinsic and acquired antibiotic resistance, which is encoded on the chromosome and plasmids or transposons, respectively (Beceiro *et al.*, 2013; Yogurtcu and Tuncer, 2013). The acquired antibiotic resistance genes can be horizontally transferred by mobile genetic elements from other strains, either distantly or closely related. In recent years, attention to the presence of antibiotic resistance in non-pathogenic bacteria has increased because they may act as reservoirs for antibiotic resistance genes (Talon and Leroy, 2011). The human gastrointestinal tract provides an ideal combination of factors for antibiotic resistance genes to arise and spread through bacterial populations (Huddleston, 2014).

Limited data are available on the antibiotic resistance of enterococci isolated from Sucuk and the presence of their antibiotic resistance genes. The objective of the present investigation was to isolate enterococci in Sucuk produced without using starter culture and to analyze isolates for their biodiversity, antibiotic resistance patterns and the presence of some antibiotic resistance genes.

Materials and Method

Sample collection and processing

A total of 20 Sucuk samples produced without using a starter culture were obtained from 20 different local manufacturers in Afyonkarahisar province, Turkey. Sucuk samples were purchased randomly between May and Septem-

ber 2013. Each 25 g Sucuk sample was homogenized with 225 mL sterile physiological water (0.85% NaCl, w/v) in a Waring blender (8011 ES HGB2WTS3, USA). 100 µL of serial decimal dilutions from homogenized Sucuk samples were inoculated onto Kanamycin Aesculin Azide (KAA) agar (LAB M Ltd., UK) and incubated at 37°C for 24-48 h. Typical enterococcal colonies were randomly selected from KAA agar and transferred into de Man, Rogosa and Sharpe (MRS) broth (LAB M). Stock cultures were stored at -20°C in MRS broth with 20% glycerol.

DNA Extraction

Total DNAs of presumptive enterococci were extracted from overnight cultures, and grown in MRS broth at 37°C, as previously described by Cancilla *et al.* (1992).

Identification of the isolates and phylogenetic analysis

The isolates were identified at the genus level, using Gram staining, catalase and cultural tests such as, growth in MRS broth at 10°C, 45°C, pH 9.6 and 6.5% NaCl (w/v). In addition, isolates were tested for resistance to heat at 60°C for 30 min (Morandi *et al.*, 2006). Identification of isolates at species level was done by 16S rDNA sequencing with pA and pE' universal bacterial primers (Edwards *et al.*, 1989). Polymerase chain reaction (PCR) was performed in 50 µL reaction mixtures, using 3 µL of DNA solution, 1 µL of each primer, 20 µL nuclease-free water and 25 µL PCR master mix (Fermentas, Lithuania). PCR was performed using the following cycling parameters: a cycle denaturation (94°C, 2 min); 30 cycles of denaturation (94°C, 30 s), annealing (55°C, 1 min) and extension (72°C, 90 s), followed by a final extension step (72°C, 10 min). The PCR products were analyzed on 1.5% (w/v) agarose gel in Tris-acetate-EDTA buffer. The gels were stained with 0.2 µg/mL of ethidium bromide (Amresco Cat no. 0492, USA) and photographed under UV light. Sequencing of the PCR products was done by Ref Gen Ltd. (Ankara University Technopolis, Turkey) using an automated gene sequencer ABI PRISM 3730XL (Perkin Elmer, USA). The sequences of the PCR products were compared to the 16S rDNA gene sequences of Genbank using the BLAST program for detection of the closest relatives. Phylogenetic analysis was conducted with MEGA software version 4.0 (MEGA; <http://www.megasoftware.net>). The tree was generated by neighbor-joining using the maximum composite likelihood model (Tamura *et al.*, 2007).

Antibiotic resistance patterns of enterococci

Antibiotic resistance patterns of isolates were detected by the disc diffusion method on Mueller-Hinton agar (Cariolato *et al.*, 2008). A total of 18 commercial antibiotic discs of ampicillin (10 µg), chloramphenicol (30 µg), ciprofloxacin (5 µg), doxycycline (30 µg), erythromycin (15 µg), gentamicin (120 µg), levofloxacin (5 µg), linezolid (30 µg), minocycline (30 µg), nitrofurantoin (300 µg), norfloxacin (10 µg), penicillin G (10 U), quinupristin/dalfopristin (15 µg), rifampin (5 µg), streptomycin (300 µg), teicoplanin (30 µg), tetracycline (30 µg) and vancomycin (30 µg) were used. All antibiotics were obtained from Oxoid Ltd. (UK). Susceptibility or resistance of enterococci was determined according to the Clinical and Laboratory Standards Institute (2012).

PCR detection of antibiotic resistance genes

The presence of chloramphenicol (*cat*), ciprofloxacin (*gyrA*), erythromycin (*ermA*, *ermB*, *ermC*), tetracycline (*tetM*, *tetL*, *tetK*, *tetS*, *tetO*) and vancomycin (*vanA*, *vanB*, *vanC*) resistance genes in enterococci was investigated by PCR. PCR primers and annealing temperatures for detec-

tion of antibiotic resistance genes are listed in Table 1. PCR was performed in 50 µL reaction mixtures. PCR conditions involved the following cycling parameters: initial denaturation cycle at 94°C for 2 min (95°C for 5 min for gene *cat*), next 30 cycles of denaturation at 94°C for 60 s (95°C for 30 s for gene *cat*), annealing at an appropriate temperature for 60 s (30 s for gene *cat*) and elongation at 72°C for 60 s (30 s for gene *cat*), and a final extension cycle at 72°C for 10 min (7 min for gene *cat*) (Dutka-Malen *et al.*, 1995; Kim *et al.*, 2005; Ouoba *et al.*, 2008; Reviriego *et al.*, 2005).

Results and Discussion

Identification of isolates

A total of 63 presumptive enterococci isolates, which appear surrounded by a black halo on KAA agar, were isolated from 20 Sucuk samples obtained from Afyonkarahisar province, Turkey. Three isolates Gram-positive rods were eliminated and the remaining 60 isolates Gram-positive cocci (single, pairs or in short chains) were selected for further analyses. All the Gram-positive cocci isolates

Table 1. PCR primers and annealing temperatures for detection of antibiotic resistance genes

Genes	Primer sequence (5' to 3')	Annealing temperature (°C)	Reference
<i>cat</i>	GCGAACGAAAAACAATTGCA TGAAGCTGTAAGGCAACTGG	55	(Kim <i>et al.</i> , 2005)
<i>gyrA</i>	GAYTATGCWATGTCAGTTATTGT GGAATRTTRGAYGTCATACCAAC	45	(Ouba <i>et al.</i> , 2008)
<i>ermA</i>	AAGCGGTAAAACCCCTCTGAG TCAAAGCCTGTCGGAATTGG	55	(Ouba <i>et al.</i> , 2008)
<i>ermB</i>	CATTTAACGACGAACTGGC GGAACATCTGTGGTATGGCG	52	(Ouba <i>et al.</i> , 2008)
<i>ermC</i>	CAAACCCGTATTCCACGATT ATCTTTGAAATCGGCTCAGG	48	(Ouba <i>et al.</i> , 2008)
<i>tetM</i>	GTAAATAGTGTTCTTGGAG CTAAGATATGGCTCTAACA	45	(Ouba <i>et al.</i> , 2008)
<i>tetL</i>	GTTGCGCGCTATATTCCAAA TTAAGCAAACCTATTCCAGC	54	(Ouba <i>et al.</i> , 2008)
<i>tetS</i>	TGGAACGCCAGAGAGGTATT ACATAGACAAGCCGTTGACC	55	(Ouba <i>et al.</i> , 2008)
<i>tetK</i>	TTAGGTGAAGGGTTAGGTCC GCAAACCTATTCCAGAAGCA	55	(Ouba <i>et al.</i> , 2008)
<i>tetO</i>	GATGGCATAACAGGCACAGAC CAATATCACCAGAGCAGGCT	55	(Ouba <i>et al.</i> , 2008)
<i>vanA</i>	GGGAAAACGACAATTGC GTACAATGCGGCCGTTA	54	(Dutka-Malen <i>et al.</i> , 1995)
<i>vanB</i>	GTGCTGCGAGATACCACAGA CGAACACCATGCAACATTTC	54	(Reviriego <i>et al.</i> , 2005)
<i>vanC</i>	GGTATCAAGGAAACCTC CTTCCGCCATCATAGCT	54	(Dutka-Malen <i>et al.</i> , 1995)

Y, C or T; R, A or G; W, A or T.

Fig. 1. Phylogenetic analysis of 16S rRNA gene sequences of *Enterococcus* strains.

were catalase-negative and resistant to heat at 60°C for 30 min. In addition, they were grown in MRS broth at 10°C, 45°C, pH 9.6 and presence of 6.5% NaCl. The results of Gram staining, catalase and cultural tests showed that these 60 strains isolated from Sucuk samples were mem-

bers of the *Enterococcus* genus. 60 presumptive *Enterococcus* strains were identified at species level by 16S rDNA sequence analyses. Presumptive *Enterococcus* strains were identified as 44 *E. faecium* (73.33%), 7 *E. faecalis* (11.67%), 5 *E. hirae* (8.33%), 2 *E. durans* (3.33%), 1 *E. mundtii*

(1.67%) and 1 *E. thailandicus* (1.67%). Phylogenetic analysis of 16S rRNA gene sequences of *Enterococcus* strains was shown in Fig. 1. Enterococci are associated with some traditional sausages produced in different European countries (Jahan *et al.*, 2013; Landeta *et al.*, 2013; Yüceer and Özden Tuncer, 2015), as confirmed in this study. Dominant *Enterococcus* species in Sucuk samples were determined as *E. faecium* (73.33%). Similar to our results, most previous studies reported *E. faecium* as a dominant microbiota in fermented sausages (Landeta *et al.*, 2013; Valenzuela *et al.*, 2009; Yüceer and Özden Tuncer, 2015). In contrast, some researchers found a higher percentage of *E. faecalis* than other species in animal originated foods (Jahan *et al.*, 2013; Peters *et al.*, 2003).

Antibiotic resistance patterns of *Enterococcus* strains

Sixty *Enterococcus* strains were tested for their resistance to 18 different antibiotics using the disc diffusion method.

Antibiotic resistance test results of *Enterococcus* strains were given in Table 2. Only *E. faecium* FYE4 and FYE60 strains displayed susceptibility to all antibiotics. All of the strains were found sensitive only to ampicillin. Similar to our result, some researchers showed that enterococci isolated from different traditional fermented sausages and cheeses were completely sensitive to ampicillin (Jahan *et al.*, 2013; Landeta *et al.*, 2013; Yogurtcu and Tuncer, 2013; Yüceer and Özden Tuncer, 2015). On the other hand, Chajęcka-Wierzchowska *et al.* (2012) showed that low numbers of enterococci isolated from animal originated food were resistant to ampicillin. *Enterococcus* strains showed different susceptibility patterns to the other antibiotics used in this study. Antibiotic susceptibility and resistance percentage of *Enterococcus* strains were given in Table 3.

Most of the strains were found sensitive to gentamicin (93.33%), streptomycin (93.33%), chloramphenicol (90.00%), minocycline (90.00%), penicillin G (90.00%),

Table 2. Antibiotic resistance of *Enterococcus* strains isolated from Sucuk and positive PCR for resistance genes in these strains

Strains	Antibiotic susceptibility ⁽¹⁾																		Positive PCR/ resistance genes
	AMP	DO	E	CN	C	LEV	LZD	QD	MH	F	NOR	P	RD	CIP	S	TEC	TE	VA	
<i>E. faecalis</i> FYE1	S ²⁾	I	R	R	R	S	S	R	I	S	I	S	S	R	R	I	I	S	<i>ermB</i> , <i>tetM</i>
<i>E. faecium</i> FYE2	S	S	I	S	S	I	S	S	S	R	I	S	R	R	S	I	S	S	<i>ermC</i>
<i>E. faecalis</i> FYE3	S	I	R	R	R	I	I	R	R	S	I	S	S	R	R	R	I	S	<i>ermB</i> , <i>tetM</i>
<i>E. faecium</i> FYE4	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	<i>ermB</i>
<i>E. faecium</i> FYE5	S	S	R	S	S	R	I	S	S	R	R	S	R	R	S	I	S	S	<i>ermC</i> , <i>gyrA</i>
<i>E. faecium</i> FYE6	S	S	I	S	S	S	S	S	S	S	S	S	S	I	S	S	S	S	none
<i>E. faecium</i> FYE7	S	S	R	S	S	S	S	S	S	S	R	S	S	S	S	I	S	S	none
<i>E. faecium</i> FYE8	S	S	I	S	S	S	S	S	S	S	S	S	S	I	S	I	S	S	<i>ermC</i>
<i>E. faecium</i> FYE9	S	S	S	S	S	S	S	S	S	I	S	S	S	I	S	I	S	S	none
<i>E. faecium</i> FYE10	S	S	I	S	S	I	S	S	S	S	I	S	S	I	S	I	S	S	none
<i>E. faecalis</i> FYE11	S	S	I	S	S	I	I	S	S	R	I	S	R	R	S	I	S	S	<i>ermC</i>
<i>E. faecium</i> FYE12	S	S	I	S	S	I	I	S	S	R	I	S	R	R	S	I	S	S	none
<i>E. faecium</i> FYE13	S	S	I	S	S	S	S	S	S	R	S	S	S	S	S	I	S	S	none
<i>E. faecalis</i> FYE14	S	R	R	S	R	I	I	R	R	S	I	S	R	I	R	I	R	I	<i>ermB</i> , <i>tetL</i>
<i>E. faecium</i> FYE15	S	S	I	S	S	I	S	S	S	R	I	S	R	I	S	I	S	S	<i>tetM</i>
<i>E. faecium</i> FYE16	S	S	S	S	S	R	S	I	S	R	S	S	S	R	S	R	S	I	<i>ermC</i> , <i>vanA</i>
<i>E. thailandicus</i> FYE17	S	S	R	S	S	I	I	I	S	S	I	S	R	I	S	I	S	S	<i>gyrA</i>
<i>E. faecium</i> FYE18	S	S	I	S	S	I	I	I	S	R	I	S	R	I	S	I	S	I	none
<i>E. hirae</i> FYE19	S	S	S	S	S	S	S	I	S	I	S	S	I	I	S	S	S	S	<i>gyrA</i>
<i>E. faecium</i> FYE20	S	S	I	S	S	S	I	S	S	R	R	S	R	R	S	I	S	S	<i>ermC</i>

Table 2. Antibiotic resistance of *Enterococcus* strains isolated from Sucuk and positive PCR for resistance genes in these strains (Continued)

Strains	Antibiotic susceptibility ¹⁾																		Positive PCR/ resistance genes
	AMP	DO	E	CN	C	LEV	LZD	QD	MH	F	NOR	P	RD	CIP	S	TEC	TE	VA	
<i>E. faecium</i> FYE21	S	S	I	S	S	I	S	S	S	R	R	R	R	R	S	I	S	S	none
<i>E. faecium</i> FYE22	S	S	I	S	S	I	I	S	S	R	I	R	R	R	S	I	S	S	ermC
<i>E. faecium</i> FYE23	S	S	I	S	S	I	S	S	S	R	I	R	R	R	S	I	S	S	ermC
<i>E. faecium</i> FYE24	S	S	I	S	S	S	S	S	S	R	I	S	R	I	S	I	S	S	ermC
<i>E. mundtii</i> FYE25	S	S	I	S	S	S	S	S	S	S	S	S	S	I	S	I	S	S	none
<i>E. faecium</i> FYE26	S	S	I	S	S	I	S	S	S	I	I	S	R	R	S	I	S	S	none
<i>E. faecalis</i> FYE27	S	I	R	I	I	S	I	R	I	S	I	S	I	I	S	I	I	I	ermB
<i>E. faecium</i> FYE28	S	S	I	S	S	I	I	I	S	R	I	R	S	R	S	R	S	S	ermC
<i>E. faecalis</i> FYE29	S	S	I	S	S	I	R	R	S	S	I	S	R	I	S	R	S	R	none
<i>E. hirae</i> FYE30	S	S	S	S	S	S	S	I	S	R	S	S	I	I	S	I	S	S	gyrA
<i>E. hirae</i> FYE31	S	S	S	S	S	S	S	I	S	R	S	S	S	I	S	I	S	S	gyrA
<i>E. faecium</i> FYE32	S	R	S	S	S	S	S	S	S	S	S	S	R	S	S	S	R	S	tetL
<i>E. faecium</i> FYE33	S	S	I	S	S	S	I	I	S	S	I	S	S	I	S	I	S	I	tetM, gyrA
<i>E. durans</i> FYE34	S	S	I	S	S	I	I	I	S	S	I	R	R	I	S	I	S	S	none
<i>E. faecium</i> FYE35	S	S	I	S	S	R	I	S	S	I	R	S	S	R	S	I	S	I	none
<i>E. hirae</i> FYE36	S	S	I	S	S	S	S	S	S	S	S	S	I	S	S	I	S	S	gyrA
<i>E. faecium</i> FYE37	S	I	R	S	R	I	I	I	I	S	R	S	S	R	S	I	R	S	ermB, tetL
<i>E. faecium</i> FYE38	S	S	I	S	S	S	S	S	S	S	S	S	R	S	S	I	S	S	tetM
<i>E. faecium</i> FYE39	S	S	I	S	S	I	S	S	S	I	S	S	I	I	S	I	S	S	none
<i>E. faecium</i> FYE40	S	S	I	S	S	S	S	S	S	R	I	S	R	I	S	S	S	S	ermC
<i>E. faecium</i> FYE41	S	S	R	S	S	S	S	S	S	S	S	S	R	S	S	I	S	S	ermC, tetM, tetL
<i>E. faecium</i> FYE42	S	S	I	S	S	I	S	S	S	S	S	S	R	I	S	I	S	S	none
<i>E. faecium</i> FYE43	S	S	I	S	S	I	I	S	S	S	I	S	R	R	S	I	S	S	none
<i>E. durans</i> FYE44	S	S	R	S	S	I	I	I	S	R	S	S	R	I	S	R	S	S	gyrA
<i>E. faecium</i> FYE45	S	S	R	S	S	I	S	S	S	S	I	S	R	R	S	I	S	S	none
<i>E. faecium</i> FYE46	S	S	R	S	S	R	S	S	S	R	I	R	I	R	S	I	S	S	ermC
<i>E. faecium</i> FYE47	S	S	I	S	S	S	S	I	S	S	S	S	S	I	S	I	S	S	none
<i>E. faecium</i> FYE48	S	S	I	S	S	S	S	I	S	S	S	S	S	I	S	I	S	S	none
<i>E. faecium</i> FYE49	S	S	I	S	S	S	S	I	S	S	S	S	S	I	S	S	S	S	none
<i>E. faecium</i> FYE50	S	S	I	S	S	S	S	I	S	S	S	S	S	I	S	I	S	S	none
<i>E. faecalis</i> FYE51	S	I	R	R	R	S	S	R	R	S	I	S	I	I	R	I	I	I	ermB, tetM
<i>E. faecium</i> FYE52	S	S	I	S	S	I	S	S	S	I	I	S	R	R	S	S	S	S	none
<i>E. faecium</i> FYE53	S	S	I	S	S	S	S	S	S	I	I	S	R	R	S	S	S	S	none
<i>E. faecium</i> FYE54	S	S	I	S	S	S	S	S	S	S	I	S	R	I	S	S	S	S	none
<i>E. faecium</i> FYE55	S	S	I	S	S	I	S	S	S	I	I	S	R	R	S	S	S	S	none
<i>E. faecium</i> FYE56	S	S	I	S	S	I	S	S	S	S	S	S	R	I	S	S	S	S	none
<i>E. faecium</i> FYE57	S	S	I	S	S	S	S	S	S	I	I	S	R	R	S	S	S	S	none
<i>E. faecium</i> FYE58	S	S	I	S	S	S	S	S	S	I	I	S	R	R	S	S	S	S	none
<i>E. hirae</i> FYE59	S	S	S	S	S	S	S	I	S	R	S	S	I	I	S	I	S	S	gyrA
<i>E. faecium</i> FYE60	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	none

¹⁾AMP, Ampicillin (10 µg); C, Chloramphenicol (30 µg); CIP, Ciprofloxacin (5 µg); CN, Gentamicin (120 µg); DO, Doxycycline (30 µg); E, Erythromycin (15 µg); F, Nitrofurantoin (300 µg); LEV, Levofloxacin (5 µg); LZD, Linezolid (30 µg); MH, Minocycline (30 µg); NOR, Norfloxacin (10 µg); P, Penicillin G (10 U); QD, Quinupristin/dalfopristin (15 µg); RD, Rifampin (5 µg); S, Streptomycin (300 µg); TEC, Teicoplanin (30 µg); TE, Tetracycline (30 µg); VA, Vancomycin (30 µg).

²⁾S, susceptible; I, intermediary; R, resistance.

Table 3. Antibiotic susceptibility and resistance percentage of *Enterococcus* strains

Antibiotics	Concentration (µg/disc)	Sensitive (%)	Intermediary (%)	Resistance (%)
Ampicillin	10	100	0	0
Chloramphenicol	30	90	1.67	8.33
Ciprofloxacin	5	13.33	48.33	38.33
Doxycycline	30	88.33	8.33	3.33
Erythromycin	15	15	63.33	21.67
Gentamicin	120	93.33	1.67	5
Levofloxacin	5	51.67	41.67	6.67
Linezolid	30	70	28.33	1.67
Minocycline	30	90	5	5
Nitrofurantoin	300	50	16.67	33.33
Norfloxacin	10	40	50	10
Penicillin G	10 ¹⁾	90	0	10
Quinupristin/dalfopristin	15	63.33	26.67	10
Rifampin	5	35	13.33	51.67
Streptomycin	300	93.33	0	6.67
Teicoplanin	30	23.33	68.33	8.33
Tetracycline	30	88.33	6.67	5
Vancomycin	30	86.67	11.67	1.67

¹⁾Penicillin G 10 U/disc.

doxycycline (88.33%), tetracycline (88.33%) and vancomycin (86.67%), followed by other antibiotics. On the contrary, *Enterococcus* strains were found resistant to rifampin (51.67%), ciprofloxacin (38.33%), nitrofurantoin (33.33%) and erythromycin (21.67%), followed by other antibiotics except for ampicillin. Recently, similar observations to our results have been reported by several researchers. Peters *et al.* (2003) indicated that 22.03% of the *Enterococcus* strains isolated from animal originated foods were sensitive to erythromycin. Chajęcka-Wierżchowska *et al.* (2012) reported that enterococci from food of animal origin were sensitive to chloramphenicol (96.73%), streptomycin (91.30%), gentamicin (85.86%) and quinupristin/dalfopristin (66.66%). In addition, Yüceer and Özden Tuncer (2015) showed that *Enterococcus* strains from Sucuk were sensitive to chloramphenicol (100%), doxycycline (100%), gentamicin (100%), minocycline (100%), streptomycin (100%), tetracycline (100%), penicillin G (88%), levofloxacin (52%), rifampin (28%) and ciprofloxacin (12%). Conversely to our results, some researchers showed that *Enterococcus* strains isolated from animal originated foods were 100% sensitive to levofloxacin, quinupristin/dalfopristin, teicoplanin or vancomycin (Chajęcka-Wierżchowska *et al.*, 2012; Landeta *et al.*, 2013; Peters *et al.*, 2003; Yogurtcu and Tuncer, 2013; Yüceer and Özden Tuncer, 2015).

Antibiotic susceptibility and resistance percentage of enterococci at species level was shown in Table 4. Among

the *Enterococcus* species, *E. faecalis* showed the highest percentage of resistance to the tested antibiotics. *E. faecalis* strains displayed resistance to quinupristin/dalfopristin (85.71%), erythromycin (71.43%), chloramphenicol (57.14%) and streptomycin (57.14%). On the other hand, *E. faecium* strains were found resistant to rifampin (56.82%), ciprofloxacin (45.45%) and nitrofurantoin (34.09%). *E. hirae* strains showed resistance only to nitrofurantoin. Three out of five *E. hirae* strains were found resistant to nitrofurantoin. Both of the *E. durans* strains isolated from Sucuk exhibited resistance to rifampin. In addition, *E. durans* FYE44 was found resistant to erythromycin, nitrofurantoin, penicillin G and teicoplanin. *E. mundtii* FYE25 did not display resistance to any antibiotics used in this study. *E. thailandicus* FYE17 showed resistance to erythromycin and rifampin. Similar to our results, Yogurtcu and Tuncer (2013) reported that *E. faecalis* strains isolated from Turkish tulum cheese were found more resistant to antibiotics than other species. On the other hand, Chajęcka-Wierżchowska *et al.* (2012) determined that *E. faecalis* and *E. faecium* strains isolated from ready-to-eat food of animal origin showed similar antibiotic resistance patterns. Conversely, some researchers reported that *E. faecium* strains were more resistant to antibiotics than other species (Landeta *et al.*, 2013; Valenzuela *et al.*, 2009).

The *Enterococcus* strains (61.67%) displayed resistance from between two and eight antibiotics. Some of the strains (11.67%) showed resistance to ciprofloxacin and rifampin.

Table 4. Antibiotic susceptibility and resistance per cent of *E. faecium*, *E. faecalis*, *E. hirae*, *E. durans*, *E. mundtii* and *E. thailandicus* strains isolated from Sucuk

Antibiotics	<i>E. faecium</i> (n ¹ : 44)			<i>E. faecalis</i> (n: 7)			<i>E. hirae</i> (n: 5)			<i>E. durans</i> (n: 2)			<i>E. mundtii</i> (n: 1)			<i>E. thailandicus</i> (n: 1)		
	S ²	I	R	S	I	R	S	I	R	S	I	R	S	I	R	S	I	R
Ampicillin	100	0	0	100	0	0	100	0	0	100	0	0	100	0	0	100	0	0
Chloramphenicol	97.73	0	2.27	28.57	14.29	57.14	100	0	0	100	0	0	100	0	0	100	0	0
Ciprofloxacin	15.91	38.64	45.45	0	57.14	42.86	20	80	0	0	100	0	0	100	0	0	100	0
Doxycycline	95.46	2.27	2.27	28.57	57.14	14.29	100	0	0	100	0	0	100	0	0	100	0	0
Erythromycin	11.36	75	13.64	0	28.57	71.43	80	20	0	0	50	50	0	100	0	0	0	100
Gentamicin	100	0	0	42.86	14.29	42.86	100	0	0	100	0	0	100	0	0	100	0	0
Levofloxacin	50	40.91	9.09	42.86	57.14	0	100	0	0	0	100	0	100	0	0	0	100	0
Linezolid	77.27	22.73	0	28.57	57.14	14.29	100	0	0	0	100	0	100	0	0	0	100	0
Minocycline	97.73	2.27	0	28.57	28.57	42.86	100	0	0	100	0	0	100	0	0	100	0	0
Nitrofurantoin	45.45	20.45	34.09	85.71	0	14.29	20	20	60	50	0	50	100	0	0	100	0	0
Norfloxacin	38.64	47.73	13.64	0	100	0	100	0	0	50	50	0	100	0	0	0	100	0
Penicillin G	88.64	0	11.36	100	0	0	100	0	0	50	0	50	100	0	0	100	0	0
Quinupristin/dalfopristin	79.55	20.45	0	14.29	0	85.71	20	80	0	0	100	0	100	0	0	0	100	0
Rifampin	38.64	4.55	56.82	28.57	28.57	42.86	20	80	0	0	0	100	100	0	0	0	0	100
Streptomycin	100	0	0	42.86	0	57.14	100	0	0	100	0	0	100	0	0	100	0	0
Teicoplanin	29.55	65.91	4.55	0	71.43	28.57	20	80	0	0	50	50	0	100	0	0	100	0
Tetracycline	95.45	0	4.55	28.57	57.14	14.29	100	0	0	100	0	0	100	0	0	100	0	0
Vancomycin	90.91	9.09	0	42.86	42.86	14.29	100	0	0	100	0	0	100	0	0	100	0	0

¹n, number of strain.²S, susceptible; I, intermediary; R, resistance.

Multiple antibiotic resistance in enterococci is not surprising. Several researchers reported that multiple antibiotic resistance in enterococci isolated from fermented meat products is common (Jahan *et al.*, 2013; Yüceer and Özden Tuncer, 2015), as confirmed in this study. Jahan *et al.* (2013) showed that *Enterococcus* strains isolated from fermented meat products exhibited a high rate (93.10%) of multiple antibiotic resistance characteristics. Researchers also reported that 58.62% of the strains displayed resistance from between three and eight antibiotics. In addition, Yüceer and Özden Tuncer (2015) determined that 68% of the *Enterococcus* strains isolated from fermented Sucuk exhibited multiple antibiotic resistance. The strains displayed resistance from between two and five antibiotics.

PCR detection of antibiotic resistance genes

A total of 60 *Enterococcus* strains were evaluated for the presence of chloramphenicol (*cat*), ciprofloxacin (*gyrA*), erythromycin (*ermA*, *ermB*, *ermC*), tetracycline (*tetM*, *tetL*, *tetK*, *tetS*, *tetO*) and vancomycin (*vanA*, *vanB*, *vanC*) resistance genes. Thirty-one out of 60 isolates showed positive PCR for *gyrA*, *ermB*, *ermC*, *tetM*, *tetL* and *vanA*. A lack of correlation between genotypic and phenotypic analysis was detected (Table 2).

A correlation between the chloramphenicol resistance

phenotype and genotype was not detected in the researched strains. The *cat* gene was not detected by PCR in five phenotypically chloramphenicol-resistant strains. This result indicated that other chloramphenicol resistance genes may be influenced in these strains. From this, we presumed that positive PCR results for *cat* gene were not detected in these strains by the primers used in this study. Similar to our results, Kim *et al.* (2005) determined that 17 out of 29 chloramphenicol-resistant *Staphylococcus intermedius* isolates contained the *cat* gene, while 12 isolates did not carry the *cat* gene by the same primer set used in our study. In addition, Jahan *et al.* (2013) reported that only *E. faecium* S15 strain was positive for the *cat* gene among three phenotypically chloramphenicol-resistant enterococci isolated from meat and fermented meat products.

The *gyrA* gene was detected in nine *Enterococcus* strains that are phenotypically one resistant, seven intermediate and one susceptible). Only one strain, *E. faecium* FYE5, was given a positive result for *gyrA* gene among 23 phenotypically ciprofloxacin-resistant *Enterococcus* strains. Similar to our results, Jahan *et al.* (2013) did not detect *gyrA* or *parC* gene in any of three ciprofloxacin-resistant *Enterococcus* strains isolated from meat and fermented meat products. In addition, PCR results showed that phenotypically ciprofloxacin-susceptible *E. hirae* FYE36 strain

contains the *gyrA* gene (Table 2). In Gram positive bacteria, both *gyrA* and *parC* genes are defined to be the primary target in quinolone resistance. Mutations in Ser84 locus of GyrA in *Staphylococcus aureus* resulted in increased of ciprofloxacin-resistance. Similarly, single mutations taking place in the Ser83 and the Glu87 loci of GyrA were related to an increase of minimal inhibitory concentration values of ciprofloxacin in *E. faecalis* strains. The mutations taking place in both *gyrA* and *parC* genes were found in high-level quinolone-resistant *E. faecium* (Petersen *et al.*, 2004).

The erythromycin resistance gene *ermA* was not detected in any strains. However, *ermB* and *ermC* genes were detected in seven (11.67%) and 13 (21.67%) strains, respectively. These strains were showed the expected amplification fragments of 425 bp (Fig. 2) and 295 bp when their DNAs were amplified with specific primers for *ermB* and *ermC* genes, respectively. Conversely to our results, previous studies showed that the *ermB* gene is the most common erythromycin-resistant gene defined in *Enterococcus* species (Aarestrup *et al.*, 2000; Jahan *et al.*, 2013). In our study, the *ermC* gene was found at a higher frequency than the *ermB* gene. A lack of correlation between phenotypic and genotypic analysis for erythromycin resistance was detected (Table 2). The *ermB* gene was detected in six erythromycin-resistant and one erythromycin-susceptible *Enterococcus* strains. In addition, the *ermC* gene was found in nine erythromycin-intermediate, two erythromycin-resistant and one erythromycin-susceptible strains. The erythromycin-resistant strains *E. faecium* FYE7, *E. thailandicus* FYE17, *E. durans* FYE44 and *E.*

faecium FYE45 did not present any of the three *erm* genes (Table 2), indicating that other factors influence resistance in these strains. Similar to these results, Jahan *et al.* (2013) demonstrated the presence of *ermB* and/or *mefA/B* genes in seven of 13 phenotypically erythromycin-resistant *Enterococcus* strains isolated from meat and fermented meat products. However, *erm* genes (*ermA*, *ermB*, *ermC* or *mefA/B*) were not found in the remaining six *Enterococcus* strains. In our study, it is interesting that the presence of *ermB* and *ermC* genes was determined in phenotypically erythromycin-susceptible *E. faecium* FYE4 and *E. faecium* FYE16 strains, respectively. A similar observation was recently reported by Ding *et al.* (2012). In this study, researchers determined the presence of *ermB* and *ermB/ermC* genes in six and two erythromycin-susceptible *Staphylococcus* strains, respectively. The researchers showed that the mutation of 23S *rRNA* gives rise to many erythromycin-susceptible isolates when it has *erm* genes, the mutation on 23S *rRNA* possibly disturbing the site of the methylation, allowing erythromycin to bind to the ribosome. The reason for lack of correlation between genotypic and phenotypic analysis in *E. faecium* FYE4 and *E. faecium* FYE16 strains may be due to mutation of the 23S *rRNA* gene in these strains. Further investigations are needed for the determination of erythromycin susceptibility mechanisms in these strains.

The tetracycline resistance genes *tetM* and *tetL* were detected in seven (11.67%) and four (6.67%) strains, respectively. In tetracycline-resistant *E. faecium* FYE32 and *E. faecium* FYE37 strains carried the *tetL* gene and *E. faecalis* FYE14 strain carried both *tetM* and *tetL* genes. More-

Fig. 2. PCR screen for *ermB* gene from *Enterococcus* strains. Order line 1, *E. faecalis* FYE1; line 2, *E. faecium* FYE2; line 3, *E. faecalis* FYE3; line 4, *E. faecium* FYE4; line 5, *E. faecium* FYE5; line 6, *E. faecium* FYE6; line 7, *E. faecium* FYE7; line 8, *E. faecium* FYE8; line 9, *E. faecium* FYE9; line 10, *E. faecium* FYE10; line M, 100 bp DNA ladder (Fermentas); line 11, *E. faecalis* FYE11; line 12, *E. faecium* FYE12; line 13, *E. faecium* FYE13; line 14, *E. faecalis* FYE14; line 15, *E. faecium* FYE15; line 16, *E. faecium* FYE16; line 17, *E. thailandicus* FYE17; line 18, *E. faecium* FYE18; line 19, negative control.

over, *tetM* gene was detected in three out of four tetracycline-intermediate strains. The *tetK*, *tetS* and *tetO* genes were not detected in any strains. Different researchers have reported that the *tetM* gene is more common in tetracycline-resistant *Enterococcus* strains (Aarestrup *et al.*, 2000; Cauwerts *et al.*, 2007; Jahan *et al.*, 2013), as confirmed in this study. Similar to our results, Aarestrup *et al.* (2000) detected *tetM* and *tetL* genes in tetracycline-resistant *E. faecalis* and *E. faecium* strains isolated from humans, broilers and pigs. Researchers did not detect *tetK* and *tetS* genes in any animal-origin isolates. Hummel *et al.* (2008) reported that 94%, 63% and 56% of tetracycline-resistant *Enterococcus* strains isolated from cheese contained *tetL*, *tetM* and *tetK* genes, respectively. In addition, researchers reported that none of the strains contained *tetO* or *tetS* genes, as confirmed in this study. In another study, Jahan *et al.* (2013) also detected *tetM* (16 strains), *tetL* (six strains), and *tetK* (six strains), genes in 19 *Enterococcus* strains isolated from meat and fermented meat products, respectively. The *tetO* and *tetS* genes were not found in any of the strains. A lack of correlation between phenotypic and genotypic analysis for tetracycline resistance was detected (Table 2). In our study, the presence of both *tetM* and *tetL* genes was determined in the phenotypically tetracycline-susceptible *E. faecium* FYE41 strain. Similar observations were detected by Cauwerts *et al.* (2007). Researchers determined that two strains contained *tetM* and one strain contained *tetL* among three phenotypically tetracycline-sensitive *E. faecium* strains isolated from broilers.

The vancomycin resistance genes *vanB* and *vanC* were not detected in any tested strains. However, *vanA* gene was detected only in vancomycin-intermediate *E. faecium* FYE16 strain. One vancomycin-resistant and six vancomycin-intermediate *Enterococcus* strains did not present any of the three vancomycin resistance genes (Table 2). The *vanA* is the most common glycopeptide resistance gene in enterococci and it is usually related to high-level vancomycin resistance. Most of the VanA type resistant strains are also teicoplanin resistant (Garrido *et al.*, 2014). In this study, the phenotypically vancomycin-resistant strain, *E. faecalis* FYE29, was also found to be teicoplanin-resistant but, none of the vancomycin resistance genes researched in this study were found, indicating that other factors influence resistance in this strain. Baylan *et al.* (2011) reported that one out of eight phenotypically vancomycin- and teicoplanin-resistant urinary *E. faecium* isolates did not contain *vanA* or *vanB* genes, as confirmed in this study. The *vanA*, *vanB* and *vanC* genes were det-

ected in none of the phenotypically vancomycin-sensitive *Enterococcus* strains as expected. Conversely to our findings, Szakacs *et al.* (2014) determined the presence of the *vanA* gene in clinical isolate of vancomycin-sensitive *E. faecium*.

Conclusion

The results of this study indicated that *E. faecium* is the dominant *Enterococcus* species present in Sucuk. *Enterococcus* strains were found resistant to the clinically relevant antibiotics except ampicillin. Most of *Enterococcus* strains displayed multiple antibiotic resistance. The *ermB*, *ermC*, *gyrA*, *tetM*, *tetL* and *vanA* genes were detected in some strains. These strains may play a role in the spread of antibiotic resistance among bacteria, and so could create a health risk for consumers. The results of this study indicated that Sucuk manufactured without using a starter culture is a reservoir of multiple antibiotic resistant enterococci. Consequently, Sucuk is a potential reservoir for the transmission of antibiotic resistance genes from animals to humans. Other potential risk factors (biogenic amine production and virulence factors) of enterococci isolated from Sucuk should be researched to protect consumer health. Moreover, further investigations are also needed for the determination of the lack of correlation between antibiotic resistance genotypes and phenotypes in enterococci to clarify antibiotic resistance and susceptibility mechanisms of these bacteria.

Acknowledgements

This study was financially supported by the Süleyman Demirel University Scientific Research Project Coordination Unit through project number 3284-YL2-12. Part of this study was presented as a poster presentation at the 2nd International Congress on Food Technology in Turkey, and the abstract was published in the Congress's book of abstracts.

References

- Aarestrup, F. M., Agerso, Y., Gerner-Smidt, P., Madsen, M., and Jensen, L. B. (2000) Comparison of antimicrobial resistance phenotypes and resistance genes in *Enterococcus faecalis* and *Enterococcus faecium* from humans in the community, broilers, and pigs in Denmark. *Diagn. Microbiol. Infect. Dis.* **37**, 127-137.
- Baylan, O., Nazik, H., Bektöre, B., Çitil, B. E., Turan, D., Öngen,

- B., Özyurt, M., Açikel, C. H., and Haznedaroğlu, T. (2011) The relationship between antibiotic resistance and virulence factors in urinary *Enterococcus* isolates (in Turkish). *Mikrobiyol Bul.* **45**, 430-445.
- Beceiro, A., Tomás, M., and Bou, G. (2013) Antimicrobial resistance and virulence: A successful or deleterious association in the bacterial world? *Clin. Microbial. Rev.* **26**, 185-230.
- Cancilla, M. R., Powell, L.B., Hillier, A. J., and Davidson, B. E. (1992) Rapid genomic fingerprinting of *Lactococcus lastis* strains by arbitrarily primed polymerase chain reaction with 32P and fluorescent labels. *Appl. Environ. Microbiol.* **58**, 1772-1775.
- Cariolato, D., Andrighetto, C., and Lombardi, A. (2008) Occurrence of virulence factors and antibiotic resistances in *Enterococcus faecalis* and *Enterococcus faecium* collected from dairy and human samples in North Italy. *Food Control* **19**, 886-892.
- Cauwerts, K., Decostere, A., De Graef, E. M., Haesebrouck, F., and Pasmans, F. (2007) High prevalence of tetracycline resistance in *Enterococcus* isolates from broilers carrying the *erm* (B) gene. *Avian Pathol.* **36**, 395-399.
- Chajęcka-Wierzchowska, W., Zadernowska, A., Nalepa, B., and Laniewska-Trokenheim, L. (2012) Occurrence and antibiotic resistance of enterococci in ready-to-eat food of animal origin. *Afr. J. Microbiol. Res.* **6**, 6773-6780.
- CLSI. (2012) Performance Standards for Antimicrobial Susceptibility Testing; 22th Informational Supplement M100-S-22 NCCLS. Pennsylvania: Wayne, United States of America.
- Ding, Z. F., Zhang, H., Tang, W., Tong, C. Y., Li, R. T., Chen, L. X., Pu, L. J., Zhu, Z. B., and Cui, Y. D. (2012) Methylase genes-mediated erythromycin resistance in *Staphylococcus aureus* from bovine mastitis in China. *Isr. J. Vet. Med.* **67**, 170-179.
- Dutka-Malen, S., Evers, S., and Courvalin, P. (1995) Detection of glycopeptide resistance genotypes and identification to the species level of clinically relevant enterococci by PCR. *J. Clin. Microbiol.* **33**, 24-27.
- Edwards, U., Rogall, T., Blocker, H., Emde, M., and Bottger, E. C. (1989) Isolation and direct complete nucleotide determination of entire genes. Characterization of a gene coding for 16S ribosomal RNA. *Nucleic Acids Res.* **17**, 7843-7853.
- Ercoşkun, H. and Özkal, S. G. (2011) Kinetics of traditional Turkish sausage quality aspects during fermentation. *Food Control* **22**, 165-172.
- Garrido, A. M., Gálvez, A., and Pulido, R. P. (2014) Antimicrobial resistance in enterococci. *J. Infect. Dis. Ther.* **2**, 1-7.
- Huddleston, J. R. (2014) Horizontal gene transfer in the human gastrointestinal tract: Potential spread of antibiotic resistance genes. *Infect. Drug Resist.* **7**, 167-176.
- Hummel, A., Holzapfel, W. H., and Franz, C. M. A. P. (2008) Characterisation and transfer of antibiotic resistance genes from enterococci isolated from food. *Syst. Appl. Microbiol.* **30**, 1-7.
- Jahan, M., Krause, D. O., and Holley, R. A. (2013) Antimicrobial resistance of *Enterococcus* species from meat and fermented meat products isolated by a PCR-based rapid screening method. *Int. J. Food Microbiol.* **163**, 89-95.
- Kaban, G. (2013) Sucuk and pastırma: microbiological changes and formation of volatile compounds. *Meat Sci.* **95**, 912-918.
- Kim, T. J., Na, Y. R., and Lee, J. I. (2005) Investigation into the basis of chloramphenicol and tetracycline resistance in *Staphylococcus intermedius* isolated from cases of pyoderma in dogs. *J. Vet. Med. B* **52**, 119-124.
- Landeta, G., Curiel, J. A., Carrascosa, A. V., Muñoz, R., and de las Rivas, B. (2013) Technological and safety properties of lactic acid bacteria isolated from Spanish dry-cured sausages. *Meat Sci.* **95**, 272-280.
- Morandi, S., Brasca, M., Andrighetto, C., Lombardi, A., and Lodi, R. (2006) Technological and molecular characterisation of enterococci isolated from North-West Italian dairy products. *Int. Dairy J.* **16**, 867-875.
- Ogier, J. C. and Serror, P. (2008) Safety assessment of dairy microorganisms: the *Enterococcus* genus. *Int. J. Food Microbiol.* **126**, 291-301.
- Ojha, K. S., Kerry, J. P., Duffy, G., Beresford, T., and Tiwari, B. K. (2015) Technological advances for enhancing quality and safety of fermented meat products. *Trends Food Sci. Tech.* **44**, 105-116.
- Ouoba, L. I. I., Lei, V., and Jensen, L. B. (2008) Resistance of potential probiotic lactic acid bacteria and bifidobacteria of African and European origin to antimicrobials: Determination and transferability of the resistance genes to other bacteria. *Int. J. Food Microbiol.* **121**, 217-224.
- Peters, J., Mac, K., Wichmann-Schauer, H., Klein, G., and Ellerbroek, L. (2003) Species distribution and antibiotic resistance patterns of enterococci isolated from food of animal origin in Germany. *Int. J. Food Microbiol.* **88**, 311-314.
- Petersen, A. and Jensen, L. B. (2004) Analysis of *gyrA* and *parC* mutations in enterococci from environmental samples with reduced susceptibility to ciprofloxacin. *FEMS Microbiol. Lett.* **231**, 73-76.
- Reviriego, C., Eaton, T., Martín, R., Jiménez, E., Fernández, L., Gasson, M. J., and Rodríguez, J. M. (2005) Screening of virulence determinants in *Enterococcus faecium* strains isolated from breast milk. *J. Hum. Lac.* **21**, 131-137.
- Szakacs, T. A., Kalan, L., McConnell Eshaghi, A., Shahinas, D., McGeer, A., Wright, G. D., Low D. E., and Patel, S. N. (2014) Outbreak of vancomycin-susceptible *Enterococcus faecium* containing the wild type *vanA* gene. *J. Clin. Microbiol.* **52**, 1682-1686.
- Talon, R. and Leroy, S. (2011) Diversity and safety hazards of bacteria involved in meat fermentations. *Meat Sci.* **89**, 303-309.
- Tamura, K., Dudley, J., Nei, M., and Kumar, S. (2007) MEGA 4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* **24**, 1596-1599.
- Valenzuela, A. S., Omar, N., Abriouel, H., López, R. L., Veljovic, K., Cañamero, M. M., Topisirovic, M. K. L., and Gálvez, A. (2009) Virulence factors, antibiotic resistance and bacteriocins in enterococci from artisan food of animal origin. *Food*

Control **20**, 381-385.

Yogurtcu, N. N and Tuncer, Y. (2013) Antibiotic susceptibility patterns of *Enterococcus* strains isolated from Turkish Tulum cheese. *Int. J. Dairy Technol.* **66**, 236-242.

Yüceer, Ö. and Özden Tuncer, B. (2015) Determination of antibiotic resistance and biogenic amine production of lactic acid bacteria isolated from fermented Turkish sausage (sucuk). *J. Food Safety* **35**, 276-285.