ARTICLE

Purification and Identification of a Natural Antioxidant Protein from Fertilized Eggs

Shaohua Yang^{1†}, Lulu Wang^{1†}, Ying Wang^{1†}, Xiaoqian Ou¹, Zhaoyuan Shi¹,
Chongchong Lu¹, Wei Wang², and Guoqing Liu^{1,3*}

¹College of Food Science and Engineering, Hefei University of Technology, 193 Tunxi Road, Hefei, Anhui 230009, P. R. China

²Agricultural Products Quality and Safety Supervision and Management Bureau, Xuancheng, Anhui 242000, P. R. China

³Wanjiang Institute of Poultry Technology, Hefei University of Technology, Xuancheng Campus, Xuancheng, Anhui 242000, P. R. China

OPEN ACCESS

Received June 26, 2017

Revised October 4, 2017

Accepted October 6, 2017

*Corresponding author

Guoqing Liu

College of Food Science and Engineering, Hefei University of Technology, Hefei, Anhui 230009, China
Tel/Fax: +86-551-65785485

E-mail: 13675512000@163.com

[†]These authors contributed equally to this work.

Copyright © Korean Society for Food Science of Animal Resources

This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Fertilized hen eggs are rich in a variety of bioactive ingredients. In this study, we aimed to obtain an antioxidant protein from fertilized eggs and the radical scavenging abilities on 1, 1-diphenyl-2-picrylhydrazyl (DPPH), hydroxyl radical ($\cdot\text{OH}$), superoxide anion ($\text{O}_2^{\cdot-}$) were used to evaluate the antioxidant activity of the purified protein. During 20 d of incubation, the radical scavenging ability of protein extracted from fertilized eggs exhibited significantly differences and the protein on day 16 showed higher antioxidant capacity. Based on this, the antioxidant protein of the samples on day 16 were isolated for the follow-up study. With a molecular weight 43.22 kDa, the antioxidant protein was purified by Diethylaminoethyl cellulose -52 (DEAE-52) column and Sephadex G-100. The LC-MS analysis showed that the purified protein molecular weight was 43.22 kDa, named D2-S. The sequence of amino acids was highly similar to ovalbumin and the coverage reached to 84%. The purified protein showed a radical scavenging rate of $52.34 \pm 3.27\%$ on DPPH and $63.49 \pm 0.25\%$ on $\cdot\text{OH}$, respectively. Furthermore, the C-terminal amino acid sequence was NAVLFFGRCVSP, which was consistent with the sequence of ovabumin. These results here indicated that purified protein may be a potential resource as a natural antioxidant.

Keywords fertilized eggs, antioxidant protein, purification, identification

Introduction

Reactive oxygen species (ROS) are products of cellular metabolism in living organisms (Yigit *et al.*, 2014), and high levels of ROS can cause lipid peroxidation (Yamaguchi *et al.*, 1994), protein carbonylation (Repine *et al.*, 1981) and DNA damage (Cerutti, 1985). Meanwhile, ROS may lead to a series of diseases, such as diabetes, hypertension, cancer and atherosclerotic cardiovascular disease (Finkel *et al.*, 2000; Memarpour-Yazdi *et al.*, 2012). Recently, more and more researchers had focused on extraction of novel, safe and natural antioxidants, for instance, carotenoids and vitamin A (Choudhry *et al.*, 2016; Korekar *et al.*, 2014; Maadane *et al.*, 2015; Páramo *et al.*, 2012). Such antioxidants are mainly obtained from vegetables, fruits, foods (Geetha *et al.*, 2002; Podsedk, 2007), and are regarded as the second antioxidant defense line in living organisms (Yigit *et al.*, 2014). In addition to these anti-

oxidants, there also contain antioxidant glutathione peroxidase (GSH-Px) enzyme, superoxide dismutase (SOD) and catalase (CAT) (Lin *et al.*, 2005; Surai *et al.*, 1996; Wilson *et al.*, 1992). As the first antioxidant defense line, these components act as ROS scavenging to protect the cell structure with the second antioxidant defense line together. Therefore, bioactive proteins from foods have gained extensive attentions due to their safety and health benefits.

Considering its rich and highly digestible proteins (Lechevalier *et al.*, 2017), egg is an important nutrient due to many desirable functional attributes in daily diet (Baldwin *et al.*, 1986). Of these, hen eggs produce many bioactive ingredients during incubation stage (Surai *et al.*, 2002). Proteins derived from hen eggs and embryos mainly include ovalbumin, (Duan *et al.*, 2013), GSH-Px, SOD, and so on (Suria *et al.*, 2002). Researchers showed that these proteins had various activities, such as, antioxidant, antibacterial and anti inflammatory activities (Duan *et al.*, 2014a; Duan *et al.*, 2014b; Erdmann *et al.*, 2008; Miguel *et al.*, 2007; Mine *et al.*, 2007). While embryos are developing, more oxygen was consumed, increasing the risk of being affected by ROS. Previous study has obtained the antioxidant peptide from embryo, and the peptide showed a high inhibition on linoleic acid autoxidation and ABTS radicals (Duan *et al.*, 2014a). Furthermore, the differences of antioxidant capacity in different tissues of chick embryo are also detected during each embryonic development of the entire stage (Chay *et al.*, 2011; Surai *et al.*, 2000). However, the information of antioxidant protein during embryonic development of the entire stage is still unclear. Thus, the objective of this study was to isolate and characterize a natural antioxidant protein from fertilized eggs.

Materials and Methods

Materials

Fertilized eggs were obtained from a hatchery (Changlv Native Products Co., China). Eggs were incubated at 37.8°C and 60% relative humidity in incubator (Photoshop Solar Energy Co., China). All eggs were turned through 90° every 90 min. Fertilized eggs were collected on days 0, 5, 7, 10, 12, 14, 16, 18, 20 of incubation. Six samples on same day were homogenized in a high-speed tissue triturator (Langyue instrument, China) and were freeze-dried at -80°C for 24 h. The totally dried samples were stored at -70°C for further analysis. All the testing chemicals were of analytical grade and purchased in Hefei Chemical Reagents Company (China).

Extraction of protein

The method to extract protein was conducted according to the method of Zhong *et al.* (2014) with some modifications. Dried samples were defatted by 20 volume (W/V) of acetone for 48 h and dried at 4°C for 12 h. Dried powder was collected and stored at -80°C. Based on previous work, proteins were extracted from the defatted dried powder by mixing with 31 volumes of PBS (0.2 M, pH 6.8) and maintaining at 55°C for 93 min. After centrifuged at 6000 rpm/min for 15 min at 4°C, the supernatant was collected and dialyzed against deionized water using a dialysis bag of 8-14 kDa for 48 h, the retentate were lyophilized and stored at -80°C for further analysis.

Determination of antioxidant activity of protein

DPPH radical scavenging activity

DPPH radical scavenging activity (RSA) was measured according to the method described by Je *et al.* (2005). 2.0 mL of each sample (10 mg/mL in PBS buffer) was added to 2 mL of 95% ethanol containing 0.1 mM DPPH radicals. The mixture was placed in the dark for 30 min, and the absorbance of the reaction solution was determined at 517 nm. The control was operated in the same way where 95% ethanol was used replaced the sample solution. Each experiment was repeated 3 times to determine the average value. The scavenging effect was calculated as the following equation:

$$\text{DPPH scavenging activity (\%)} = \frac{A_c - A_s}{A_c} \times 100$$

Where A_c is the absorbance of the control, A_s is the absorbance of the sample mixture.

O^{2•} radical scavenging activity

The O^{2•} radical scavenging activity was assayed according to the method described by Duan *et al.* (2014). Briefly, reaction solution was made up of sample solution (1.5 mL, 10 mg/mL protein in PBS buffer) and Tris-HCl buffer (4.5 mL, 50 mM, pH 8.0) were incubated at 25°C for 10 min, later 300 µL of 3 mM pyrogalllic acid (dissolved in 10 mM HCl) was added. The absorbance of reaction mixture at 325 nm was measured immediately at 30 s intervals. The autoxidation rate constant (Kb) was calculated from the curve of $A_{325\text{ nm}}$ vs time. Each experiment was repeated 3 times to determine the average value. The inhibitory actions of test samples against the autoxidation rate indicated their O^{2•} scavenging abilities.

Hydroxyl radical-scavenging activity

The Hydroxyl radical scavenging activity was assayed according to the method described by Li *et al.* (2008) with a few modifications. The •OH was generated by mixing 1.0 mL FeSO₄ (5 mM) and 1.0 mL H₂O₂ (0.1%), then, added 1 mL phenanthroline solution (5 mM), Mixed well and incubated at 37°C for 1 h, the absorbance of mixed solution was measured at 536 nm. The percentage scavenging effect was calculated as the following equation:

$$\text{Hydroxyl radical-scavenging activity (\%)} = \frac{A_3 - A_1}{A_2 - A_1} \times 100$$

Where A_1 absorbance of the solution containing FeSO₄, H₂O₂, phenanthroline solution; A_2 absorbance of solution containing phenanthroline and FeSO₄; A_3 absorbance of solution which has 1 mL sample solution.

Purification of antioxidant protein

Ion exchange chromatography

The crude protein samples on day 16 were dissolved in 10 mM phosphate buffer (pH 7.2), and centrifuged at 10,000 rpm for 15 min. The supernatant was collected and loaded on a DEAE cellulose 52 column (16×500 mm) previously equilibrated with 10 mM phosphate buffer (pH 7.2). The unbound protein was washed with the same buffer and the bound protein was eluted by increasing the concentration of NaCl (0.1, 0.3, 0.5 M) in phosphate buffer at a flow rate of 5 mL/min. After elution, the absorbance of tube was read at a wavelength of 280 nm, and four peaks including D1, D2, D3, D4 were collected. Each peak was pooled, dialyzed, freeze-dried, and the totally dried powders were used for DPPH radical scavenging assay.

Gel filtration chromatography

The most active fraction D2 was dissolved in phosphate buffer (10 mM, pH 7.2) and applied to a Sephadex G-100 column (10×300 mm). Then, eluted with 10 mM phosphate buffer (pH 7.2) at a flow of 1 mL/min, the elution was detected at 280 nm. The peak was pooled, dialyzed, freeze-dried for further study.

Identification of the purified protein

Electrophoresis

The molecular weight of protein was evaluated according to method of Laemmli (1970). Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was

performed with 5% stacking gel and 12% separating gel. The gels were stained with coomassie brilliant blue R-250 for 2 h, then, the gels were de-stained for 24 h. The protein was separated at 80 V for 30 min and at 120 V for 100 min. A pre-stained standard protein marker (18.0-94.0 kDa) was run in the gel to determine molecular weight of the proteins.

Liquid-phase mass spectrometry

The band was excised from the gel and identified by a LC-LTQ Orbitrap Velos Pro (Thermo Scientific, USA). The digestion of the band was detected according to Katayama (2001). After the digestion treatment, the supernatant was collected and completely dried for LC-MS analysis. Sample was separated by the Nano-HPLC liquid phase system EASY-n LC1000 and analyzed by LTQ Orbitrap Velos Pro (Thermo Scientific, USA). The mass spectrometry parameters were as follows: analysis of time 105 min, spray voltage 1.8 kV, interface heater temperature 250°C, with parent ion scanning range of 350-1800 m/z. The parameters of Information Dependent Acquisition (IDA) were as follows: fracture mode of collision-induced dissociation, normal chemical energy 35%, the q value 0.25, activation time 30 ms. The MASCOT V2.3 search engine (Matrix Science Ltd., U.K.) was adopted to analysis data, using the Gallus gallus database as search parameters.

Protein N-terminal sequencing

The D2-S protein (20 mg/mL) was added to SDS-PAGE. After electrophoresis, the strip was cut and placed in a centrifuge tube, then discolored at 50% CAN/25 mM NH₄HCO₃ solutions for 30 min. 15 ng/μL of trypsin and chymotrypsin were added and incubated for 40 min at 4°C and then, 5-10 μL of 25 mmol/L NH₄HCO₃ solution to each tube was added and water bath 37°C for 12 h. Subsequently, 1% TFA-50% ACN-50% water 200 μL were added for 1 h at 37°C water bath and transferred to another new EP tube. The lyophilized sample was dissolved in 0.1% formic acid, 2% acetonitrile, mixed and centrifugated at 13,200 rpm for 10 min at 4°C. The supernatant was subjected to Q Exactive mass spectrometer (Thermo Scientific, USA). Proteome Discoverer software was adopted to analyze the original data of the mass spectrometry to obtain the target protein sequence information.

Statistical analysis

The experiment was conducted in triplicate. All results were expressed as mean±standard deviation (n=3) and

Fig. 1. Antioxidant activity of proteins derived from fertilized eggs during different incubation days. The protein concentration was 10 mg/mL for determination of the radical scavenging abilities on DPPH and •OH. The value of each fraction is the mean $\pm$ standard deviation ($n=3$), different letters (a, b) above columns indicated significant differences ($p<0.05$).

were analyzed using SPSS 16.0 software (SPSS Inc., USA). One-way analysis of variance (ANOVA) was used to analyze data, the differences between the means were separated using Duncan's multiple range test. Results were considered statistically significant at $p<0.05$.

Results

Change in antioxidant activity of proteins

As shown in Fig. 1, the DPPH radical scavenging activity had no significant change before day 7, and then con-

Table 1. Inhibition of pyrogallol autoxidation rate of proteins derived from fertilized eggs

Incubation time (d)	Kb value ($\times 10^{-6}$)
0	52.39 $\pm$ 2.34 ^{de}
5	56.70 $\pm$ 2.31 ^f
7	50.00 $\pm$ 1.77 ^d
10	53.34 $\pm$ 2.62 ^{de}
12	41.78 $\pm$ 3.75 ^c
14	34.62 $\pm$ 1.16 ^b
16	21.24 $\pm$ 4.13 ^a
18	33.31 $\pm$ 1.29 ^b
20	30.70 $\pm$ 0.72 ^b

The results were expressed as means $\pm$ standard deviation ($n=4$), different characters (a, b) represent significant differences ($p<0.05$). Kb value means auto-oxidation rate of pyrogallol.

tinuously increased with the change of hatching days until day 16 and reached its maximum 37.89 $\pm$ 3.60%. Meanwhile, the protein on day 16 possessed the highest scavenging rates on •OH, 47.87 $\pm$ 1.88% in Fig. 1. As shown in Table 1, the value of Kb on day 16 was significantly smaller than the other days ($p<0.05$). The results suggested that protein on day 16 had the highest capacity to scavenging radicals.

Purification of antioxidant peptides

The protein on day 16 was dissolved in 10 mM phosphate buffer and applied to DEAE cellulose column, then, eluted with the same buffer. As shown in Fig. 2A, four peaks, number fraction D1 to D4, were collected, dialyzed and lyophilized separately. These fractions were detected their radical scavenging ability on DPPH and •OH

Fig. 2. Purification of antioxidant proteins by Ion Exchange Chromatography. (A) Elution profile of crude proteins separated by DEAE cellulose 52 column at a flow rate of 1 ml/min. Four peaks were obtained (D1, D2, D3, D4) from fertilized eggs. (B) The radical scavenging capacity on DPPH and •OH of each fraction with the concentration of 10 mg/mL.

Fig. 3. Purification of antioxidant proteins by Gel Filtration. (A) Elution profile of D2 following Sephadex G-100 gel chromatography, the protein concentration of each tube was determined at a wavelength of 280 nm. (B) The radical scavenging capacity on DPPH and •OH of fraction D2-S with the concentration of 10 mg/mL. The value of each fraction is the mean $\pm$ standard deviation ($n=3$) of figure (b) and (d).

(Fig. 2B). The results showed the DPPH and •OH radical scavenging ability of each peaks, among four fractions, fraction D2 showed the highest ability (Fig. 2B), reached $57.32 \pm 3.28\%$, $49.63 \pm 0.23\%$. Thus, fraction D2 was used for further purification.

The D2 peak was dissolved in 10 mM phosphate buffer and purified further with Sephadex G-100. The active peak D2 protein was rechromatographed on Sephadex G-100 column resulted in single peak (Fig. 3A), indicating the preparation of the antioxidant protein from fertilized eggs has been accomplished. Fig. 3B showed the radical scavenging ability on DPPH and •OH of fraction D2-S were $52.34 \pm 3.27\%$, $63.49 \pm 0.25\%$, respectively.

Characterization of antioxidant protein

The SDS-PAGE results showed that it's a single band and molecular weight was about 45 kDa. Fig. 4 The protein band was obtained and analyzed by EASY-Nlc 1000-LTQ Orbitrap Velos Pro (Thermo Finnigan). The purified protein was digested, isolated and analyzed by LC-MS, the obtained peptide sequences was compared with the sequences of the protein (*Gallus gallus*) in NCBI. The result exhibited that purified protein sequence was highly similar to ovalbumin (gi|28566340|gb| AAO43266.1) (Table 2) and sequence coverage reached 84%, this was a more reliable identification result. LC-MS analysis of purified protein showed its molecular weight was 43.22 kDa and the isoelectric point of 5.19.

According to the theoretical sequence of the sample, we used trypsin and chymotrypsin for protease digestion and

Fig. 4. Identification and SDS-PAGE analysis of the purified protein. The purified protein (20 mg / mL) was loaded onto 5% stacking gel and 12% separating gel. A wide molecular weight standard protein (10-180 kDa) acts as marker, the remaining three lanes is the same sample volume of D2-S.

subsequent matching analysis. Fig. 5A and 5B showed TIC maps of protein D2-S digested by trypsin and chymotrypsin. Fig. 5C and 5D showed C-terminal sequence secondary mass spectrometry of the protein D2-S. According to results of TIC map and secondary mass spectrometry, protein D2-S C-terminal sequence was NAVLFFGRVCSP,

Table 2. The comparison of sequences of purified protein with ovalbumin (gi|28566340|gb|AAO43266.1|)

	1-10	11-20	21-30	31-40	41-50
1	MGSIGAASME	FCFDVFKELK	VHHANENIFY	CPIAIMSALA	MVYLGAKDST
51	RTQINKVVR	DKLPGFGDSI	EAQCGTSVNV	HSSLRDILNQ	ITKPNDEVYSF
101	SLASRLYAE	RYPILPEYLQ	CVKELYRGGL	EPINFQTAAD	QARELINSWV
151	ESQXNGIIRN	VLQPSSVDSQ	TAXVLVNAIV	FKGLWEKAFK	DEDTQAMPFR
201	VTEQESKPVQ	MMYQIGLFRV	ASMASEKMKI	LELPFASGTM	SMLVLLPDEV
251	SGLEQLESII	NFEKLTEWTS	SNVMEERKIK	VYFPRMKMEE	KYNLTSVLMA
301	MGITDVFSSS	ANLSGISSAE	SLKISQAVHA	AHAEINEAGR	EVVGSAEAGV
351	DAASVSEEFR	ADHPFLFCIK	HIATNAVLFF	GRCVSP	

The matched peptides shown in black, the sequence coverage reached 84%. The molecular weight was 43.220 Da and the calculated pI value was 5.19 of purified protein.

Fig. 5. the protein D2-S C-terminal sequences. The TIC map of the sample after digestion; A, Trypsin digestion; B, Chymotrypsin digestion.

Fig. 5. the protein D2-S C-terminal sequences. (Continued) The TIC map of the sample after digestion; C and D, terminal sequence secondary mass spectrometry.

which was consistent with the sequence of ovabumin.

Discussion

Fertilized hen eggs are traditionally considered as dietary supplement in many Asian countries for their nutrition value. During incubation, embryo requires more oxygen to maintain energy, especially in the formation of tissues and organs, and the improvement of respiratory function (Vlecket *et al.*, 1980). In this work, the protein derived from

fertilized eggs showed unequal radical scavenging capacity on DPPH, •OH, O₂^{•-} in different incubation periods, and the ability of radical scavenging was significantly increased after 10-d incubation, which was consistent with the results of Duan (2014a). The protein derived from fertilized eggs on day 16 with stronger radical scavenging capacity on DPPH, •OH, O₂^{•-}, which suggested that it had the strongest capacity to convert free radicals to stable products, prevent the radical chain reaction, and avoid cell damage caused by hydroxyl radicals-the most reac-

tive of all the reduced forms of dioxygen (Je *et al.*, 2007; Sivapriya *et al.*, 2007). Therefore, the protein may be a potential antioxidant that exists in the embryo to protect embryo from ROS. Furthermore, protein derived from fertilized eggs may be an efficient antioxidant.

Based on the results of antioxidant activity, an antioxidant protein was isolated and purified from fertilized eggs on day 16. The SDS-PAGE illustrated that its molecular weight was about 45 kDa. LC-MS analysis result showed that purified protein sequence was highly similar to ovalbumin (coverage reached 84%), compared with ovalbumin sequence. In order to understand more about protein D2-S, we have studied its C-terminal sequence, result showed that protein D2-S C-terminal sequence was NAVLFF-GRCVSP, which was consistent with the sequence of Ovalbumin. ovalbumin is a 386-aa residue protein, molecular weight of about 43 kDa. Duan (2013) reported that the change of ovalbumin was not significantly during incubation, Xu *et al.* (2007) reported that the hydrolysates of ovalbumin showed a distinctly inhibitory action to $\cdot\text{OH}$, $\text{O}_2^{\cdot-}$ and linoleic acid autoxidation. Abeyrathne *et al.* (2014) showed that peptide derived from ovalbumin were hydrolyzed by alcalase and trypsin showed strong iron- and copper-binding capacities and antioxidant capacity. The sequence of antioxidant protein obtained from fertilized eggs on day 15 exhibited highly similar to ovalbumin (Yang *et al.*, 2016). Thus, we suspected that the purified protein in this study was a new antioxidant protein. There has been reported that antioxidant activity of some proteins was attributed to the presence of leucine, histidine and proline amino acid sequence (Peñaramos *et al.*, 2004). In this study, we discovered that the content of ovalbumin in 16 incubation days was 33.72 ± 1.33 $\mu\text{g/mL}$, and the radical scavenging ability on DPPH, $\cdot\text{OH}$ was $48.36 \pm 0.48\%$, $53.27 \pm 1.37\%$, respectively. Meanwhile, the purified protein possessed antioxidant capacity. The amino acids sequence was highly similar to ovalbumin and had leucine, histidine and proline amino acid. Therefore, we named this protein D2-S.

The outcome of present study revealed that antioxidant protein from fertilized exhibited antioxidative properties. Considering the antioxidant capacity of protein in the whole process of chicken embryo hatching, an antioxidant protein D2-S (MW: 43.22 kDa) isolated from fertilized eggs possessed potential antioxidant and free radical scavenging properties. However, the real mechanism of its antioxidant capacity is not clearly known. Thus, it is necessary to fully characterize this active mechanism and

evaluate its capacity in food system.

Acknowledgements

This work was supported by the Fundamental Research Funds for Industry-University-Research of Xuancheng Campus in Hefei University of Technology (XC2016JZ-BZ03), “the Fundamental Research Funds for the Central Universities” (JZ2017HGTA0228), Wanjiang Institute of Poultry Technology (XC2015AKKG002), the Science and Technology Bureau of Xuancheng (XC2015XKKJ02) and Science and Technology Bureau of Anhui (XC2015AK-KG003).

References

- Abeyrathne, E. D., Lee, H. Y., Jo, C., Suh J. W., and Ahn, D. U. (2014) Enzymatic hydrolysis of ovalbumin and the functional properties of the hydrolysates. *Poultry Sci.* **93**, 2678-2686.
- Baldwin, R. E. (1986) Functional properties of eggs in Foods. *Egg Sci. Technol.* 405-482.
- Cerutti, P. A. (1985) Prooxidant states and tumor promotion. *Science* **227**, 375-381.
- Chay Pak Ting, B. P., Mine, Y., Juneja, L. R., Okubo, T., Gauthier, S. F., and Pouliot, Y. (2011) Comparative composition and antioxidant activity of peptide fractions obtained by ultrafiltration of egg yolk protein enzymatic hydrolysates. *Membranes* **1**, 149-161.
- Choudhry, Q. N., Kim, M. J., Kim, T. G., Pan, J. H., Kim, J. H., Park, S. J., Lee, J. H., and Kim, Y. J. (2016) Saponin-based nano emulsification improves the antioxidant properties of Vitamin A and E in AML-12 Cells. *Molecular Sci.* **17**, 1406-1418.
- Duan, X., Li, M., Wu, F. F., Yang, N., Nikoo, M., Jin, Z., and Xu, X. (2013) Postfertilization changes in nutritional composition and protein conformation of hen egg. *J. Agric. Food. Chem.* **61**, 12092-12100.
- Duan, X., Ocen, D., Wu, F. F., Li, M., Yang, N., Xu, J., Chen, H. Y., Huang, L. Q., Jin, Z., and Xu, X. (2014a) Purification and characterization of a natural antioxidant peptide from fertilized eggs. *Food Res. Int.* **56**, 18-24.
- Duan, X., Wu, F. F., Li, M., Yang, N., Wu, C., Jin, Y. M., Yang, J. J., Jin, Z., and Xu, X. (2014b) Naturally occurring angiotensin I-Converting enzyme inhibitory peptide from a fertilized egg and its inhibitory mechanism. *J. Agric. Food. Chem.* **62**, 5500-5506.
- Erdmann, K., Cheung, B. W., and Schröder, H. (2008) The possible roles of food-derived bioactive peptides in reducing the risk of cardiovascular disease. *J. Nutr. Biochem.* **19**, 643-654.
- Geetha, S., Sai, R. M., Singh, V., Ilavazhagan, G., and Sawhney, R. C. (2002) Antioxidant and immunomodulatory properties of seabuckthorn (*Hippophae rhamnoides*)-An *in vitro* study. *J. Ethnopharmacol.* **9**, 373-378.

- Huang, W., Deng, Q., Xie, B., Shi, J., Huang, F. H., Tian, B. P., Huang, Q. D., and Xue, S. (2010) Purification and characterization of an antioxidant protein from *Ginkgo biloba* seeds. *Food Res. Int.* **43**, 86-94.
- Je, J., Park, P., Kwon, Y., and Kim, S. (2005) A novel angiotensin I Converting enzyme inhibitory peptide from alaska pollack (*Theragra chalcogramma*) frame protein hydrolysate. *J. Agric. Food Chem.* **52**, 7842-7845.
- Je, J. Y., Qian, Z. J., Byun, H. G., and Kim, S. K. (2007) Purification and characterization of an antioxidant peptide obtained from tuna backbone protein by enzymatic hydrolysis. *Process Biochem.* **42**, 840-846.
- Katayama, H., Nagasu, T., and Oda Y. (2001) Improvement of in gel digestion protocol for peptide mass fingerprinting by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.* **15**, 1416-1421.
- Laemmli, U. K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685.
- Lechevalier, V., Guérin-Dubiard, C., Anton, M., Beaumalc, V., Briand E. D., Gillard, A., Le Gouar, Y., Musikaphun, N., Tanguy, G., Pasco, M., Dupont, D., and Nau, F. (2017) Pasteurisation of liquid whole egg: Optimal heat treatments in relation to its functional, nutritional and allergenic properties. *J. Food Eng.* **195**, 137-149.
- Li, Y., Jiang, B., Zhang, T., Wu, W., and Liu, J. (2008) Antioxidant and free radical-scavenging activities of chickpea protein hydrolysate (CPH). *Food Chem.* **106**, 444-450.
- Lin, Y. F., Tsai, H. L., Lee, Y. C., and Chang, S. J. (2005) Maternal vitamin, E supplementation affects the antioxidant capability and oxidative status of hatching chicks. *J. Nutr.* **135**, 2457-2461.
- Maadane, A., Merghoub, N., Ainane, T., Arroussi, H. E., Benhima, R., Amzazib, S., Bakrib, Y., and Wahby, I. (2015) Antioxidant activity of some Moroccan marine microalgae: Pufa profiles, carotenoids and phenolic content. *J. Biotechnol.* **215**, 13-19.
- Memarpour-Yazdi, M., Asoodeh, A., and Chamani, J.K. (2012) A novel antioxidant and antimicrobial peptide from hen egg white lysozyme hydrolysates. *J. Funct. Foods* **4**, 278-286.
- Miguel, M., Manso, M., Aleixandre, A., Alonso, M. J., Salaices, M., and López-Fandiño, R. (2007) Vascular effects, angiotensin I-converting enzyme (ACE)-inhibitory activity, and anti-hypertensive properties of peptides derived from egg white. *J. Agric. Food Chem.* **55**, 10615-10621.
- Mine, Y. (2007) Egg proteins and peptides in human health-Chemistry, bioactivity and production. *Curr. Pharmaceut. Design* **13**, 875-884.
- Peñaramos, E. A., Xiong, Y. L., and Arteaga, G. E. (2004) Fractionation and characterization for antioxidant activity of hydrolyzed whey protein. *J. Sci. Food Agric.* **84**, 1908-1918.
- Podsedek, A. (2007) Natural antioxidants and antioxidant capacity of Brassica vegetables: A review. *LWT-Food Sci. Technol.* **40**, 1-11.
- Qian, Z. J., Jung, W. K., and Kim, S. K. (2008) Free radical scavenging activity of a novel antioxidative peptide purified from hydrolysate of bullfrog skin, *Rana catesbeiana* Shaw. *Biore-sour. Technol.* **99**, 1690-1698.
- Repine, J. E. Pfenninger, O. W., Talmage, D. W., Berger, E. M., and Pettijohn, D. E. (1981) Dimethyl sulfoxide prevents DNA nicking mediated by ionizing radiation or iron/hydrogen peroxide-generated hydroxyl radical. *Proc. Natl. Acad. Sci.* **78**, 1001-1003.
- Sivapriya, M. and Leela, S. (2007) Isolation and purification of a novel antioxidant protein from the water extract of Sundakai (*Solanum torvum*) seeds. *Food Chem.* **104**, 510-517.
- Surai, P. F. (2000) Effect of selenium and vitamin E content of the maternal diet on the antioxidant system of the yolk and the developing chick. *Brit. Poultry Sci.* **41**, 235-243.
- Surai, P. F. (2002) Natural Antioxidants in avian nutrition and reproduction. *Nottingham University Press*. Nottingham.
- Surai, P. F., Noble, R. C., and Speake, B. K. (1996) Tissue-specific differences in antioxidant distribution and susceptibility to lipid peroxidation during development of the chick embryo. *Biochimica et Biophysica Acta.* **1304**, 1-10.
- Turrens, J. F. (2003) Mitochondrial formation of reactive oxygen species. *J. Physiol.* **552**, 335-344.
- Vleck, C. M., Vleck, D., and Hoyt, D. F. (1980) Patterns of metabolism and growth in avian embryos. *Am. Zoologist* **20**, 405-416.
- Wassmann, S., Wassmann, K., and Nichenig, G. (2004) Modulation of oxidant and antioxidant enzyme expression and function in vascular cells. *Hypertension* **44**, 381-386.
- Wilson, J. X., Lui, E. M., and Del Maestro, R. F. (1992) Developmental profiles of antioxidant enzymes and trace metals in chick embryo. *Mech. Ageing Development* **65**, 51-64.
- Xu, M., Shangguan, X., Wang, W., and Chen, J. (2007) Antioxidative activity of hen egg ovalbumin hydrolysates. *Asia Pacific J. Clin. Nutr.* **16**, 178-182.
- Yamaguchi, S., Sakurada, S., and Nagumo, M. (1994) Role of intracellular SOD in protecting human leukemic and cancer cells against superoxide and radiation. *Free Rad. Biol. Med.* **17**, 389-395.
- Yang, S. H., Tang, Z. J., Tang, S. H., Zhang, T. F., Tang, F., Wu, Y., Wang, Y., Wang, L. L., and Liu, G. Q. (2016) Purification and characterization of an antioxidant protein from fertilized eggs. *Korean Soc. Food Sci. Anim. Resour.* **36**, 791-798.
- Yigit, A. A., Panda A. K., and Cherian G. (2014) The avian embryo and its antioxidant defense system. *World's Poultry Sci.* **70**, 563-574.
- Zhong, C., Zhou, Z., Wang, R., Han, P. P., and Jia, S. R. (2014) Studies on protein extraction from *Caragana korshinskii* Kom. using acetone precipitation method. *Springer Berlin Heidelberg* **251**, 1609-1616.