

We examined the empirical relationship between stellar activity and rotation period found in our earlier work for the lower main sequence stars to interpret it on the basis of a simplified astronomical dynamo theory.

From the analysis it is found that the magnetic field strength B_0 activated under the dynamo action is proportional to $(\Omega v \rho l)^{1/2}$ where Ω , v , ρ , l are the characteristic angular velocity, flow velocity, mass density and the scale of the dynamo active region, respectively.

Hydrogen Emission Spectra of Quiescent Prominences

Kap-Sung Kim and Young-Key Minn

Department of Astronomy & Space Science, Kyung Hee University, Yong-In

We have calculated the models of quiescent prominences which satisfy simultaneously the constraints of radiative transfer, statistical equilibrium and charge-particle conservation. The prominence is treated as an infinite slab of finite thickness, standing vertically on the solar surface illuminated on both sides by the photospheric, chromospheric and coronal radiation field. The effect of filaments in prominence on the spectral line emission has been examined by considering the fine geometrical structure. We will present the result of our calculation on hydrogen line emissions for the prominence model with filamentary structure.

회전하는 항성내부 철핵의 수축 모형

육 인 수 · 현 정 준

서울대학교 천문학과

초신성 폭발은, 질량이 $8M_{\odot}$ 보다 큰별의 급격한 중력수축으로 시작된다. 기존의 구대칭 수축 모형은 상부 맨틀을 날리보낼 만큼 충분한 운동에너지가 중력에너지로부터 변환되지 않고 있다. 따라서 우리는 또 다른 에너지 변환기구로서 별의 회전효과를 고려하였다.

수축 모형은 크게 유체역학 방정식, 상태 방정식 그리고 중성미자 전달 방정식 등 세 부분으로 구성된다. 이들은 구형 좌표계에서 기술되었으며 특히 중성미자와 물질과의 상호작용은 상세히 취급되었다.

Uncertainties in the Star-Count Analysis

Seung Soo Hong and See-Woo Lee

Department of Astronomy, Seoul National University

We have examined how sensitively the extinction derived from the star-count depends on such factors as the plate limit, the size of counting reseaus, the non-linearity in the number distribution of stars with magnitude, and the angular resolution demanded by the given problem. We let the Poisson distribution portray the statistical nature of the countings, and chose the region containing globule Barnard 361 as an example field. Uncertainties due to combinations of the various factors involved in the star-count analysis are presented in graphic forms: Dynamic range in the extinction measurements by the star-count analysis is evaluated as a function of reseau size for varying plate