

Our results indicate as follows: 1) In the case of negligible diffusion, clusters are tidally truncated to the theoretical tidal radius at a perigalacticon. 2) There is no apparent orbital phase-dependence of the tidal radius for clusters with eccentric orbits. 3) In timescales of moderately efficient two-body relaxation, diffusion processes significantly modify the structure of outer regions such that the limiting radius is likely to be comparable to the tidal radius at an apogalacticon. 4) Galactic tidal torque gives a rise to the isotropy in velocity dispersion of the outer cluster region. For relaxed clusters, the velocity dispersions are likely to be isotropic in the cores, anisotropic in the envelopes, and isotropic near the limiting radii. 5) Stars with direct orbits are less stable so that the prolonged tidal interaction can lead to an apparent retrograde rotation in the outer cluster regions. For anisotropic clusters, the effect of tidally induced retrograde rotation likely to be extended into relatively small radii by stars of highly eccentric orbits.

Successive Merging of Stars in Dense Stellar Systems

Lee, Hyung Mok

(Department of Earth Sciences, Pusan University)

In dense stellar systems, close encounters between stars can lead to mergers if the velocity dispersion is sufficiently smaller than the escape velocity from the stellar surface. Simple theory describing evolution of the mass function in the presence of successive merging is presented. This process transforms low-mass stars into high-mass stars in the 'merging' time scale. Eventually a significant fraction of the stellar system's core mass can be locked into in a small number of high mass stars, provided that stellar evolution is slower than the rate of merging. The mass function becomes a power law when the merging cross section is assumed to be a power law on the stellar mass. The implications of the simple analysis for a realistic stellar system are discussed. However, it is not yet clear whether a central black hole can be formed through this process in nuclei of galaxies because of many other uncertainties.

Spherically Symmetric Radiation Hydrodynamics near Eddington Flux

Park, Myeong-Gu

(Kyungpook National University)

Time-independent spherical accretion of gas onto the surface of compact star is studied. The previous, simplified, special relativistic treatment is generalized under general relativity. The flow velocity is allowed to approach the speed of light and the gravitational field can be arbitrarily strong if the field can be represented by the Schwarzschild metric. This general relativistic effects become increasingly important when the flux is near Eddington limit.

Results in current literatures are found to be incorrect in many circumstances. When the flux is sufficiently close to the Eddington limit, the flow velocity increases with decreasing radius far from the compact star, reaches a maximum at an intermediate radius, and decreases at small radii. If the external luminosity (luminosity not related to the accreting gas) is centralized, steady accretion is possible only when the luminosity at infinity does not exceed 76% of the

Eddington luminosity for a 10km neutron star with mass 1.4 times solar mass.

On the A-type Contact System DO Cas

Kyu-Dong Oh

(Chunnam National University)

Young-Sook Ahn

(Institute of Space Science and Astronomy)

New UBV photoelectric observations of EB-type contact binary DO Cas were obtained at the Sobacksan Observatory. These observations covered the eclipse portions of the light curves and twice the minimum light. Both the light curves and the radial velocity curve for the primary of DO Cas (Mannino 1958) have been solved simultaneously using the Wilson-Devinney synthesis code. New geometric, photometric, and absolute elements for this system are given.

Abundance and Chemistry of Interstellar HOCO⁺

Minh, Young Chol

(Daeduk Radio Astronomy Observatory Institute of Space Science and Astronomy)

M.K. Brewer, W.M. Irvine

(Five College Radio Astronomy Observatory University of Massachusetts, U.S.A.)

P. Friberg, L.E.B. Johansson

(Onsala Space Observatory Chalmers University of Technology, Sweden)

We derive HOCO⁺ column densities $\sim 10^{15} \text{cm}^{-2}$ toward the Galactic center and $< 10^{12} \text{cm}^{-2}$ for cold dark clouds from observations and an LVG model. We mapped the HOCO⁺ $4_{04}-3_{03}$ line toward Sgr A. The fractional abundance of HOCO⁺ in the Galactic center region is three orders of magnitude larger than predicted by quiescent ion-molecule chemistry and an order of magnitude larger than predicted by a MHD shock model. If HOCO⁺ traces interstellar CO₂, the implied high abundance ($[\text{CO}_2] \sim [\text{CO}]$) in the Galactic center may result from UV photolysis of grain mantles.

세포형 구조 우주와 Fractal

송 두 종

(천문우주과학연구소)

뜨거운 보이지않는 물질이 지배하는 표준 우주 모형 안에서 관측된 은하 및 은하단분포의 두점상 관관계 함수로 대표되는 큰크기 구조는 “세포형” 구조를 보임을 알 수 있다. “세포형” 큰크기 구조를 fractal 구조와 비교함으로써 우주의 균질성의 한계를 규정 짓는 “한계거리”가 존재해야 함을 보였고, 이에 의한 우주 배경 복사 불균질성의 각의 크기를 계산할 수 있었다.

Compact HII region K3-50 주변의 분자운에 대한 연구

노덕규 · 김효령 · 김봉규 · 민영철

(천문우주과학연구소)