

G802

**P Element-Mediated Germ-Line Transformation of *Bombyx mori***

Sun Young Kim\*, Seung Hyun Sung, Jin Sung Lee, Sang Mong Lee,  
Jae Sam Whang and Dong Sang Suh  
Department of Genetic Engineering, Sungkyunkwan university

This study was use of P element of the *Drosophila melanogaster* to obtain transformed *Bombyx mori*. To induce transformations by artificial introduction of genetic materials, a simple method of microinjection of DNA into eggs was developed using a microinjector and micromanipulator. The DNA to be injected is comprised of two components. The first consists of a helper plasmid containing a defective P element that although capable of producing the P transposase which can act in trans on another P transposon, is itself immobile. The second component consists of a transposon construct. The vector for this study is pUCHsneo. pUCHsneo has *lacZ* which has been used in many different kinds of expression assay. Amount of coinjected DNA is 0.06ng. The hatchability is about 1%. The larvae derived from DNA injected eggs and their progenies. To identify transformants, we analyzed by PCR and X-gal staining.

G803

**Production of the Anti-HLA Monoclonal Antibody Using Human Hybrid Myeloma**

Jee Young Ahn\*, Hye Joung Lee, Kwang Min Lee and Dong Sang Suh  
Department of Genetic Engineering, Sungkyunkwan University

During transplant operation, we need specific antibodies against HLA antigens for the HLA typing between donor and recipient. On the production of the anti-HLA monoclonal antibody in the human system, up to this point available partner cell was not developed. We must develop more available fusion partner cell, therefore we fused SKO-007, human myeloma, and HeLa cell, human cervix carcinoma. We made more effective clones as human fusion partner cells. As a result, their doubling time is shorter than SKO-007 and they are 6-thioguanine resistant. The most available clone, HS-7, of them has the shortest doubling time, 35 hours, and its chromosome number is 70. It has been using for the production of the anti-HLA monoclonal antibody secreting hybridoma. It can be used as fusion partner cell for production of several monoclonal antibodies.