

F317

Isolation of the *Vibrio vulnificus* DNA sequences which complement the phenotype of *Escherichia coli* defective in starvation sigma factor

Kyoung-Suk Shin^{1*}, Hee-Joon Myung¹, and Kyu-Ho Lee²
¹Dept. Microbiol., ²Dept. Environ. Sci., Hankuk Univ. Foreign Studies.

As an effort on the identification of the genes required for starvation-survival of *V. vulnificus*, the library was prepared of *V. vulnificus* genomic DNA partially digested by *Sau3AI*. Several clones expressing high activity of β -galactosidase were isolated from the transformants of *rpoS*-defective *E. coli* cell containing *rpoS*-dependent promoter, *bolA::lacZ* fusion. One of the *rpoS*-complementing plasmids, pSK22 includes a small sized, 403bp, DNA insert. The deduced polypeptide sequence of the putative ORF is composed of 56aa residues of which size (about 6kDa) was confirmed by SDS-PAGE of IPTG-induced cell extract, and does not show any significant similarity with the known bacterial proteins. In addition to induction of *bolA* promoter by pSK22, another *rpoS*-dependent promoter, *katE* is also induced more than 25-folds by the presence of this plasmid in *E. coli* cell. We have constructed a series of plasmids including the different lengths of the coding region. The effect of these deleted plasmids on the expression of *rpoS*-dependent promoters is under the investigation. We discuss the possible role of this short DAN sequence in ecological and physiological cycle of *V. vulnificus* in nature.

F318

Molecular Analysis of the *aroB*, a Gene Encoding DHQ Synthase in *Corynebacterium glutamicum*

Han Min-Ah*, Lee Myeong-Sok Dept. of Biology,
Sookmyung Women's Univ., Seoul 140-742, Korea

The *aroB* gene encoding dehydroquinase synthase of *Corynebacterium glutamicum* encoding dehydroquinase synthase has been cloned by complementation of the auxotrophic mutant of *Escherichia coli* with the genomic DNA library. The recombinant plasmid containing 1.4 kb fragment complemented the DHQ synthase-deficient mutant of *E. coli*. The nucleotide sequence analysis of the subcloned DNA has been determined. The sequence contained an open reading frame of 360 codons, from which a protein with a molecular weight of about 38 kDa could be predicted. Alignment of different prokaryotic and eukaryotic *aroB* gene products reveals an overall identity ranging from 32 to 56%.