

The Distribution of Chlorinated Pesticides in Surface Sediments from Masan-Chinhae Bay

홍상희*, 오재룡², 임운혁², 이인숙¹, 이수형²
¹이화여자대학교 생물학과
²한국해양연구소 화학연구단

The chlorinated pesticides concentrations in surface sediment from Masan-Chinhae Bay were the highest among that of ten major harbors in Korea (Sokch'o, Uljin, P'ohang, Ulsan, Pusan, Masan, Chinhae, Mokp'o, Kunsan, and Manrip'o). In this study, the surface sediments from twenty station in Masan-Chinhae Bay were investigated.

The concentrations of total chlorinated pesticides (27 compounds) in sediments were in the range of 0.67 - 94.2 ng/g. The highest concentrations were found in J1 and M5. The concentrations were higher in Masan Bay than in Chinhae Bay and tend to higher near urban areas. Chlorinated pesticides in twenty stations were dominated by DDT compounds (o,p'-DDT, p,p'-DDT, o,p'-DDD, p,p'-DDD, o,p'-DDE, p,p'-DDE). The average concentrations of DDE, DDD, and DDT were 6.46, 3.58, and 4.35ng/g dry weight respectively. β -HCH and γ -chlordane were the dominant in hexacyclohexane and chlordane compounds respectively. The concentrations of dieldrin (degradation product of aldrin) were higher than those of aldrin.

Species Diversity and Niche Breadth of the *Zalkova serrata* community in Mt. Daedun

이 윤원, 이 민순
 중부대학교 생명자원학부

As general, the species diversity (H'), probability of interspecific encounter (Δ_1), Levin's niche breadth (Sh), Simpson's measure of diversity (Δ_2), ratio of interspecific to intraspecific competition (Δ_4) were decreasing as the altitude develops. Whereas the intraspecific competition ($1-\Delta_1$) was increasing. And by Shannon-Weaver formula, the maximum possible diversity (H'_{max}), evenness (J') and dominance were represented as irrelevant to the altitude.

On the other hand, upper layer was decreasing as Simpson's measure of diversity (Δ_2) going higher, but it was increasing as D.B.H. getting lower. And Levin's niche breadth (Sh) was proportioned to the size of D.B.H..