

carbamate. On the other hand, cinnamyl 4-methoxybenzyl ether reacted with CSI to give cinnamyl N-(4-methoxybenzyl) carbamate.

[PD1-24] [ 04/21/2000 (Fri) 14:50 - 15:50 / [1st Fl, Bldg 3] ]

**The New Erythromycin A derivatives with C-9 oxime as a treatment of Helicobacter Pylori.**

Nam GS, Kim SH, Kim JH, Shin JH

Biochemicals research center, Korea Institute of Science & Technology; <sup>1</sup>Department of chemistry Seoul National University

Clarithromycin is used as a H. Pylori treatment and is one of the top five best-selling antibiotics in 1997.

Roxithromycin is known as more stable than Erythromycin A under acid conditions like gastric environment. In this regards, we designed compounds to resist the strong acidic condition and to have excellent activity against H. Pylori-active. A series of erythromycin A derivatives were synthesized and tested for acid-resistant property. Biological activity against H. Pylori was evaluated.

[PD1-25] [ 04/21/2000 (Fri) 14:50 - 15:50 / [1st Fl, Bldg 3] ]

**Importance of phenyl moiety for cytotoxicity of 4-Phenyl-1-arylsulfonylimidazolidinones**

Kim JM, Kim IH, Jung SH, Whang HS'

College of Pharmacy, Chungnam National University, <sup>1</sup>Dong-Wha Pharm. Ind. Co. Ltd.

Novel 4-Phenyl-1-arylsulfonylimidazolidinones have been reported to show highly potent antitumor activity against the various cancer cell lines.

As a result of the structural modification of these compounds, the small aromatic moiety such as phenyl ring at 4-position of imidazolone ring had been identified as a structurally essential necessity for cytotoxicity.

However, the derivatives removed phenyl ring at 4-position have not been investigated. The corresponding compounds were synthesized and evaluated for their antitumor activity and compared to that of 4-phenyl compounds.

[PD1-26] [ 04/21/2000 (Fri) 14:50 - 15:50 / [1st Fl, Bldg 3] ]

**Synthesis and Antibacterial Activity of New Carbapenems Containing Isoxazole Moiety**

Kang Yong Koo, Shin Kye Jung, Yoo Kyung Ho, Kim Dong Chan, Seo Kyung Jae, Lee Kyung Seok, Park Ran Hee, Kim Dong Jin, Park Sang Woo

Korea Institute of Science and Technology

1 $\beta$ -Methylcarbapenems exhibit a broad antibacterial spectrum against both Gram-positive and Gram-negative organisms and high stability to dehydropeptidase-I (DHP-I). Meropenem, which has a 1 $\beta$ -methyl group in carbapenem nucleus, is stable to renal DHP-I and it has successively been launched on the market. In recent years, several analogues such as BO-2727, S-4661, ZD-4433,