

[PE1-5] [04/19/2001 (Thr) 15:30 – 16:30 / Hall 4]

Transdermal Controlled Delivery of Tenoxicam: 1. Effects of Vehicles and Enhancers

Gwak HS^o, Chun IK

College of Pharmacy, Dongduk Women's University, Seoul 136-714, Korea

Purpose: To evaluate the effect of various vehicles and penetration enhancers on the percutaneous absorption of tenoxicam, a non-steroidal anti-inflammatory drug of oxamic type.

Methods: Permeation studies were conducted using a side-by-side permeation system. Receptor compartments of cells were filled with 40% polyethylene glycol 400 (PEG 400) in normal saline and donor compartments were dosed with saturated solutions of tenoxicam in various vehicles. Vehicles used were water, ethanol, PEG 400, propylene glycol (PG), propylene glycol laurate, diethylene glycol monoethyl ether (DGME), propylene glycol monocaprylate (PGMC), propylene glycol monolaurate (PGML), oleyl alcohol (OAI), isopropyl myristate and propylene glycol dicaprylate/caprate. Cosolvents used were mixtures of DGME/PGMC, DGME/PGML and PG/OAI. As permeation enhancers, various fatty acids including oleic acid (OAc), linoleic acid (LOAc), capric acid, caprylic acid and lauric acid were added to some vehicles. Triethanolamine (TEA) or tromethamine was also added at 1 % concentration as a solubilizer. Samples (100 μ l) were taken over 24 hours and drug concentrations were quantitated using HPLC. Flux, lag time and amount permeated were calculated.

Results: Fluxes were low with neat vehicles (0.1–1.1 μ g/cm²/hr). In the effects of cosolvents, both of DGME/PGML and DGME/PGMC showed the highest flux of 20.3 \pm 8.0 and 13.5 \pm 8.0 μ g/cm²/hr at a 4:6 ratio, respectively. The cosolvents of PG/OAI increased the flux up to 100 times and 70 times at the ratios of 8:2 and 5:5, respectively, compared to OAI alone (0.3 \pm 0.1 μ g/cm²/hr). The addition of OAc and LOAc at 3% concentration to PG markedly increased the fluxes which were 41.8 \pm 4.0 and 28.5 \pm 5.0 μ g/cm²/hr, respectively, while the other fatty acids did not show any enhancing effects. Also, the addition of TEA or tromethamine, rather, decreased the flux when added to PG with OAc or LOAc in spite of improving the solubility of tenoxicam.

Conclusions: It was suggested that these results would be applied to the fabrication of tenoxicam transdermal delivery system.

[PE1-6] [04/19/2001 (Thr) 15:30 – 16:30 / Hall 4]

Development of novel thermosensitive and mucoadhesive diclofenac liquid suppository using sodium chloride

Jung SH^o, Yong CS¹, Rhee JD¹, Kim CK², Choi HG¹

¹ College of Pharmacy, Yeungnam University, 214-1, Dae-Dong, Gyongsan 712-749, South Korea

Liquid suppository system composed of poloxamers and bioadhesive polymers such as carbopol, polycarbophil and sodium alginate was easy to administer to the anus and mucoadhesive to the rectal tissues without leakage after the dose. However, a liquid suppository containing diclofenac sodium could not be developed using the bioadhesive polymers, since the drug was precipitated in this preparation. To solve this problem, it has been attempted to develop a novel thermosensitive and mucoadhesive liquid suppository system using sodium chloride instead of bioadhesive polymers. Poloxamer P 407 or/and P 188 were used to confer the thermosensitive gelation property. The mixtures of P 407 (15%) and P 188 (15–20%) existed as liquid at room temperature, but gelled at physiological temperatures. Diclofenac sodium significantly increased the gelation temperature and weakened the gel strength and bioadhesive force, while sodium chloride did the opposite. Furthermore, the diclofenac sodium liquid suppository with less than 1.0% of sodium chloride, in which the drug was not precipitated, was inserted into the rectum without difficulty and leakage and retained in the rectum for at least 6 h. These results suggested that the novel thermosensitive and

mucoadhesive liquid suppository system using sodium chloride could be further developed as a more convenient and effective rectal dosage form for diclofenac sodium.

[PE1-7] [04/19/2001 (Thr) 15:30 - 16:30 / Hall 4]

Synthesis and In vitro Evaluation of Diclofenac Prodrugs for Improved Dermal Delivery

Lee SH^o, Quan QZ, Rhee, JD, Choi HG, Yong CS

College of Pharmacy, Yeungnam University, 214-1 Gyongsan, Gyongbook

Synthesis and In vitro Evaluation of Diclofenac Prodrugs for Improved Dermal Delivery

Seung-Ho Lee*, Qi-Zhe Quan, Han-Gon Choi, Jong Dal Rhee, and Chul Soon Yong.

College of Pharmacy, Yeungnam University, 214-1 Gyongsan, Gyongbook

Purpose. To synthesize and evaluate alkyl esters of diclofenac as potential dermal prodrugs of diclofenac.

Methods. The prodrugs of diclofenac were synthesized, and their physicochemical properties such as solubilities, pKa's and stabilities in buffered solution and in hairless skin and liver were investigated. The permeation study of prodrugs across hairless mouse skin was carried out employing flow-through diffusion cell.

Results. The methyl- and ethyl ester prodrugs showed higher lipid solubilities in terms of octanol-buffer partition coefficients(log P_{apparent}) of 5.5 and 5.1 at pH7.0, respectively, when compared with diclofenac. They possessed moderate chemical stability in aqueous solutions of various pH's except strong acidic and basic conditions and were readily converted to diclofenac in hairless mouse skin and liver. The prodrugs showed a higher flux across the hairless mouse skin than diclofenac, with a maximum enhancement of 2.6-fold compared to diclofenac. They, however, showed shorter lag time than diclofenac did, and poor aqueous solubilities. They were ca. 1000 times more soluble in propylene glycol than in aqueous solution. Methyl- and ethyl ester prodrugs had the pKa of 6.9 and 7.2, respectively.

Conclusion. Alkyl ester prodrugs of diclofenac could be a potential candidate for improved dermal delivery of diclofenac.

[PE1-8] [04/19/2001 (Thr) 15:30 - 16:30 / Hall 4]

In Vivo Absorption of Poorly Soluble Drug According to Use of pH-Dependently Soluble Polymer as Carrier

Wang HS^o, Jang SW, Bae WT, Kwak HH, Kim JH, Kwon JW, Kim WB

Dong-A Research Laboratories

It has been examined the possibility of water insoluble drug into more soluble form using various polymers, which show pH-dependent solubility, i.e., entero-soluble or gastro-soluble polymers.

The final compositions manufactured using both kinds of polymers revealed very different dissolution profiles according to the pH of media as expected. In the case of using Eudragit L100 as an enteric polymer, the model drug was released in nearly intermediate pH much faster and more than the case of using gastro-soluble one, Eudragit E100. This was vice versa in the acidic pH, i.e., faster and more dissolution of latter composition.

According to the followed in vivo absorption experiment with rat, both compositions showed greatly different absorption behavior. Unlikely as expected, the composition containing gastro-soluble polymer revealed to show very reduced absorption, which was lower than that of extract itself. On the other hand, enteric polymer-containing composition had a good absorption behavior, much higher