

B13

**Production of the Wild Entomopathogenic Fungi,
Cordyceps militaris, in the Silkworm, *Bombyx mori***

Sang Mong Lee, Nam Sook Park¹, Sae Yun Cho², Jae Sam Hwang²
and Byung Rae Jin¹

*Department of Sericultural and Entomological Biology, Faculty of
Agriculture, Miryang National University, Miryang 627-702, Korea.*

¹*College of Natural Resources and Life Science, Dong-A University, Pusan
604-714, Korea.,* ²*Department of Sericulture and Entomology, National
Institute of Agricultural Science and Technology, R.D.A, Suwon 441-400,
Korea.*

The wild entomopathogenic fungi, *Cordyceps militaris*, were collected at the Whawang mountain, Korea. The pupae of the silkworm, *Bombyx mori*, were used as infecting hosts for the production of the *silkworm-militaris dongchunghacho*, silkworm vegetable wasps and plant worms with *C. militaris*. Three inoculation methods in terms of injection, spray and immersion were tested against the silkworm pupae. Three inoculation methods revealed 100% infectivity to the silkworm pupae tested. Of the three inoculation methods, the injection method was highly effective in the reduction of the period required for the endosclerotium and the completion of fruiting body formation. These results indicate that the silkworm pupae are very effective host insects for the production of *C. militaris*.