

P-48

**ACTIVATION OF PI3K IS NOT SUFFICIENT, BUT REQUIRED FOR  
H-Ras-INDUCED INVASIVE PHENOTYPE IN MCF10A CELLS**Il-Chung Shin and Aree Moon

College of Pharmacy, Duksung Womens University, Seoul 132-714

We have previously shown that H-ras, but N-ras, induces an invasiveness and cell motility in human breast epithelial cells (MCF10A), while both H-ras and N-ras induce transformed phenotype. It has been recently shown that phosphatidylinositol 3-kinase (PI3K) plays an important role on cell migration. In the present study, we wished to investigate the functional role of PI3K in H-ras-induced invasive phenotype in MCF10A cells. The activation of PI3K was examined by detecting phosphorylation of Akt, a downstream molecule of PI3K, by Western blot analysis. We show that phosphorylated Akt level was upregulated both in H-ras MCF10A cells and N-ras MCF10A cells comparing to the parental MCF10A cells while the amount of Akt was equal in the parental, H-ras- and N-ras MCF10A cells. The results suggest that activation of PI3K is not sufficient for invasiveness and motility since PI3K is also activated in the N-ras MCF10A cells which have been shown to be non-invasive and non-motile. We then further investigated the functional significance of PI3K activation in invasion and motility by using the known PI3K inhibitors, LY294002 and wortmannin. Treatment of LY294002 and wortmannin significantly reduced invasive phenotype and motility of H-ras MCF10A cells, suggesting that activation of PI3K is not sufficient, but may be required for H-ras-induced invasion and motility.