

P-51

**PREVENTIVE EFFECT OF MUSHROOM PHELLINUS LINTEUS ON
THE INHIBITION OF GAP JUNCTIONAL INTERCELLULAR
COMMUNICATION BY H₂O₂ IS INVOLVED IN THE
UP-REGULATION OF ERK2 AND p38**

Kyung-Sun Kang¹, Jong-Ho Cho¹, Sung-Dae Cho¹, Kyung-Bae Kim¹, Ji-Hae Lee¹,
Nam-Shik Ahn¹, Ji-Won Jung¹, Se-Ran Yang¹, Joon-Suk Park¹, Byung Su Yoon², Sung-hoon
Kim³, and Yong-Soon Lee¹

¹Department of Veterinary Public Health College of Veterinary Medicine, Brain Korea 21
agricultural biotechnology, Seoul National University 103 Seodun-Dong, Kwonsun-Ku, Su-won
441-744, Korea, ²Department of Biology, Kyunggi University, Su-won 442-760, Korea,
³Graduate School of East-west medical Science, Kyunghee University, Su-won 449-711, Korea
E-mail: leeys@snu.ac.kr. Fax: 031-292-7610

Gap junctional intercellular communication (GJIC) is a cellular event underlying the tumor promotion process and that treatment to prevent the down-regulation or to up-regulate GJIC is important in preventing tumor promotion. We evaluated the potential preventive effect of Mushroom Phellinus Linteus (PL) against the promoting action of hydrogen peroxide (H₂O₂) in WB-F344 rat liver epithelial cells. Cells were preincubated with PL (1-100 μ g/ml) for 24h followed by a stimulating treatment with PL and H₂O₂ (500 μ M) for 1h. Fluorescent dye (Lucifer Yellow) coupling between adjacent cells was evaluated by SL/DT (Scrape-Loading/Dye-Transfer). The distribution, quantity and phosphorylation pattern of connexin 43 were detected using immunofluorescence analysis and Western blotting. PL (5-100 μ g/ml) significantly prevented down regulation and hyperphosphorylation of connexin 43 by H₂O₂. Immunofluorescence analysis for connexin 43 demonstrated numerous punctate fluorescent spots along the intercellular plasma membrane in controls, which were significantly decreased by H₂O₂. PL prevented the decrease of Connexin 43. Interestingly, we also found that PL increased quantity of some MAPK (ERK2, p38). In conclusion, we suggest that PL might act as an

anti-tumor promoter via its ability of prevent down-regulation of GJIC through activation of ERK2 and p38.