

P-77

**SYNERGISTIC EFFECT OF HUMAN CYTOCHROME B5
COEXPRESSION ON THE METABOLIC ACTIVITY OF CYP1A2 IN
CHINESE HAMSTER OVARY CELLS**

Jin Sun Kang, Hyuck-Joon Kang, Mi-Sook Dong*, and Chang-Hwan Park

College of Medicine and Institute of Biomedical Sciences, Hanyang University and *Graduate
School of Biotechnology, Korea University, Seoul, Korea

E-mail: chshpark@hanyang.ac.kr

Human cytochrome B5 (CYB5) was coexpressed with cytochrome P450 1A2 (CYP1A2), NADPH-CYP450 reductase (CYPR) and *N*-acetyltransferase 2 (NAT2) in Chinese hamster ovary (CHO) cells. The expression of four proteins was determined by Western blot analyses. The introduction of cDNAs to CHO cells were transduced via retroviral vectors. The cytotoxicity assay of 2-aminoanthracene (2-AA) and aflatoxin B₁ were approximately 4-fold more sensitive than CYB5 free cells. But there were no difference treated with 2-amono-3,4-dimethylimidazo[4,5-*f*]quinoxaline (MeIQx) between CYB5 transduced and CYB5 free cells.