

F834 Genetic Relationship among Six Species of the Genus *Haliotis* by Random Amplified Polymorphic DNA (RAPD) Analysis

Hye-Suck An¹, Young-Ju Jee²,
Seock-Jung Han³, Sung-Real Park⁴,
Ho-Yong Ryu¹, Bong-Seok Kim¹ and
Bong-Lae Kim³

Biotechnology Division¹, Aquaculture Division²,
Cheju Marine Hatchery³, TongYeong Laboratory⁴,
National Fisheries Research & Development
Institute, Pusan 619-900

The RAPD technique was used to identify genetic relationships among six species of the genus *Haliotis* distributed in Korea. A dendrogram was constructed using UPGMA from the polymorphic patterns generated by RAPD profiles. The molecular data clustered into two groups. Cluster I included *Haliotis discus hannai*, *H. discus*, *H. madaka* and *H. gigantea*, which was subsequently divided into two subclusters. Subcluster I included *Haliotis discus hannai*, *H. discus* and *H. madaka*, subcluster II with *H. gigantea*. Cluster II contained *H. diversicolor supertexta* and *H. diversicolor diversicolor*. The RAPD markers were found to be a useful tool for detecting genetic relationship within the six species of the genus *Haliotis*.

F835 Population Genetic Data on the Thirteen CODIS Short Tandem Repeat Loci in Koreans

Myun-Soo Han¹, Pil-Won Kang¹,
Sang-Kyu Choi¹, Youl-Hey Cho², Han-Jun
Jin³, Kyoung-Don Kwak³ and Wook Kim^{3*}

¹DNA Analysis Section, National Institute of
Scientific Investigation, Seoul 158-097, Korea

²Department of Genetics, Hanyang University
School of Medicine, Seoul 133-791, Korea;

³Department of Biology, College of Advanced
Sciences, Dankook University, Cheonan 330-714,
Korea

We analyzed variations at thirteen
Combined DNA Index System (CODIS)
short tandem repeat (STR) loci (CSF1PO,

FGA, TH01, TPOX, vWA, D3S1358,
D5S818, D7S820, D8S1179, D13S317,
D16S539, D18S51, and D21S11) in a sample
from 130 unrelated individuals in the
Korean population. Allele and genotype
frequencies were determined with
commercial PCR-based DNA profiling kits.
The Exact Test demonstrated that all loci
were found to be no deviations from
Hardy-Weinberg expectations ($P > 0.05$). For
forensic testing, the discriminating powers
(PD) were 0.866 for CSF1PO, 0.961 for
FGA, 0.826 for TH01, 0.760 for TPOX,
0.931 for vWA, 0.863 for D3S1358, 0.909 for
D5S818, 0.904 for D7S820, 0.948 for
D8S1179, 0.930 for D13S317, 0.915 for
D16S539, 0.958 for D18S51, and 0.917 for
D21S11, respectively (combined PD:
0.9999999999999998). Therefore, the Korean 13
CODIS STR data could be useful for the
regional specific and prerequisite references
to the forensic community.

F836 Requirement of Mediator complex for gene-specific transcriptional activation during *Drosophila* development.

Young-Joon Kim, Jin Mo Park, Byeong
Soo Gim

Yonsei University Department of Biochemistry
National Creative Research Center for Genome
Regulation

Mediator of transcriptional regulation is the
evolutionary conserved coactivator complex
that plays the central role in the integration
and recruitment of diverse regulatory
signals and transcription machinery to
certain promoters. In yeast, each Mediator
subunit is required for transcriptional
regulation of a distinct group of genes. In
order to decipher the mechanistic roles of
Mediator proteins in regulating
developmental specific gene expression, we
isolated, and analyzed a multiprotein
complex containing *Drosophila* Mediator
homologs (dMediator). dMediator interacts
with several sequence-specific transcription
factors and basal transcription machinery,

and is critical for activated transcription in response to diverse transcriptional activators. In order to elucidate the function of Mediator in metazoan development, we isolated mutants of a conserved Mediator subunit, *Drosophila* Med6 (dMed6). dMed6 null homozygotes failed to pupate and died in the third larval instar. Larval mitotic cells and most imaginal discs showed severe defects in proliferation, but no apparent morphological defect was observed in other larval tissues. Clonal analysis of dMed6 mutant cells revealed that dMed6 is essential for cell viability and proliferation of most adult cell types. *Drosophila* cDNA microarray, quantitative RT-PCR, and in situ expression analyses of developmentally regulated genes in dMed6 mutants showed that transcriptional activation of a subset of genes involved in neuroblast proliferation in the larval brain were most affected. Our results suggest that dMed6 is required in most cells for transcriptional regulation of a subset of genes important for cell proliferation and metabolism.

F837 The chromosomal study of native plants in Korea

Jae-Wook Bang^{*}, Soo-Young Kim, Joong-Yeon Cho, Woo-Ku Lee, Mi-Na Yu, June-Young Kim, Mi-Sung Jeon and Jee-Young Park

Dept. of Biology, Chungnam National University, 220 Gung-Dong, Yousung-gu, Daejeon 305-764, Korea

There are about 90 families and 3600 species of native plants in Korea. In this study, chromosomes of 18 families and 46 species were observed by Feulgen staining. The numbers in the family Compositae chromosome were observed diversely. The somatic chromosome number of *Carduus crispus* was $2n=16$, *Lactuca sativa* was $2n=18$, *Atractylodes japonica* and *A. ovata* were $2n=24$, *Matricaria chamomilla* was $2n=28$, *Taraxacum coreanum* was $2n=32$, *Arctium lappa* and *Achillea sibirica* were $2n=36$ and *Aster ageratoides* was $2n=72$.

The chromosome numbers of 4 species in the family Umbelliferae were $2n=22$. The other families and species also have different chromosome numbers.

F838 Construction of transgenic silkworm using P element based expression vector in *Bombyx mori*.

Soonjeung Kim^{*}, Seunghyun Sung, Ji-yeon Sung, Seungshic Yum and Dongsang Suh.

Department of Genetic engineering, SungKyunKwan Univ. 440-746 Suwon, Korea.

Transgenesis is a powerful method of studying the role and expression mechanism of genes and organism. It is also a way to confer genetically useful characteristics to animals and plants that can be used in biotechnological applications. Introducing new genes into silkworms has proved difficult, but we have developed an efficient method of transgenesis for the silkworm *Bombyx mori*. The method makes use of the microinjection technique and P-derived vector to transfer the foreign genes into the chromosomes. We constructed the expression vector using fibroin gene promoter and P transposon vector containing luciferase as reporter genes (pFpLuc). We microinjected into eggs laid at the preblastoderm stage. 29 of 6815 microinjected eggs were survived. After PCR analysis, 3 of them were tured out transgenic silkworms. Also, F1 were assayed by PCR. We assayed F2 and F5 transgenic silkworms and got the positive PCR results and did PCR-sequencing. As for ClustalW result, PCR products had the sequence of luciferase. The studies on the gene expression using fibroin gene promoter may help to understand mechanism in fibroin genes, i.e. transcriptional regulation, or many advantages to produce useful biological materials. Transgenic silkworm technique will be very useful for basic research of silkworm and may be used for the massive production of proteins for diagnostic and therapeutic purposes.