

acid. There are two isoforms of COX, designated as COX-1 and COX-2. While COX-1 is constitutively expressed in most tissues, COX-2 can be induced transiently by proinflammatory cytokines, endotoxins, growth factors, oncogenes, UV and mitogens. Elevated levels of COX-2 have been observed in cancers of breast, colon, and lung as compared with the surrounding normal tissues. Based on these findings, it is conceivable that targeted inhibition of inappropriate or abnormal up-regulation of COX-2 is one of the most broadly effective and promising approaches to cancer chemoprevention. Celecoxib, a selective COX-2 inhibitor, has been reported to prevent experimentally induced colon, breast and skin carcinogenesis. Moreover, daily intake of celecoxib resulted in significant reduction of polyps in patients with familial adenomatous polyposis. In the present study, we examined the effect of celecoxib on COX-2 induction in 12-*O*-tetradecanoylphorbol-13-acetate (TPA)-treated mouse skin. Topical application of 0.1, 1.0, or 10 μ mole celecoxib onto shaven backs of female ICR mice (6 to 7 wk of age) 30 min prior to 10 nmole TPA inhibited expression of COX-2 protein and subsequent production of prostaglandin E₂ in a dose-related manner. Celecoxib inhibited activation of AP-1 and C/EBP transcription factors in a dose dependent manner. Furthermore, celecoxib inhibited both catalytic activity and phosphorylation of ERK1/2. These results suggest that celecoxib suppresses TPA-induced COX-2 expression in mouse skin by blocking activation of ERK, which appears to be mediated by transcription factors such as AP-1 and C/EBP.

[OC-2] [04/19/2002 (Fri) 14:50 - 15:00 / Hall A]

BETA-AMYLOID INDUCES OXIDATIVE DNA DAMAGE AND CELL DEATH: POSSIBLE INVOLVEMENT OF INFLAMMATORY CASCADES

Jang JungHee^o, Surh YoungJoon

College of Pharmacy, Seoul National University, Korea

Oxidative stress induced by reactive oxygen and/or nitrogen species has been considered as a major cause of cellular injuries in a variety of neurodegenerative disorders including Alzheimer's disease (AD). Inflammatory as well as oxidative tissue damage has been implicated in pathophysiology of AD, and non-steroidal anti-inflammatory drugs have been reported to have beneficial effects in the treatment or prevention of AD. In this study, we have investigated the molecular mechanisms underlying oxidative and inflammatory cell death induced by beta-amyloid, a neurotoxic peptide associated with senile plaques formed in the brains of patients with AD. Rat pheochromocytoma (PC12) cells treated with beta-amyloid exhibited increased intracellular accumulation of reactive oxygen species and underwent apoptotic death as determined by characteristic morphological features, internucleosomal DNA fragmentation and positive in situ terminal end-labeling (TUNEL staining). beta-Amyloid treatment also led to the cleavage of poly(ADP-ribose) polymerase, the increased Bax/Bcl-XL ratio and the decreased mitochondrial membrane potential in PC12 cells. Furthermore, transfection of PC12 cells with *bcl-2* rescued these cells from apoptotic death induced by beta-amyloid. beta-Amyloid caused activation of NF- κ B and AP-1, which appeared to be preceded by activation of mitogen-activated protein kinases (MAPKs), such as extracellular signal-regulated kinase 1/2 (ERK1/2), p38 MAPK and c-Jun N-terminal kinase/stress-activated protein kinase. Exposure of PC12 cells to beta-amyloid resulted in time-dependent induction of cyclooxygenase-2 (COX-2) and inducible nitric oxide synthase. beta-Amyloid-induced cell death was attenuated by pretreatment with the selective COX-2 inhibitor celecoxib or the peroxynitrite scavenger ergothioneine. These results suggest that prooxidative and proinflammatory mechanisms are involved in the oxidative and/or nitritative cell death in mediating the neurodegeneration associated with AD.

[OD-1] [04/19/2002 (Fri) 15:00 - 15:10 / Hall A]

Cytotoxic Terpenoids from the Sponge *Sarcotragus* Species

Liu Yonghong^o, Hong Jongki, Lee Chong-O, Im Kwang Sik, Kim Nam Deuk, Choi Jae Sue, Jung Jee Hyung

College of Pharmacy, Pusan National University, Pusan 609-735, Korea, Korea Basic Science Institute, Seoul, Korea, Korea Research Institute of Chemical Technology, Taejon, Korea, and Pukyung National University, Pusan, Korea

Twenty-one new and two known terpenoids were isolated from the marine sponge *Sarcotragus* sp., including twelve furanosesterpene and their derivatives, seven new pyrrolosesterterpenes two trinosesterterpenes, and two diterpenes were isolated from the same sponge by bioactivity-guided

fractionation. The gross structures were established based on NMR and MS analysis. The stereochemistry was defined by combined use of NMR, CD spectroscopy, or chemical degradation, optical rotation. The compounds were evaluated for cytotoxicity against five human tumor cell lines to exhibit moderate to significant activity.

[OD-2] [04/19/2002 (Fri) 15:10 - 15:20 / Hall A]

Curtisians block the neurodegeneration by neurotoxin through iron chelation

Lee In-Kyoung⁰, Yun Bong-Sik, Kim Young-Ho¹, Yoo Ick-Dong

Korea Research Institute of Bioscience and Biotechnology, ¹College of Pharmacy, Chungnam National University

Among the neuroprotective approach for stroke treatment, calcium-induced excitotoxicity was widely accepted as a key event after cerebral ischemia. Activation of excitatory receptors by glutamate increase Ca^{2+} entry into neurons and accumulation of the Ca^{2+} into mitochondria followed oxidative damage leads to the cell death. To select neuroprotective agents, we concentrated on antioxidants with lipid peroxidation inhibitory activity. In our continuing investigation for lipid peroxidation inhibitors from mushroom extracts, we have isolated *p*-terphenyls named curtisians from *Paxillus curtisii*. Curtisians showed very significant lipid peroxidation inhibition activity, especially in lipid peroxidative assay system used iron as oxidant, comparing with superoxide and DPPH radical scavenging effects. It suggested that curtisians were implicated in iron chelating properties.

In the present study, we have investigated the mode of action of curtisians including superoxide and DPPH radical scavenging activity, neuronal cell protective activity in mouse cortical cell cultures, and iron-chelating activity using DNA single strand breakage method. The neuroprotective effect of iron chelator desferrioxamine in the hypoxia-ischemic immature mouse brain recently reported supported the fact that iron chelation was a prominent effect of curtisians. In conclusion, our results suggested that neuroprotective activity of curtisians would act as an iron chelator but not as a free radical scavenger.

[OD-3] [04/19/2002 (Fri) 15:20 - 15:30 / Hall A]

Diagnostic Patterns of Capillary Electrophoretic Urinary Nucleoside Profiles from Patients with Thyroid cancer

La Sookie⁰, Cho JungHwan, Kim Kyoung-Rae, Jung Han Kim

College of Pharmacy, Sungkyunkwan University, College of Pharmacy, Sookmyung Womens University, Department of Biotechnology, College of Engineering and Bioproducts Research Center, Yonsei University

An efficient capillary electrophoretic profiling method in micellar electrokinetic capillary chromatography (MEKC) mode was combined with simple pattern recognition methods for the correlation between urinary nucleosides and thyroid cancer. For the urinary nucleoside profiles, free nucleosides were extracted by SPE in affinity mode with the subsequent analysis by MEKC. A total of 15 nucleosides were positively identified urine samples from patients with thyroid cancer and normal subjects studied. Multivariate statistical analyses appropriate for the correlation between urinary nucleosides profiles and thyroid cancer was investigated. The present nucleoside profiling and simple pattern recognition methods appear to be useful for the comparative analysis of urinary nucleosides among groups of normal subjects and patients with thyroid cancer.