

Purpose: It is generally accepted the fact that pharmacological activity would be different in terms of the method of extraction. Antidiabetic activities between SP water extract and 70% ethanol extract were compared in streptozotocin (STZ)-induced diabetic rats.

Experimental methods: SP water or ethanol extract, 1 g/kg each, was administered orally 3 days before intraperitoneal injection of STZ (Day 0). Until Day 5, 20 mg/kg of STZ, which is dissolved in 100 mM citrate buffer (pH 4.5), was injected once a day. SP extracts were administered for 3-week period, and plasma glucose level and body weight change were determined periodically. At the end of the treatment, plasma levels of total cholesterol, triglyceride and free fatty acid were investigated. Urinary glucose and albumin excretion levels were also determined at 2 and 3 weeks. Kidney hypertrophy and mesangium expansion were compared between groups using PAS staining and immunohistochemical analyses.

Results: SP water- and ethanol extract-treated groups both significantly reduced the plasma glucose levels at 3rd week as compared with the diabetic control group. Water intakes in SP water- and ethanol treated-groups were also improved as compared with the diabetic control group. SP ethanol extract treated group showed more potent kidney hypertrophy protection activity than water extract-treated group. SP ethanol extract treated group also had more urine albumin and glucose lowering activity than water extract treated group.

Taken together, we may conclude that SP ethanol extract treated group showed more potent antidiabetic activity in STZ-induced diabetic rats than water extract treated group.

[PA1-42] [ 04/18/2002 (Thr) 14:00 - 17:00 / Hall E ]

#### Antiangiogenic activities of isoflavonoids isolated from the rhizomes of *Belamcanda chinensis*

Jung SangHoon<sup>o</sup>, Lee YeonSil, Lee SangHyun, Lim SoonSung, Kim YeongShik, Shin KukHyun

Natural Products Research Institute, Seoul National University

Many evidences proved the correlation between angiogenesis and tumor growth, therefore, searching for natural or synthetic angiogenic inhibitors as potential anti-cancer drugs has been extensively carried out, and it has recently been reported that COX-2 induces angiogenesis, which is essential for tumor growth. Increased prostaglandin production and enhanced release of angiogenic growth factor by COX-2 may induce neovascularization.

In the previous study, we reported that tectoridin and tectorigenin, two isoflavonoids, isolated from the rhizomes of *Belamcanda chinensis* inhibited PGE<sub>2</sub> production in TPA- or thapsigargin stimulated rat peritoneal macrophages by inhibiting the induction of COX-2 protein.

Based on these results, the present study was carried out to clarify whether tectorigenin and tectoridin, inhibit angiogenesis by experimental method *in vitro* and *in vivo*. As a results, both compounds were found to exhibit a significant antiangiogenic activity when measured by chorioallantoic membrane (CAM) assay and by its effect on the proliferation of calf pulmonary arterial endothelial cells. Both compounds were also found to possess matrix metalloproteinase inhibitory activity *in vitro*. *In vivo*, a matrigel plug assay showed that both compounds suppressed basic fibroblast growth factor (bFGF)-stimulated angiogenesis and lowered the hemoglobin content inside the plug.

From these experimental data, it is suggested that tectorigenin is attributed to be a promising compound for the prevention and/or treatment of angiogenesis.

[PA1-43] [ 04/18/2002 (Thr) 14:00 - 17:00 / Hall E ]

#### Isolation of Aldose Reductase Inhibitors from the Rhizomes of *Belamcanda chinensis*

Jung SangHoon<sup>o</sup>, Lee YeonSil, Lee SangHyun, Lim SoonSung, Kim YeongShik, Shin KukHyun

Natural Products Research Institute, Seoul National University

Aldose reductase, the key enzyme of the polyol pathway, is known to play important roles in the diabetic complication. The inhibitors of aldose reductase, therefore, would be a most promising drug for the