

## BandW : The 2-DE image analysis software using MATLAB

Park Dae Ui<sup>o</sup>, Kim Chul Hong, Hong Seong Eui, Jung An Seong, Chang Gregory Y., Jung Hae Young

Pusan Bioinformatics and Biocomplexity Research Center, Pusan National University, Pusan, Korea. College of Pharmacy, National University, Pusan, Korea. Dept. of Neurology, University of Southern California, Los Angeles, USA

Since genomics represents only the first step in understanding cellular physiology, it needs to be complemented by systematic analysis of the proteins, termed proteomics. The primary means studying proteomics has been with use of 2-dimensional electrophoresis (2DE) since 1975. 2DE initially separate each protein according to its electric charge content by using isoelectric focusing and further separation of similar charged proteins are accomplished by using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). But Analyzing two-dimensional gel electrophoresis (2DE) is complicated and time-consuming. In order to facilitate the process, we compiled "BandW" which is an image analysis tool made using MATLAB (Matrix oriented computing engine). BandW analyzes the input image using M by N matrix which has a value between 0 to 255. Each value represents a pixel intensity. Image output is processed to TIF (Tagged Indexed Fileformat) format. BandW has simplified layout which utilizes image processing toolbox and graphic user interface engine embedded in MATLAB 5.3. During analysis, spot volume is calculated using contour algorithm. Volume calculation starts with combining outer boundary of same pixel intensity in the adjacent matrix until the circular loop is completed. The area within the loop actually represents a three-dimensional volume. We tested BandW by analyzing 2DE in liver mitochondria of young (LCY) or liver mitochondria of old (LCO) rats. Our results, using BandW, were identical to values obtained from available commercial softwares. In addition, BandW is 3D capable as only one other highly priced software.

[PC1-50] [ 04/18/2002 (Thr) 14:00 - 17:00 / Hall E ]

### Activation of PI3K is not Sufficient but Required for H-ras-induced Invasive Phenotype of MCF10A Cells

Shin Il-Chung<sup>o</sup>, Moon Aree

College of Pharmacy, Duksung Women's University, Seoul, Korea

We have previously shown that H-ras, but N-ras, induces an invasiveness and cell motility in human breast epithelial cells (MCF10A), while both H-ras and N-ras induce transformed phenotype. It has been recently shown that phosphatidylinositol 3-kinase (PI3K) plays an important role on cell migration. In the present study, we wished to investigate the functional role of PI3K in H-ras-induced invasive phenotype in MCF10A cells. The activation of PI3K was examined by detecting phosphorylation of Akt, a downstream molecule of PI3K, by Western blot analysis. We show that phosphorylated Akt level was upregulated both in H-ras MCF10A cells and N-ras MCF10A cells comparing to the parental MCF10A cells while the amount of Akt was equal in the parental, H-ras and N-ras MCF10A cells. The data indicate that activation of PI3K is not sufficient for invasiveness and motility since PI3K is also activated in the non-invasive and non-motile N-ras MCF10A cells. We investigated the functional significance of PI3K activation in invasion and motility by using PI3K inhibitors, LY294002 and wortmannin. Treatment of LY294002 and wortmannin significantly reduced invasive phenotype and motility of H-ras MCF10A cells, suggesting the requirement of PI3K activation for H-ras-induced invasion and motility. We then examined the effect of the PI3K inhibitors on matrix metalloproteinase (MMP) expression. Treatment of LY294002 inhibited secretion of MMP-2 and MMP-9 in a dose-dependent manner while wortmannin did not affect MMP levels in H-ras MCF10A cells. The possible role of Rac1 in H-ras-induced invasive phenotype in MCF10A cells are currently under investigation.

[PC1-51] [ 04/18/2002 (Thr) 14:00 - 17:00 / Hall E ]

### Enhancement of Proliferation and Migration of Glioma Cells by Glial Cell-derived Neurotrophic Factor (GDNF) for the Development of an Artificial Nerve Tubing

Song Hyun<sup>o</sup>, Dong June Chung<sup>1</sup>, Pill-Hoon Choung<sup>2</sup>, Aree Moon

College of Pharmacy, Duksung Women's University, <sup>1</sup>Sungkyunkwan University, <sup>2</sup>Seoul National University

In an attempt to provide useful information on the development of an artificial nerve tubing, proliferative and migrative properties of two glioma cell lines, C6 rat glioma cells and Hs683 human glioma cells, were examined. The present study shows that C6 cells proliferated more rapidly than Hs683 cells. The Hs683 cells are more adequate for the development of nerve tubing since unlike C6 cells, they are of human origin and known to be non-tumorigenic. In order to enhance proliferative and migrative abilities of Hs683 cells for the application as an artificial nerve tubing, we studied the effect of glial cell-derived neurotrophic factor (GDNF) on C6 and Hs683 cells. GDNF increased proliferation and migration of Hs683 cells in a dose-dependent manner. As an approach to develop artificial nerve tubing, we wished to determine if GDNF stimulate proliferation of glioma cells in the scaffolds. Cells were seeded in the scaffolds (polymer constructs), fabricated with type I collagen and alginate modified with cinnamoyl moiety, in the presence or absence of GDNF. Compared to control, cell proliferation was greatly enhanced by GDNF treatment of scaffolds as evidenced by staining of the cells in paraffin block. We then tested cytotoxicity of scaffolds used in this study. Hs683 cell growth was not inhibited by scaffold, proving that scaffold is not cytotoxic. Taken together, we show that GDNF treatment of scaffolds effectively increased Hs683 cell proliferation, suggesting a possible use of GDNF for developing artificial nerve tubing.

Poster Presentations – Field C2. Microbiology

[PC2-1] [ 04/18/2002 (Thr) 14:00 – 17:00 / Hall E ]

Isolation, Identification and Characterization of vancomycin-resistant *Streptococcus equinus* from raw milk sample

Lee JinWoo<sup>o</sup>, Choi SungSook, Kim KyungJae, Ha NamJoo

<sup>1</sup>Dept of Pharmacy, Sahmyook University, Seoul 139-742, Korea <sup>2</sup>Dept. of Food Science, Sahmyook College

To determine the occurrence of vancomycin-resistant *Enterococcus* and *Streptococcus* in raw milk samples, we examined raw milk samples for three month. Resistant strains were isolated directly from *Enterococcal* selective agar plates supplemented with 2mg of vancomycin per liter. 6 strains having high resistance were isolated. 5 of six were identified as *Streptococcus equinus* and 1 of 6 was identified as *Enterococcus faecium*.

To determined resistance, 6 isolates were tested with vancomycin and teicoplanin. Vancomycin resistant were genotyped by PCR analysis and *Enterococcus faecium* was VanC type

[PC2-2] [ 04/18/2002 (Thr) 14:00 – 17:00 / Hall E ]

Antihyperlipidemic Effect of *Alpinia officinarum*

Shin JeEun<sup>o</sup>, Kim DongHyun, Han Myungjoo

Department of Food and Nutrition, and College of Pharmacy, Kyung Hee University

The inhibition of lipase improves the condition of hyperlipidemia, obesity, hypertension, atherosclerosis and many other cardiovascular diseases. Therefore, the objective of this study was to investigate antihyperlipidemic effects of functional food ingredients.

*Cinnamomum cassia*, *Rheum palmatum*, *Alpinia officinarum* and *Chrysanthemum indicum* were selected as the the potent lipase inhibitor in vitro. However, *Cinnamomum cassia*, *Rheum palmatum* and *Chrysanthemum indicum* showed no significant antihyperlipidemic activity in high cholesterol diet induced hyperlipidemic mice. However, *Alpinia officinarum* improved serum TG, HDL and LDL level in corn oil