

The roots of *Juglans mandshurica* has been used as a folk medicine for treatment of cancer in Korea. In the course of isolating cytotoxic compounds from this plant, we isolated two new and two known diarylheptanoids along with one known sesquiterpenoid and their structures were elucidated on the basis of spectroscopic studies. Four of these compounds exhibited moderate cytotoxicities in ranges of IC₅₀ from 2 to 25 µg/ml against human colon carcinoma and human lung carcinoma cell lines.

[PD2-14] [04/19/2002 (Fri) 10:00 – 13:00 / Hall E]

Constituents from the roots of *Hemerocallis fulva*

Kim JuSun^o, Kang SamSik, Son KunHo, Chang HyeunWook, Kim HyunPyo, Bae KiHwan

Natural Products Research Institute, Seoul National University, 1Department of Food and Nutrition, Andong National University, College of Pharmacy, 2Yeongnam University, 3Kangwon National University and 4Chungnam National University

Besides chrysophanol and friedelin, five mixtures of *n*-hydrocarbons [pentacosane(72.6%), heptacosane (14.6%), tetracosane(5.8%), nonacosane(4.1%) and hexacosane(2.9%)], *n*-hydrocarbon alcohols [octacosanol(70.5%) and hexacosanol(29.5%)], 1-monoacyl glycerols [acyl part, behenic acid(43.5%), lignoceric acid(32.4%), cerotic acid (9.3%), tricosanoic acid(8.9%), pentacosanoic acid(2.6%), octacosanoic acid(2.3%), heneicosanoic acid(1.0%)], waxes [behenic acid(56.3%); lignoceric acid(23.0%) cerotic acid(19.8%), tricosanoic acid(4.6%), octacosanoic acid(4.0%), pentacosanoic acid(1.7%), triacontanoic acid(0.6%)/ octacosanol(33.7%), hexacosanol(21.0%), tetracosanol(15.6%), triacontanol (10.5%); docosanol(6.0%), tricosanol(6.0%), heptacosanol(4.2%), nonacosanol(3.0%)] and sterols [β -sitosterol(73.2%), stigmasterol(14.6%), campesterol(12.2%)] were isolated from the roots of *Hemerocallis fulva*. A mixture of 1-monoacyl glycerols is the first isolation from this plant. All compounds were identified on the basis of spectral data and chemical reactions.

[PD2-15] [04/19/2002 (Fri) 10:00 – 13:00 / Hall E]

Two new non-glycosidic iridoids from *Patrinia saniculaefolia*

An RenBo^o, Na MinKyun, Jin WenYi, Lee HyunYong, Bui Trong Dat, Kim YongHo, Bae KiHwan

College of Pharmacy, Chungnam National University, Taejeon 305-764, Korea

Patrinia saniculaefolia Hemsley (Valerianaceae) is an endemic species in Korea. The whole plant was extracted with methanol, then suspended in H₂O and successively partitioned with hexane, CH₂Cl₂ and *n*-BuOH. Repeated column chromatography of the hexane soluble fraction afforded two new non-glycosidic iridoids. On the basis of ¹H, ¹³C-NMR, HMQC, HMBC and ¹H-¹H ROESY spectral data, their structures were established as butanoic acid, 3-methyl-1-[(1R,3R,5S,7aS)-1, 3, 5, 7a-tetrahydro-3, 5-dimethoxy-7-(hydroxymethyl)-1-(3-methyl-1-oxo-but-oxy)cyclopenta[c]pyran-4-yl]methyl ester(1) and butanoic acid 3-methyl-1-[(1R,3R,5R,7aS)-1, 3, 5, 7a-tetrahydro-3, 5-dimethoxy-7-(hydroxymethyl)-1-(3-methyl-1-oxo-but-oxy)cyclopenta[c]pyran-4-yl]methyl ester(2), which were named patridoid I and patridoid II, respectively.

[PD2-16] [04/19/2002 (Fri) 10:00 – 13:00 / Hall E]

Cerebrosides and Triterpene Glycosides from the root of *Aster scaber*

Kwon HakCheol^o, Choi SangZin, Min YongDeuk, Lee SungOk, Lee WonBin, Yang MinCheol, Chung AeKyung, Lee KyuHa, Nam JungHwan, Lee JongSoon, Lee KangRo

Natural Products Laboratory, College of Pharmacy, SungKyunKwan University, Suwon 440-746, Korea and Department of Biochemistry, YeungNam University, GyongSan, 712-749, Korea.