

inhibitory effect against MAO in a dose dependant manner with the IC50 value of 4 µg, and inhibited both MAO-A and B with the IC50 value of 4 µg and 3 µg respectively. Compound 1 was found to be competitive MAO inhibitor.

[PD2-45] [04/19/2002 (Fri) 10:00 – 13:00 / Hall E]

Phenolic compounds from Needles of *Pinus densiflora* and Their Cytotoxic Activities on Mouse Melanoma Cell line

Kim JeeHun^o, Kwon YoungMin, Choi JongMoon, Yeom SeungHwan, Lee JaeHee, Lee MinWon

Choong-Ang University

Phytochemical examination of needles of *Pinus densiflora* isolated eight phenolic compounds and the cytotoxic activity of these compounds on mouse melanoma cell line were evaluated by 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) colorimetric method. Several compounds showed significant cytotoxic activity. These result suggested that some phenolic compounds from needles of *Pinus densiflora* might be developed to anti-cancer agent.

[PD2-46] [04/19/2002 (Fri) 10:00 – 13:00 / Hall E]

Inhibitory effects of medicinal herbs on cytochrome P450 drug metabolizing enzymes

Jeong Hye Gwang, You Ho Jin¹, Chang Young-Su^o, Park Sung Jun, Moon Young Hee, Woo Eun-Rhan

College of Pharmacy, ¹College of Medicine, Chosun Univ., Kwangju 501-759, Korea

The MeOH ext., CH₂Cl₂Frac., EtOAc Frac., n-BuOH Frac., and H₂O Frac. of 23 Korean medicinal herbs were prepared and were tested the inhibitory effects on Cytochrome P450 (Cyp) 1A1/2, 2B1/2, 2E1. Among the tested samples, the extracts of *Selaginella tamariscina*, *Euonymus alatus*, *Salvia miltiorhiza*, *Angelica acutiloba*, *Rheum palmatum*, *Paeonia moutan*, *Scutellaria barbata*, *Tribulus terrestris*, *Hedyotis diffusa*, *Curcuma zedoaria*, *Rehmania glutinosa*, *Trogopterus xanthipes*, *Melandryum firmum*, *Achyranthes bidentata*, *Leonurus sibiricus*, *Panax ginseng*, *Paeonia lactiflora*, *Poncirus trifoliata*, *Cnidium officinale*, *Cyperus rotundus*, *Corydalis ternata* showed significant inhibitory effects on Cyp 1A1/2, 2B1/2, 2E1. The IC₅₀ values of those extracts were found to be below 50 µg/ml.

[PD2-47] [04/19/2002 (Fri) 10:00 – 13:00 / Hall E]

Effect of *Panax ginseng* head butanol fraction on collagen-induced arthritis in DBA/1J mice

Jeong ChoonSik^o, Hyun JinEe, Suh InOk, Cho SoYean, Kim YeongShik

¹College of Pharmacy, Duksung Women's University, ²Natural Products Research Institute, Seoul National University, ³Korea Food and Drug Administration

Head of *Panax ginseng* C. A. Meyer indicates its growth number of years and it has been widely used for supplying energy to weak person. In the previous study, we reported that butanol fraction of *Panax ginseng* head not only has antigastric and anti-ulcerative properties but also showed anti-inflammatory activity. It is widely known that arthritis has relevance to inflammatory, thus we inclined to investigate the effect of *Panax ginseng* head butanol fraction on arthritis animal model. Collagen-induced arthritis is recognized as an in vivo tool in researching the mechanism of RA. Male DBA/1J mice, aged 5-6 weeks, were treated under the intradermally with bovine type II collagen emulsified in Freund's complete adjuvant, and a booster injection was given under the same conditions on the 21th day. Butanol fraction of *Panax ginseng* head showed significant inhibition on hind paw edema test and anti-