

Modulation of carcinogen-activating enzymes by synthetic trans-stilbene analogs

Lee SangKwang^O Kim SangHee Kim MieYoung Chun YoungJin

College of Pharmacy, Chungang University, Seoul, 156-756, Korea and Natural Products Research Institute, Seoul National University, Seoul, 110-460, Korea

Previous studies have demonstrated that 2,3',4,5'-tetramethoxystilbene (TMS) and 3,3',4',5,5'-pentamethoxystilbene (PMS) showed selective inhibition of human cytochrome P450 1B1 and 1A1 in vitro, respectively. In the present study, the effects of synthetic stilbene analogs on the expression of cytochrome P450 1A1 or 1B1 were investigated in human tumor cell lines such as HepG2, MCF-7 and MCF-10A. TCDD caused a dramatic increase in the amount of P450 1A1 or 1B1 proteins and mRNA levels. TMS suppressed TCDD-induced P450 1B1 expression in MCF-7 and MCF-10A cells in a dose-dependent manner. PMS also showed a dose-dependent decrease in P450 1A1 protein and mRNA levels induced by TCDD in HepG2 cells. The cytotoxic effects of these analogs were determined in these cell lines. Taken together, our results indicate that PMS and TMS are strong modulators of P450s gene expression as well as potent and selective inhibitors of P450 1A1 or 1B1, respectively. The detailed mechanisms of suppression of P450s gene expression needs to be determined.

[PC1-2] [10/17/2002 (Thr) 13:30 - 16:30 / Hall C]

Submicrosecond dynamics of nucleic acids studied with a long-lifetime metal-ligand complex

Kang Jung Sook^O, ‡Son Woo-Sung, §Kostov Yordan

Dept. of Oral Biochem. & ‡Orthodontics, Col. of Dentistry, Pusan Nat'l Univ., Pusan 602-739, Korea; §Dept. of Chem. & Biochem. Eng., Univ. of Maryland Baltimore County, Baltimore, MD 21225, USA

The metal-ligand complex, $[\text{Ru}(\text{phen})_2(\text{dppz})]^{2+}$ (phen = 1,10-phenanthroline, dppz = dipyrro[3,2-a:2',3'-c]phenazine) (RuPD), was used as a spectroscopic probe for studying nucleic acid dynamics. The RuPD complex displays a long lifetime and a molecular light switch property upon DNA binding due to shielding of its dppz ligand from water. To show the usefulness of this luminophore (RuPD) for probing nucleic acid dynamics, we examined its intensity and anisotropy decays when intercalated into tRNA^{val} and pBlueScript SK(+) plasmids using frequency-domain fluorometry with a blue light-emitting diode (LED) as the modulated light source. The mean lifetime for the tRNA^{val} was much shorter than that for the pBlueScript SK(+) plasmids, suggesting a more efficient shielding from water by the plasmids. Because of their size difference, the anisotropy decay data also showed a much shorter slow rotational correlation time for the tRNA^{val} than for the pBlueScript SK(+) plasmids. These results indicate that RuPD can be useful for studying nucleic acid dynamics.

[PC1-3] [10/17/2002 (Thr) 13:30 - 16:30 / Hall C]

POTENT INHIBITION OF HUMAN CYTOCHROME P450 1 ENZYMES BY DIMETHOXYPHENYL VINYL THIOPHENE

Lee SangKwang^O, Kim SangHee, Kim MieYoung, Chun YoungJin

College of Pharmacy, Chungang University, Seoul, 156-756, Korea and Natural Products Research Institute, Seoul National University, Seoul, 110-460, Korea

Recently we have reported that various hydroxystilbenes show strong inhibition of human P450 1 activity. A series of synthetic trans-stilbene derivatives were prepared and their inhibitory potentials were evaluated with the bacterial membrane of recombinant human P450 1A1, 1A2 or 1B1 coexpressed with human NADPH-P450 reductase to find new candidates for cancer chemoprevention. Of the compounds tested, SY-021 (3,5-dimethoxyphenyl vinyl thiophene) exhibited a potent inhibition of human P450 1B1 with an IC₅₀ value of 2 nM. SY-021 also showed the inhibition of P450 1A1 with IC₅₀ value of 61 nM and P450 1A2 with IC₅₀ value of 11 nM. SY-021 showed 31-fold selectivity for P450 1B1 over P450 1A1 and 6-fold selectivity for P450 1B1 over 1A2. We have further investigated the inhibition kinetics of P450 1A1, 1A2 and 1B1 by SY-021. The modes of inhibition by SY-021 were non-competitive for all three P450 1 enzymes. Effect of preincubation with NADPH on inhibition of

P450 1B1 by SY-021 was determined. These results suggest that SY-021 is one of the most potent inhibitor of human P450 1 enzymes and may be considered as a good candidate for a cancer chemopreventive agent in human

[PC1-4] [10/17/2002 (Thr) 13:30 - 16:30 / Hall C]

Ceramide analogs inhibit inducible nitric oxide synthase expression and nitric oxide production in interferon-gamma and lipopolysaccharide-stimulated RAW 264.7 macrophages.

Park SungSik^o, Kim Haejong, Yim Chulbu, Kim MieYoung, Chun YoungJin

College of Pharmacy, Chungang University, 156-756, Seoul, Korea

Nitric oxide (NO) production through the inducible nitric-oxide synthase (iNOS) pathway has been implicated in inflammatory diseases and cellular injury. Inhibition of various genes related to inflammation, including iNOS is one of the major roles of well-known anti-inflammatory drugs. In the present study, the effects of ceramide analogs on iNOS expression and NO production were evaluated to investigate how ceramide and its structurally related analogs modulate NO-mediated cellular signals and inflammation. Of ten compounds tested, KY3336 and KY3436 significantly inhibited the NO production in RAW 264.7 murine macrophage cells stimulated with lipopolysaccharides (LPS) and interferon (IFN)-gamma in a dose-dependent manner. Expression of iNOS was also significantly suppressed by these analogs. In conclusion, synthetic ceramide analogs decrease iNOS-dependent NO production in LPS and IFN gamma-stimulated RAW 264.7 macrophages, prompting investigation of its potential use as anti-inflammatory drugs.

[PC1-5] [10/17/2002 (Thr) 13:30 - 16:30 / Hall C]

Environmentally benign, Background-free protein staining in SDS-polyacrylamide gels using an counter ion-dye complex solution.

Jin Li-Tai^o, Hwang Sun-Young, Yoo Gyung-Soo, Choi Jung-Kap

College of Pharmacy, Chonnam National University, Kwangju 500-757, South Korea

Environmentally benign protein staining method in sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) using both an acidic dye, zincon (ZC) and a basic dye, ethyl violet (EV) is described. It is based on a counter ion-dye staining technique that employs oppositely charged two dyes to form an ion-pair complex. The selective binding of the free dye molecules to proteins in an acidic solution produces bluish violet colored bands. It is a rapid procedure, involving only fixing and staining steps that are completed in 45 min. The sensitivity of this method is 5-10 ng of protein which is four-fold better than that of the conventional Coomassie brilliant blue R-250 (CBBR) staining and is comparable to the sensitivities of the colloidal Coomassie brilliant blue G (CBBG) staining, rapid silver staining and SYPRO fluorescence staining procedures. This staining method can be applied to detect for the trace amount of protein in 2D-PAGE. Compatibility of the counter ion-dye stain with MALDI-TOF MS has been demonstrated. Due to the use of nontoxic solvent, ethanol, high sensitivity and rapidity, this stain may be more practical than any other dye-based stains for routine laboratory purposes.

[PC1-6] [10/17/2002 (Thr) 13:30 - 16:30 / Hall C]

ROLES OF PGE₂ AND 15-DEOXY-Δ^{12, 14} PROSTAGLANDIN J₂ IN ET-18-O-CH₃-INDUCED INFLAMMATORY CELL DEATH

Na HyeKyung^o, Surh YoungJoon

College of Pharmacy, Seoul National University, Korea