

them, exhibited strong NF- κ B activity in transfectant human HaCat cells as well as raw 264.7 cells.

[PD3-7] [10/18/2002 (Fri) 13:30 – 16:30 / Hall C]

Triterpene Components from the Leaves of *Acanthopanax sessiliflorus* .

Ryoo Seon-Hyo^{O1}, Chang Seung-Yeup², Park Sang-Young², Oh O-Jin, Yang Ki-Sook, Liu Xiang-Qian, Ro Sook-Hi, Yook Chang-Soo

College of Pharmacy, Kyung Hee University, Seoul 130-704. Korea

Acanthopanax species (Araliaceae) are widely distributed in Asia, which used as tonic and sedative as well as a drug with ginseng-like activities from olden time. There are many reports on the studies of these plants, but there seems no reported about components from the leaves of *Acanthopanax sessiliflorus*, which is indigenous plant to Korea. We have now characterized three triterpenoid compounds from MeOH extract of the leaves of this plant. Based on the physicochemical and spectroscopic data, their structures were identified as chiisanogenin, chiisanoside and 22- α -hydroxychiisanoside.

[PD3-8] [10/18/2002 (Fri) 13:30 – 16:30 / Hall C]

Regulation of the absorption of dietary sugar by α -glucosidase inhibitors from herbal medicines

Yunha Yun^O, ShinSeok Lee, NamYong Chung, Dongsool Yim, Sookyeon Lee, Namjoo Ha, Kyungjae Kim

Department of Pharmacy, Sahmyook University, Seoul 139-742. Korea

The inhibitory activity of several crude drugs on α -glucosidase, which are the key enzyme for carbohydrate digestion and the prevention of diabetic complications, was investigated. This experiment was designed to examine the hypoglycaemic effect of four water extracts crude drugs. We found two drugs, *Mori radidis Cortex* and *Cudrania radidis Cortex* in several crude drugs remarkably inhibited α -glucosidase. Two crude drugs were examined in streptozotocin induced high blood glucose mice. Oral administration of *Mori radidis Cortex* and *Cudrania radidis Cortex* lowered the blood glucose level in the high blood glucose mice. High blood glucose was induced in mice by *Cudrania radidis Cortex* intraperitoneal injections of streptozotocin (STZ, 150 mg/kg). *Mori radidis Cortex* and *Cudrania radidis Cortex* strongly showed inhibitory activity by 36.4 and 21.9% in mice loaded with starch. In the case of the maltose load test, *Mori radidis Cortex* and *Cudrania radidis Cortex* showed inhibitory activity by 19.5 and 6.1%. We used acarbose for positive standard. We compared with acarbose and starch groups but also compared with acarbose and maltose groups. When compared with acarbose and starch groups, *Mori radidis Cortex* was 1.2 times higher than acarbose but *Cudrania radidis Cortex* was lower than acarbose. When compared with acarbose and maltose groups, *Mori radidis Cortex* was about 7 times higher than acarbose and *Cudrania radidis Cortex* was about 2 times higher than acarbose.

[PD3-9] [10/18/2002 (Fri) 13:30 – 16:30 / Hall C]

Effect of Ethyl Acetate Extract from *Caesalpinia sappan* L. on Melanogenesis in Melan-a cells

Chun HyunJa^O, Choi WonHyung, Choi EunYoung*, Yang HyunOk**, Baek SeungHwa

Dept. of Herbal Resources, Professional Graduate School of Oriental Medicine, Wonkwang University, *Dept. of Beauty and Skin Care, Kimcheon Science College, **Dept. of Cosmetics, Wonkwang Health Science College

Melanin is a main pigment found in skin, hair and eyes, and tyrosinase plays an important role in the process of melanin polymer biosynthesis. *Caesalpinia sappan* L. (*C. sappan*) has been commonly used in Oriental folk medicines to promote blood circulation and as analgesic as well as remedy for thrombosis. This present study was designed to investigate the effect of ethyl acetate extract from *C. sappan* on melanogenesis in Melan-a cells. The cells showed a dose-dependent inhibition in their proliferation without apoptosis after treatment with ethyl acetate extracts. Therefore, the growth retardation by the extract may be due to the cell arrest or cell differentiation. The melanin content and