

## Effects of Epidermal Growth Factor on the Apoptosis and Implantation Related Genes in Bovine Embryos Developing *in vitro*

Sae Young Park, Jin Cheol Tae, Eun Young Kim, SePill Park, Jin Ho Lim<sup>1</sup>, Nam Hyung Kim<sup>2</sup>

*Maria Infertility Hospital/ Medical Institute/ Maria biotechnology, <sup>1</sup>Maria Infertility Hospital, <sup>2</sup>Chungbuk National University*

Epidermal growth factor (EGF) induces well-documented mitogenic and differentiating effects on murine and bovine preimplantation embryos. However, the effects of EGF on apoptosis and implantation-related gene expression in bovine embryos developing *in vitro* have not been evaluated. The objective of this study was to determine the effects of exogenous EGF in the presence and absence of BSA on the preimplantation development of bovine embryos. In addition, we measured cell number, apoptosis, and expression of apoptosis and implantation-related genes of the blastocysts that developed in these culture conditions. *In vitro* produced bovine embryos were randomly cultured in the same medium containing 0 or 10 ng/ml EGF in the presence and absence of 0.8% BSA. More 2-cell embryos developed into blastocysts at day 7 when BSA was present than when BSA was absent. The addition of 10 ng/ml EGF into the medium did not significantly increase the developmental rate and the cell numbers per blastocyst. However, addition of EGF in the presence of 0.8% BSA significantly reduced the degree of apoptosis in the blastocysts ( $P < 0.01$ ). To investigate whether EGF modulates mRNA expression of apoptosis-related genes, mRNA was prepared from single blastocysts and each preparation was subjected to RT-PCR for Bcl-2 and Bax transcripts. EGF did not alter the relative abundance of Bax gene expression in the presence of BSA, but increase Bcl-2 ( $P < 0.01$ ). The relative abundance of Interferon tau expression was increased by EGF treatment in the presence of BSA. These results suggest that EGF and BSA synergistically enhance Bcl-2 and interferone tau gene expression, which may result in a net increase in viability in bovine embryos.

Key words) *Bovine embryo, Epidermal growth factor, BSA, Preimplantation development, Apoptosis*