

A case study of red tide detection around Korean waters using satellite remote sensing

Y. S. Suh · N. K. Lee · L. H. Jang · H. G. Kim · J. D. Hwang

Development of Oceanography and Marine Environment, National Fisheries Research & Development
408-1 Sirang-ri, Gijang-eup, Busan 619-902, Korea
yssuh@nfrdi.re.kr

Abstract: Korea has experienced 10 a *Cochlodinium polykrikoides* red tide outbreaks during the last 10 years (1993-2002). The monitoring activities at National Fisheries Research and Development Institute (NFRDI) in Korea have been extended to all the coastal waters after the worst of fish killing by *C. polykrikoides* blooms in 1995. NFRDI is looking forward to finding out the feasibility of red tide detection around Korean waters using satellite remote sensing of NOAA/AVHRR, Orbview-2/SeaWiFS, IRS-P4/OCM and Terra/MODIS on real time base. In this study, we used several alternative methods including climatological analysis, spectral and optical methods which may offer a potential detection of the major species of red tide in Korean waters. The relationship between the distribution of SST and *C. polykrikoides* bloom areas was studied. In climatological analysis, NOAA, SeaWiFS, OCM satellite data in 20th and 26th August 2001 were chosen using the known *C. polykrikoides* red tide bloom area mapped by helicopter reconnaissance and ground observation. The 26th August, 2001 SeaWiFS chlorophyll *a* anomaly image against the image of non-occurring red tide for August 20, 2001 showed the areas *C. polykrikoides* occurred. The anomalies of chlorophyll *a* concentration from satellite data between before and after red tide outbreaks showed the similar distribution of *C. polykrikoides* red tide in 26th August, 2001. The distribution of the difference in SST between daytime and nighttime also showed the possibility of red tide detection. We used corrected vegetation index (CVI) to detect floating vegetation and submerged vegetation containing algal blooms. The simple result of optical absorption from *C. polykrikoides* showed that if we use the optical characteristics of each red tide we will be able to get the feasibility of the red tide detection.

Keywords: *Cochlodinium polykrikoides*, Red tide, Satellite remote sensing.

1. Introduction

Korea has experienced a *Cochlodinium polykrikoides* red tide 10 out of the last 10 years (1993-2002). This species of red tide has been experienced at least once by all the southern coast of the Korean peninsular and has been transported up the southeastern coast all the way to the northeastern coast since 1995. The impression is that red tide is spreading and becoming more common, not only in the nearest coastal water, but also in the offshore.

The monitoring activities at National Fisheries Research and Development Institute (NFRDI) in Korea have been extended to all the coastal water of Korean waters after the worst outbreaks of fish killing *C. polykrikoides* blooms in 1995. Mitigating red tide impacts

requires an ability to monitor for blooms and to forecast occurring and spreading of red tide. Nowadays NFRDI is looking forward to finding out the feasibility of red tide detection around Korean waters using satellite remote sensing of NOAA/AVHRR (Advanced Very High Resolution Radiometer), Orbview-2/SeaWiFS (Sea Viewing Wide Field-of-View Sensor), IRS-P4/OCM (Ocean Colour Monitor) and Terra/MODIS (Moderate Resolution Imaging Spectroradiometer) on real time base.

For remote sensing to be effective, red tide must be detectable, either directly through its effect on the water color, or indirectly by correlation with algal blooms or association with a water mass that can be monitored by other remotely sensed characteristic such as sea surface temperature (Tester *et al.*, 1991; Keafer and Anderson 1993).

In this study, I used several alternative methods including a climatological analysis, spectral and optical methods which may offer a potential detection of the major species of red tide in Korean waters.

2. Material and Method

The sea surface temperature (SST) data were derived from the infrared channels of AVHRR sensor on NOAA satellites from 1995 to 2002 (Suh *et al.*, 2000; McClain *et al.*, 1985). There were two channels (channel 1 (ch1): 571-686 nm, channel 2 (ch 2): 713-989 nm) on AVHRR to apply ocean color remote sensing in 1995. At that time, I just used the band ratio (ch1 of AVHRR/ ch2 of AVHRR) to detect the major species of algal bloom, *C. polykrikoides*, *Ceratium furca* and *Notiluca* in 1995 as Gallegos (1990) did it.

To estimate chlorophyll *a* derived from SeaWiFS and OCM were analyzed by using ocean color chlorophyll 2 algorithm, OC2 (O'Reilly *et al.*, 1998). In case of Terra/MODIS data, we used visible channels and near-infrared channel to get the spectral band ratios to estimate distribution of red tide.

In climatological analysis, the estimated chlorophyll *a* concentration derived from the ocean color satellites for red tide occurring day is compared to that of before red tide occurring day. Distribution of the SST differences derived from NOAA infrared data between daytime and nighttime in the red tide occurring coastal water was made to detect indirectly the distribution of algal bloom

using the biological characteristics of daily vertical migration of phytoplankton.

Absorption spectra between 250 nm and 800 nm of *C. polykrikoides*, *H. akashiwo* and *P. minimum* were measured by spectrophotometer (Varian Carry 100).

3. Conclusions

The relationship between the distribution of SST and *C. polykrikoides* bloom areas was studied. The cold water was formed by upwelling water from deep layer in the southeastern coast of the Korean peninsula played a role in blocking the spreading of red tides during summer season. The red tides that initially bloomed at the coast of Pohang on September 21, 1995 moved to the coast of Uljin on September 26, 1995. The skipped appearance of the red tides in the areas between Pohang and Uljin was due to the East Korean Warm Current, which was moving offshore from Pohang to approach to Uljin.

We used corrected vegetation index (CVI) to detect floating vegetation and submerged vegetation containing algal blooms as Sonia Gallegos did it in 1990. However, we have got different result because of different species of algal bloom. Algal bloom areas were estimated by ratios of ch1 digital values (60-80) and ch2 digital values (90-120). The major species of algal bloom were *C. polykrikoides*, *Ceratium furca* and *Noctiluca*. Among the major species, *Noctiluca* was detected by CVI in 1995.

In climatological analysis, NOAA, SeaWiFS, OCM satellite data in 20th and 26th August 2001 were chosen using the known *C. polykrikoides* red tide bloom area mapped by helicopter reconnaissance and ground observation. The August 26, 2001 SeaWiFS chlorophyll *a* anomaly imageries against the imageries of non-occurring red tide for August 20, 2001 showed the areas *C. polykrikoides* occurred.

The standard global chlorophyll algorithm (Ocean Color 2) tends to overestimate chlorophyll in the southern coast of Korea like case 2 waters (Suh *et al.*, 2002). When a coastal correction is not applied, may increase this error another two fold. However, the anomalies of chlorophyll *a* concentration from satellite data between after occurring red tide and before occurring red tide showed the similar distribution of *C. polykrikoides* red tide in August 26, 2001.

The distribution of the difference in SST between daytime and nighttime also showed the possibility of red tide detection. The simple result of optical absorption from *C. polykrikoides* showed that if we use the characteristics of response for the red tide we will be able to get the feasibility of the red tide detection.

Acknowledgement

This study was funded by the research project "The application of satellite data for fisheries oceanography" of the National Fisheries Research and Development

Institute and the Korea Aerospace Research Institute.

References

- [1] Gallegos S. C., 1990. Evaluation of the potential of the NOAA-n AVHRR reflective data in oceanography. Ph. D. paper, University Texas A. & M., pp. 1-88.
- [2] Keafer B. A. and Anderson D. M., 1993. The use of remotely-sensed sea surface temperatures for studies of *Alexandrium* bloom dynamics in the southwestern Gulf of Maine. In: Smayda T. and Shimizu Y. (eds), Toxic Phytoplankton Blooms in Sea, pp 763-8. Elsevier, Amsterdam, The Netherlands
- [3] McClain E. P., W. G. Pichel and C. C. Walton, 1985. Comparative performance AVHRR based multi-channel sea surface temperatures, *J. Geophys. Res.*, 90: 11,587-11,601.
- [4] O'Reilly J. E., S. Maritorena, B. G. Mitchell, D. A. Siegel, K. L. Carder, S. A. Garver, M. Kahru and C. R. McClain, 1998. Ocean color chlorophyll algorithms for SeaWiFS. *J. Geophys. Res.*, 103: 24,937-24,953.
- [5] Suh Y. S., J. H. Kim and H. G. Kim, 2000. Relationship between sea surface temperature derived from NOAA satellites and *Cochlodinium polykrikoides* red tide occurrence in Korean coastal waters. *J. Korean Environ. Sci. Soc.*, 9(3): 215-221.
- [6] Suh Y. S., L. H. Jang, N. K. Lee and B. K. Kim, 2002. Development of the regional algorithms to quantify chlorophyll *a* and suspended solid in the Korean waters using ocean color. *J. Korean Fish. Soc.* 35(3): 207-215.
- [7] Tester P. A. and Stumpf R. P., Vukovich F. M. *et al.*, 1991. An expatriate red tide bloom: transport, distribution, and persistence. *Limnol. Oceanogr.* 36(5): 1,053-61.