

Effect of IVM Medium and Protein Source on *In Vitro* Maturation of Canine Oocytes

Hyo-Sang Lee, Xi-Jun Yin, Young-Ho Lee, Se-Jin Chun, Young-Il Suh, Keum-Ju Park, Jin-Sung Seo, Su-Jin Jo and Il-Keun Kong
Department of Animal Science and Technology, College of Agriculture and Life Science, Sunchon National University

This study was conducted to determine the ability of nuclear development of canine oocytes depend on the kind of maturation media and addition of serum sources. Ovaries were collected from a bitches at various stages of estrus cycle by an ovariohysterectomy. Oocytes were collected of cumulus oocytes complexes after slicing of ovaries with blade. The maturation medium was containing 0.6 mM/ml cysteine, 0.2 mM pyruvic acid, 20 ng/ml E₂ and 1 µg/ml rST. Exp. 1, the oocytes were matured in four different maturation medium as follows: 1) TCM-199, 2) DMEM, 3) NCSU37 and 4) modified-NCSU37 with 10% FBS. Exp. 2: the oocytes were matured in mNCSU37 supplemented with different protein sources (10% FBS, 10% EDS, 0.3% BSA and 0.1% PVA) to select the optimal one. Oocytes were matured in a humidified atmosphere containing 5% CO₂ at 39°C for 72 hrs. The maturation rate were analyzed by Duncan's multiple range test using General Linear Models procedure in SAS.

The rates of meiotic resumption to MI-MII depend on different culture media were achieved with TCM-199 (5.2%), DMEM (5.0%), NCSU37 (7.2%) and m-NCSU37 (5.9%), respectively. The rates of meiotic resumption to MI-MII according to addition of protein source were 10% FBS (13.3%), 10% EDS (25.0%), 0.3% BSA (25.0%) and 0.1% PVA (15.4%), respectively. In conclusion, the results obtained showed that *in vitro* maturation media and protein supplement to m-NCSU37 culture medium tested did not promote the final steps of IVM in canine oocytes.

Key words) *Canine, Oocyte, In vitro maturation Medium*