

**[P-21]****Estrogenic activity of pomegranate extract  
in MCF7-ERE cells**

Cho minjung, An Jinyoung, Yang soyun, Sheen Yhunyhong, Lansky Ephraim<sup>1</sup>  
*College of Pharmacy, Ewha womans University, Seoul, Korea*  
*Rimonest Ltd., Science Park Technion<sup>1</sup>, Nesher, Israel*

Pomegranate, a small tree originating in Orient, belongs to Punicaceae family. The seeds contain an oil of which about 80% is rare trans 18 carbon fatty acid (punicic acid), and have highest botanical concentration of a sex steroid, estrone. Pharmacological properties of pomegranate extract have been studied, with anti-microbial, anti-parasitic, anti-viral, and anti-cancer effects.

We have examined the estrogenic activity of the pomegranate extracts using MCF-7-ERE cells. MCF-7-ERE cells, stably transfected with pERE-Luc were treated various amount of pomegranate extract and after overnight treatments, luciferase activity were measured. Estradiol(E2) dose dependently induced luciferase activity in this cell and maximal response is obtained at 1 nM E2.

82, 80 extract of pomegranate showed stronger estrogenic activity than that of 1 nM E2. And 40, 41, 44 (EtOH bot seed, fer jui polyphenols, scf peric polyphenol) showed about half inductive effect as compared to 1 nM E2. These increase in luciferase activity by pomegranate was inhibited by Tamoxifene concomitant treatment. Therefore is reasonable to interpret that pomegranate extract increased luciferase activity via ER.