

Molecular Cytogenetics of rDNA Genes in the Youngia sonchifolia

Soo Young Kim¹, Sung Ran Min², Dong Su In³, Jang Ryol Liu^{2,3}, Jae Wook Bang^{1*}

Department of Biology, Chungnam National University, Daejeon 305-764, Korea

²Korea Research Institute of Bioscience and Biotechnology (KRIBB), 52-Oun Dong, Yusong, Daejeon 305-333, Korea

³Eugentech Inc. 52-Oun Dong, Yusong, Daejeon 305-333, Korea

Objectives

In this study, we observed the chromosome numbers and determind the 5s and 45s rRNA gene loci of *Youngia sonchifolia* and it's regenerants.

Materials and Methods

- 1. Materials: Y. sonchifolia and it's regenerants
- Methods: Chromosome preparation and Feulgen staining were applied for karyotype analysis. The 45s rDNA and 5s rDNA were labelled with digoxigenin-11-dUTP and biotin-16-dUTP by PCR method for fluorescent in situ hybridization (FISH).

ere

Acknowledgement

This research was supported by a grant (Code #: PF001201-03: Pl J. W. Bang) from Plant Diversity Research Center of 21st Century Frontier Research Program funded by Ministry of Science and Technology of Korean government.

and its regenerants were 2n=10 and all the chromosomes pairs

were submetacentrics. NORs were observed in chromosome 3. Physical mapping of the two multigene families coding for 5s

rRNA and 45s rRNA in Y. sonchifolia and regenerants was carried


out using fluorescence in situ hybridization (FISH). The 5s rDNA

signals were located only on long arm of chromosome 3 and 45s rDNA loci were found at NORs region so the two ribosomal gene

families were generally located on same chromosome.

Results and Discussion

In karyotype analysis, chromosome numbers of Y. sonchifolia






Figure 1. Somatic metaphase chromosome complement (2n=10) and karyotype of *Y. sonchifolia* and two-color FISH pattern. Bar, 5 μ m.

^{*}Corresponding author. Tel 042-821-5497 E-mail bangjw@cnu.ac.kr