

# $^{11}\text{B}$ NMR Study of Vortex Dynamics in $\text{MgB}_2$ Superconductor

Kyuhong Lee<sup>a</sup>, B. J. Mean<sup>a</sup>, K. H. Kang<sup>a</sup>, Moohee Lee\*<sup>a</sup>, B. K. Cho<sup>b</sup>

<sup>a</sup> *Department of Physics, Konkuk University, Seoul 143-701, KOREA*

<sup>b</sup> *Department of Materials Science and Technology, GJ-IST, Kwangju, 500-712, KOREA*

Vortex structure and dynamics for Magnesium-diboride have been studied using pulsed NMR techniques. We have measured spectrum, shift, detuning frequency, and transverse relaxation rate( $1/T_2$ ) of  $^{11}\text{B}$  NMR for  $\text{MgB}_2$  powder from 4 K to the normal state under 1.8 T of external magnetic field. In the superconducting state, the spectrum shows the characteristic field distribution with diamagnetic shift and broad linewidth due to the imperfect penetration of magnetic field.  $1/T_2$  results have a single peak with small change of the rate contrary to the results of nickel borocarbides. Below 16 K, the shape of  $T_2$  decay changes from Lorentzian to Gaussian. It means that the thermal fluctuation is reduced as temperature decreases and the vortex motion is smaller than nickel borocarbides.