

E001

Characterization of Calcium-Activated Bifunctional Peptidase of the Psychrotrophic *Bacillus cereus*Jong-Il Kim^{*}, Hyo-Jin Lee, and Sun-Min Lee

Department of Food and Microbial Technology, Seoul Women's University

The protease purified from *Bacillus cereus* JH108 has the function of leucine specific endopeptidase. When the enzyme activity was measured by hydrolysis of synthetic substrate (*N*-succinyl-Ala-Ala-Pro-Leu-*p*-nitroanilide) it exhibited optimal activity at pH 9.0, 60°C. The endopeptidase activity was stimulated by Ca⁺⁺, Co⁺⁺, Mn⁺⁺, Mg⁺⁺ and Ni⁺⁺ and inhibited by metal chelating agents such as EDTA, 1,10-phenanthroline, and EGTA. Addition of serine protease inhibitor, PMSF, resulted in the elimination of the activity. The endopeptidase activity was fully recovered from the inhibition of EDTA by adding 1 mM Ca⁺⁺, and partially restored by Co⁺⁺ and Mn⁺⁺, indicating that the enzyme was stabilized and activated by divalent cations and has a serine residue at the active site. Addition of Ca⁺⁺ increased the pH and heat stability of endopeptidase activity. These results show that endopeptidase requires calcium ions for activity and/or stability. A Lineweaver-Burk plot analysis indicated that K_m value of endopeptidase is 0.315 mM and V_{max} is 0.222 μmol of *N*-succinyl-Ala-Ala-Pro-Leu-*p*-nitroanilide per min. Bestatin was shown to act as a competitive inhibitor to the endopeptidase.

E002

Conversion of Methylglyoxal to Acetol by *Escherichia coli* Aldo-keto ReductasesDongkyu Park^{*}, Jeeyeon Song, Minsuk Kwon, Jinyoung Yoon, Dongwook Choi, and Chankyu Park

Korea Advanced Institute of Science and Technology

Methylglyoxal(MG) is a toxic metabolite known to accumulate in various cell types. We detected *in vivo* conversion of MG to acetol in MG-accumulating *E. coli* cells by ¹H-NMR spectroscopy. Search for homologs of the mammalian aldo-keto reductases, reported to exhibit activity to MG, revealed nine ORFs from the *E. coli* genome. Based on both sequence similarities and preliminary screening with ¹H-NMR for crude proteins of the corresponding mutant strains, we selected five genes, *yafB*, *yqhE*, *yeaE*, *yghZ*, and *yajO*. Quantitative assessment of acetol production including its metabolite 1,2-propanediol from the crude extracts of these mutants indicated that the *yafB*, *yqhE*, *yeaE* and *yghZ* genes are involved in the conversion of MG in the presence of NADPH. Although enzymatic properties of these proteins are generally consistent with their activities on MG conversion, the *yqhE* mutation did not exhibit further sensitivity to MG in glyoxalase-deficient strain. The results imply that the glutathione-independent detoxification of MG *in vivo* can occur through multiple pathways, consisting of *yafB*, *yeaE*, and *yghZ* genes, leading to a generation of acetol.

[Supported by the BK21 and glycomics program]

E003

***Csp* Genes of *Corynebacterium glutamicum* are Involved in Cold-shock Response**Wan-Soo Kim^{1*}, Younhee Kim², and Heung-Shick Lee¹¹Department of Biotechnology and Bioinformatics, Korea University, ²Department of Oriental Medicine, Semyung University

Corynebacterium glutamicum genes which appear to play a role in cold shock response were isolated using the *aceB* promoter of *C. glutamicum*. Expression of the gene in *E. coli* resulted in 70% reduction in the expression of the *aceB* gene as monitored by the β-galactosidase activity expressed from the P_{aceB}-*lacZ* reporter plasmid. The isolated gene encoded a putative protein composed of 63 amino acids which showed high similarity with cold shock proteins of other bacteria and designated *cspA*. Scrutiny of the *C. glutamicum* ATCC13032 genome sequence identified 2 additional *csp* genes and the expression of the genes were monitored by fusing chromophenicol acetyltransferase (CAT) genes to the promoter region of the genes. For all 3 genes, the highest activity was observed in the late log phase. In addition, expression of the genes was stimulated when the cells carrying the clones were given a cold shock. These data indicate that the *cspA* gene may encode a transcriptional regulator and the *csp* genes are probably involved in the cold shock response in *C. glutamicum*.

[Supported by a grant from KOSEF.]

E004

Antibacterial and Antiviral Activity of the Culture Soup Produced from *Leuconostoc kunchii*Sung-Hyun Song^{*}, Tae-Hwan Kim, Jin-Hee Park, Dong-Il Jang, Chang-Jin Lee, and Sa-Ouk Kang

Laboratory of Biophysics, School of Biological Sciences, and Institute of Microbiology, Seoul National University

The culture soup produced from *Leuconostoc kunchii* has antibacterial activity against gram-positive bacteria *Bacillus subtilis* and gram-negative bacteria *Escherichia coli*, *Salmonella typhimurium*, *Shigella sonnei*, *Vibrio vulnificus*. The antibacterial activity of the culture soup was stable over 100°C but not showed over pH 5. Antibacterial activity of the culture soup was better at 3 days after inoculation. When treatment of Proteinase-K to the culture soup, antibacterial activity remained. Partially purified substance(s) showed antibacterial activity against *B. subtilis* and other strains. The culture soup reveals antiviral activity against influenza virus(H3N2).