

F037

Partial Reconstitution of Hepatitis C virus RNA Polymerization by Heterologous Expression of NS5B Polymerase and Template RNA in Bacterial Cell

Jong-Ho Lee*, Sangyoon Lee, Minsoo Kim, Yu Jeong Byun, and Heejoon Myung

Department of Bioscience and Biotechnology, Hankuk University of Foreign Studies

The hepatitis C virus is a major etiological agent causing chronic hepatitis in humans. The current study presents a surrogate model system using a viral polymerase and RNA template. A plasmid expressing the HCV NS5B polymerase was maintained with a plasmid containing a reporter gene in an *E. coli* cell. The reporter construct contained the HCV 5' UTR followed by a luciferase gene with a specific orientation so that a minus-sense transcript containing the luciferase fused to the 5' UTR was produced after the initial transcription. When the HCV NS5B polymerase was expressed in the same cell, the primary transcript was recognized by the polymerase due to the presence of the minus-sense 5' UTR, and a secondary transcript containing a plus-sense luciferase gene produced. Thus, a simple luciferase assay was able to measure the HCV NS5B polymerase activity. The production of minus and plus-sense transcripts was confirmed by an RT-PCR, while the production of HCV NS5B and expression of the reporter luciferase in the bacterial cell were confirmed by immunofluorescence microscopy. The polymerization occurred in the absence of any other viral/host factors.

F038

The Lethal Phenotypes of Dna2 mutations Are Rescued by Overexpression of the DNA Repair Proteins in *Saccharomyces cerevisiae*

Hyun-Sun Lee*, Sung-Hoon Kwon, In-Hwan Lee, Hyun-Jung Kim, and Sung-Ho Bae

Department of Biological Sciences, Inha University

Dna2 protein involved in Okazaki fragment processing is a multifunctional enzyme with two essential enzyme activities, endonuclease and helicase. The endonuclease activity is essential to remove RNA-DNA primers of Okazaki fragments, while the helicase facilitates removal of secondary structures formed in the primers. In an effort to identify genes interacting genetically with Dna2, multi-copy suppressors were screened in yeast that rescue the lethality of Dna2 mutants devoid of helicase (*dna2KE*) or endonuclease (*dna2DA*) activities. A putative helicase, MPH1, was identified as a suppressor for *dna2KE*. Purified Mph1 protein possessed both ATPase and helicase activities, suggesting that it might replace the function of Dna2 helicase. The suppressors for *dna2DA* mutant encode the nucleases involved in the nucleotide excision or double-strand break repair pathways. They also contain several novel proteins that associate with these enzymes. These results suggest that some DNA repair proteins may participate in removing the unprocessed RNA-DNA primers which are not cleaved by Dna2 and/or other Okazaki fragment processing enzymes.

[Supported by grant from KRF, KRF-2004-041-C00240]

F039

A Novel Interferon Resistance Mechanism of Hepatitis C Virus (HCV) Mediated by NS5B Protein

Jay Seok Park*, Minhee Seo, Jie Hye Lim, and Heejoon Myung
Department of Bioscience and Biotechnology, Hankuk University of Foreign Studies

Two mechanisms of interferon resistance for hepatitis C virus are known. One is mediated by E2 protein and the other is mediated by NS5A protein. Both target host PKR for their action. We report here a third mechanism mediated by NS5B protein. An interferon effector molecule called p56 was recently identified. It binds to p48 subunit of eIF3 complex to inhibit translation. We showed that NS5B also bound to p48 of eIF3. It was demonstrated by yeast two hybrid analysis, GST-pulldown assay. The subcellular colocalization was confirmed by immunofluorescence confocal microscopy. The inhibitory effect of interferon alpha was diminished by the presence of NS5B in a reporter assay from cultured human hepatocyte. When NS5B was present, the inhibitory effect of p56 on translation was diminished in an *in vitro* lysate assay. The inhibition was in a dose-dependent manner.

F040

Identification of Genes Regulated by *luxO* Homologue in *Vibrio vulnificus* MO6 24/O Using a Small-scale DNA Microarray

Kyung-Shin Lee¹*, Saet-Byoung Lee², Kyu-Ho Lee³, Byung-Soo Kim⁴, and Kun-Soo Kim^{1,2}

¹Program of Integrated Biotechnology and ²Departments of Life Science, Sogang University, ³Department of Environmental Science, Hankuk University of Foreign Studies, ⁴Department of Applied Statistics, Yonsei University

Vibrio vulnificus is an opportunistic pathogen, which causes a primary septicemia on human. As in other *Vibrio* species, *V. vulnificus* contains *luxO* homologue, which is known to regulate the expression of genes related with the luminescence and the virulence in *V. harveyi* and *V. cholerae*, respectively. To identify the genes regulated by LuxO in *V. vulnificus*, a small-scale DNA microarray was constructed, which comprises 160 PCR products, putatively encode functions associated with pathogenicity and gene regulations, derived from the genome of *V. vulnificus*. The cDNAs of wild-type and *luxO* null-mutant were hybridized on SGVV v1.0.1 chip which was printed with 640 spots (quadruplicates of 160 genes). We carried out two sets of hybridizations and another dye-swap of one set of hybridizations. To analyze the array profile, we employed two statistical procedures, the SAM procedure and the Lonstedt and Speed's empirical Bayes procedure using B statistic. As a result, 4 differentially expressed genes of statistical significance, including DNA helicase and DNA binding protein H-NS were identified. [Supported by the 21C Frontier Microbial Genomics and Applications Center Program.]