

## P-32

**Ca<sup>2+</sup> and K<sup>+</sup> concentrations change during early embryonic development in mouse**

Dawon Kang, Chang-Gi Hur, Chang-Rok Choi  
Jae-Yong Park, Seong-Geun Hong and Jaehee Han

*Department of physiology, College of Medicine and Institute of Health Science,  
Gyeongsang National University, Jinju, 660-751*

Ions play a fundamental role in nearly all cellular processes. However, it is little known whether how ions regulate early embryonic development. In this study, we examined changes in Ca<sup>2+</sup> and K<sup>+</sup> currents and concentrations in embryos and oviduct during mouse early embryonic development using patch clamp technique and confocal microscopy. During early embryonic development, K<sup>+</sup> concentration in oviduct increased markedly, but intracellular Ca<sup>2+</sup> concentration did not change. The membrane potential was depolarized (from -38 mV to -16 mV), and Ca<sup>2+</sup> current decreased suggesting that these events may coincide. The membrane potential induced by influx of Ca<sup>2+</sup> first hyperpolarized and then depolarized in mouse oocytes. The hyperpolarization (HP) elicited by Ca<sup>2+</sup> in membrane potential resulted from activation of K<sup>+</sup> channels. Our results showed that the K<sup>+</sup> channel was not blocked by TEA and apamin, well-known K<sup>+</sup> channel blockers. These results suggest that the K<sup>+</sup> channel that induces HP in mouse oocytes could belong to another K<sup>+</sup> channels including two-pore domain K<sup>+</sup> channel that are insensitive to TEA and apamin, and the changes in ions following embryonic development mainly play a critical role in regulation of membrane potential.

**Keywords:** *Calcium, Potassium, Mouse, Oocytes, Embryo, K<sup>+</sup> channel.*  
*Support contributed by BK21*