

S-4 [11:50-12:20]

The anti-dementia effects of herbal
medicine formula YMJ -from lab to
clinic

Prof. Hyunsu Bae
(Kyung Hee University)

[11:50 – 12:20]

The anti-dementia effects of herbal medicine fomula YMJ From lab to clinic

Hyunsu Bae M.D., Ph.D.
College of Oriental Medicine, KyungHee University

The herbal extract YMJ has been widely used as an anti-aging herbal medicine for centuries in Asian countries. Among the various modified subscriptions of YMJ, YMJd was formulated to enhance memory retention. This study has three goals: 1) to quantitatively evaluate the memory-enhancing effect of YMJd using behavior tasks; 2) use cDNA micro-array tools to identify candidate genes responsible for enhancing memory; and 3) to evaluate enhancing cognitive ability in normal human subjects and discusses its possibility to be used to treat dementia patients with deficient cognitive ability. Memory retention abilities are addressed by the passive avoidance task with SD male rat. The retention time of the YMJd group was significantly delayed (~100%), whereas *Ginkgo biloba* and Soya lecithin treatment were only delayed 20% and 10%, respectively. In order to confirm the anti-dementia effects of YMJd, ibotenic lesioned alzheimer's disease model was utilized. Ibotenic lesion of the medial septum (MS) of rat showed the impaired performance in the Morris water maze test and severe cell losses in the MS, as indicated by decreased choline acetyltransferase-immunoreactivity in the medial septum. Daily administrations of YMJd (100 mg/kg, i.p.) for 21 consecutive days produced significant reversals of ibotenic acid-induced deficit in learning and memory. The cDNAs from the hippocampi of YMJd and rat control groups were applied to an Incyte rat GEM2 cDNA microarray. The microarray results showed that transthyretin and PEP-19 were abundantly expressed in the YMJd treated group. Importantly, PEP-19 is a neuron-specific protein that inhibits apoptotic processes. On



the other hand, neuronal genes involved in neuronal death or neurodegeneration such as pentraxin and spectrin were abundantly expressed in the control group. Finally, K-WAIS tests, a Korean version of an individual intelligence quotient (IQ) test, and a P300 latency assessment of Event-Related Potential (ERP) were conducted in order to measure changes in cognitive ability before and after 6 weeks of YMJd treatment in normal young adults. The K-WAIS mean scores of the group treated with YMJd were significantly higher than those of the placebo group ($p < 0.05$), and their mean P300 latency was substantially shorter ($p < 0.005$). These results suggest that YMJd treatment accelerates the speed of information processing and enhances cognitive ability. YMJd treatment may help dementia patients or the elderly recover from cognition deficiencies or degeneration in clinic

The anti-dementia effects of herbal medicine fomula YMJ From lab to clinic

Hyunsu Bae M.D., Ph.D.
KyungHee Univeristy,
College of Oriental Medicine


1. Background

- 미국에서는 400만명 이상이 알츠하이머형 치매로 고생하고 있으며 우리나라에서는 노령화 사회로 진입함에 따라 빠른속도로 치매환자들이 증가하고 있음
- 알츠하이머병과 관련된 세계 치매치료제 시장 규모는 2002년 기준 약 4조 6천억원
- 국내치매치료시장 규모는 2000년 약 24억원, 2001년 약 48억원, 2002년도에는 약 125억원의 시장규모로 2년사이 무려 5배나 시장규모가 확대됨
- 그러나, 국내 치매치료제는 외국에서 수입하거나 유통비를 지불하는 것이 대부분임
- 이뿐만 아니라 대부분의 치매치료제의 부작용은 심각한 간독성, 소화 장애 및 운동장애임
- 구거지황탕은 전통한방 유래의 새로운 한치매제로 의학 질감 뿐 만 아니라 수천년간 임상 및 예비 임상실험에서 별다른 부작용이 없음이 밝혀져서 안전하면서도 효과 있는 새로운 치매치료제 기대




Behavior tests


Morris water maze



Passive avoidance



Radial arm maze



Morris water maze




Fig. 6. Target entry number of each group in Morris water maze

Normal: group of rats without any treatment

Control: group of rats treated with L-NAME

Sample: group of rats administrated to YMJ in the presence of L-NAME pretreatment

☆: $p < 0.05$ by student t-test

Morris water maze




Fig. 7. First latency of each group in Morris water maze

Normal: group of rats without any treatment

Control: group of rats treated with L-NAME

Sample: group of rats administrated to YMJ in the presence of L-NAME pretreatment

☆: $p < 0.05$ by student t-test

Morris water maze




Fig. 8. Memory score of each group in Morris water maze

Normal: group of rats without any treatment

Control: group of rats treated with L-NAME

Sample: group of rats administrated to YMJ in the presence of L-NAME pretreatment



☆: $p < 0.05$ by student t-test



Radial arm maze test

Trial	1	2	3	4	5	6
Normal	7.83±0.11 ^{a)}	7.83±0.17	7.67±0.22	7.08±0.65	6.33±0.89	7.67±0.33
Control	0.93±0.59	3.57±0.90	4.07±1.03	4.29±3.85	1.14±1.07	4.43±1.03
Sample	3.14±1.13	5.78±1.01	5.56±1.02	7.11±0.89	7.41±0.44	7.78±0.22
p value	0.07	0.13	0.32	0.06	0.01*	0.01*






a): mean ± SE
 Normal: Group of normal rats without any treatment
 Control: Group of rats treated with L-NAME
 Sample: Group of rats administered *Yukimjhwangtang* in the presence of L-NAME pretreatment
 *: p<0.05 by student t-test

Memory decay kinetics







cholinergic model




cholinergic model






Microarray analysis









Gene Name	BDE	Genbank*
Rat prealbumin (transthyretin)	-3.2	K03252
Mouse mRNA for type II 57 kd keratin.	-2.2	X03491
Rat phospholipidylethanolamine N-methyltransferase L14441	-2.2	
Rat neuron-specific protein PEP-19	-2.1	M24852
Mouse Purkinje cell protein-4 (Pcp-4)	-2.1	M96359
Rat Axi receptor tyrosine kinase	-2.0	AF046886
Mouse mRNA for type I 47 kd keratin.	-2.0	X03492
Sprague-Dawley (clone LRB9) RAB15 mRNA	2.0	M83679
Rat glutamate receptor subunit 2 (GLUR2) non-NMDA mRNA	2.0	M85035
Mouse p19 mRNA, complete cds. U17259	2.0	
Rat CDK108 mRNA.	2.0	Y17328
Rat neuronal olfactomedin-related ER localized protein (D2Sut1e) U03414	2.1	
Mouse A-X actin	2.1	J04181
Rat alpha-actin cardiac protein.	2.1	X80130
Rat calcium/calmodulin-dependent protein kinase type II alpha-subunit	2.3	J02942
membrane glycoprotein M6	2.3	S65735
Rat spectrin-like protein GTRAP41 AF225960	2.3	
Rat neuronal pentraxin precursor U18772	2.4	


Real Time PCR analysis



Original PCR test







Thank you