

[GC-09] **Propagation of UHE Photons through a Magnetized Large Structure**

Santabrata Das¹, Hyesung Kang², Dongsu Ryu, Jungyeon Cho³
ARCSEC, Pusan Nat'l Univ., Chungnam Nat'l Univ.

We investigate alignments between the spin axes of cosmic voids and the principal axes of nearby superclusters using the Millennium Run simulation of a LCDM cosmology. The concept of void spin was first introduced by Lee and Park in 2006 to quantify the tidal effect on voids from the surrounding matter distribution. Our numerical analysis reveals that the void spin axes are strongly aligned with the supercluster minor axes, but anti-aligned with the major axes, and have no correlations with the intermediate axes. We provide physical explanations to this numerical results on the basis of tidally induced correlations. It is expected that our work will provide a new insight into the characterization of the cosmic web on the largest scale.

[GC-10] **SHB Progenitor Life Time & Three Classes of GRB**

정수민¹, 이창환²
¹이화여자대학교 물리학과, ²부산 대학교 물리학과

Until 2005, afterglows of the short hard gamma ray burst (SHB) had not been detected because of observational difficulties in obtaining accurate localization. After detecting the afterglow of SHB, it allowed us to estimate the SHB progenitors. The most promising candidates of SHB progenitors are NS-NS mergers and NS-BH mergers. In this work, we calculated the local rate of SHBs to constrain the lifetime of SHB progenitors by using various star formation history models and presumed progenitor lifetimes. Our results confirm that the SHB progenitors are consistent with old population, about 6Gyr ~ 10Gyr lifetime. We applied this method to two possible subclasses in long soft gamma ray bursts. We found that three classes of gamma ray burst are more probable.