

Existence and Uniqueness of Solutions for the Fuzzy Differential Equations in n -Dimension Fuzzy Vector Space

n -차원 퍼지벡터공간에서의 퍼지미분방정식에 대한 해의 존재성과 유일성

권영철¹, 김외현¹, 박동근¹

동아대학교 자연과학대학 수학과
yckwun@dau.ac.kr, ouzzano@hanmail.net, dgpark@dau.ac.kr

Abstract

In this paper, we study the existence and uniqueness of solutions for the fuzzy differential equations in $(E_N)^n$ using by Banach fixed point theorem. $(E_N)^n$ is n -dimension fuzzy vector space.

Keywords : existence and uniqueness, fuzzy differential equations, n -dimension fuzzy vector space, Banach fixed point theorem

1. Introduction

Many authors have studied several concepts of fuzzy systems. Diamond and Kloeden [2] proved the fuzzy optimal control for the following system:

$$\dot{x}(t) = a(t)x(t) + u(t), \quad x(0) = x_0,$$

where $x(\cdot), u(\cdot)$ are nonempty compact interval-valued functions on E^1 . Kwun and Park [4] proved the existence of fuzzy optimal control for the nonlinear fuzzy differential system with nonlocal initial condition in E_N^1 using by Kuhn-Tucker theorems. Balasubramanian and Muralisankar [1] proved the existence and uniqueness of fuzzy solutions for the semilinear fuzzy integrodifferential equation with nonlocal initial condition. Recently, Park, Park and Kwun [6] find the sufficient condition of nonlocal controllability for the semilinear fuzzy integrodifferential equation with nonlocal initial condition.

In this paper, we study the the existence and uniqueness of solutions for the following fuzzy differential equations:

$$\frac{dx_i(t)}{dt} = a_i(t)x_i(t) + f_i(t, x_i(t)) \text{ on } E_N^i \quad (1)$$

$$x_i(0) = x_{0_i} \in E_N^i \quad (i = 1, 2, \dots, n) \quad (2)$$

where $a_i : [0, T] \rightarrow E_N^i$ is fuzzy coefficient, E_N^i is the set of all upper semi-continuously convex fuzzy numbers with $E_N^i \neq E_N^j$ ($i \neq j$), $f : [0, T] \times E_N^i \rightarrow E_N^i$

is a nonlinear regular fuzzy function and $x_{0_i} \in E_N^i$ is initial condition.

2. Preliminaries

A fuzzy subset (in short a fuzzy set) of R^n is function $u : R^n \rightarrow [0, 1]$. For each such fuzzy set u , we denote by $[u]^\alpha = \{x \in R^n : u(x) \geq \alpha\}$ for any $\alpha \in [0, 1]$, its α -level set.

Let u, v be fuzzy subset of R^n . It is well know that $[u]^\alpha = [v]^\alpha$ for each $\alpha \in [0, 1]$ implies $u = v$.

Let E^n denote the collection of all fuzzy sets of R^n that satisfy the following conditions:

- (1) u is normal, i.e., there exists an $x_0 \in R^n$ such that $u(x_0) = 1$;
- (2) u is fuzzy convex, i.e., $u(\lambda x + (1 - \lambda)y) \geq \min\{u(x), u(y)\}$ for any $x, y \in R^n, 0 \leq \lambda \leq 1$;
- (3) $u(x)$ is upper semi-continuous, i.e., $u(x_0) \geq \overline{\lim}_{k \rightarrow \infty} u(x_k)$ for any $x_k \in R^n$ ($k = 0, 1, 2, \dots$), $x_k \rightarrow x_0$;
- (4) $[u]^0$ is compact.

And we call $u \in E^n$ a n -dimension fuzzy number.

For any $u_i \in E, i = 1, 2, \dots, n$, we call the ordered one-dimension fuzzy number class u_1, u_2, \dots, u_n (i.e., the Cartesian product of one-dimension fuzzy number u_1, u_2, \dots, u_n) a n -dimension fuzzy vector, denote it as (u_1, u_2, \dots, u_n) ,

and call the collection of all n-dimension fuzzy vectors (i.e., the Cartesian product $\overbrace{E \times E \times \dots \times E}^n$) n-dimension fuzzy vector space, and denote it as $(E)^n$.

Definition 2.1[8] If $u \in E^n$, and $[u]^\alpha$ is a hyperrectangle, i.e., $[u]^\alpha$ can be represented by $\prod_{i=1}^n [u_{il}^\alpha, u_{ir}^\alpha]$, i.e., $[u_{1l}^\alpha, u_{1r}^\alpha] \times [u_{2l}^\alpha, u_{2r}^\alpha] \times \dots \times [u_{nl}^\alpha, u_{nr}^\alpha]$ for every $\alpha \in [0, 1]$, where $u_{il}^\alpha, u_{ir}^\alpha \in R$ with $u_{il}^\alpha \leq u_{ir}^\alpha$ when $\alpha \in (0, 1]$, $i = 1, 2, \dots, n$, then we call u a fuzzy n-cell number. And we denote the collection of all fuzzy n-cell numbers by $L(E^n)$.

Theorem 2.1[8] For any $u \in L(E^n)$ with $[u]^\alpha = \prod_{i=1}^n [u_{il}^\alpha, u_{ir}^\alpha]$ ($\alpha \in [0, 1]$), there exists a unique $(u_1, u_2, \dots, u_n) \in (E)^n$ such that $[u_i]^\alpha = [u_{il}^\alpha, u_{ir}^\alpha]$ ($i = 1, 2, \dots, n$ and $\alpha \in [0, 1]$).

Conversely, for any $(u_1, u_2, \dots, u_n) \in (E)^n$ with $[u_i]^\alpha = [u_{il}^\alpha, u_{ir}^\alpha]$ ($i = 1, 2, \dots, n$ and $\alpha \in [0, 1]$), there exists a unique $u \in L(E^n)$ such that $[u]^\alpha = \prod_{i=1}^n [u_{il}^\alpha, u_{ir}^\alpha]$ ($\alpha \in [0, 1]$).

Note 2.1 Theorem 2.1 indicates that fuzzy n-cell numbers and n-dimension fuzzy vectors can be represented each other, so $L(E^n)$ and $(E)^n$ may be regarded as identity. If $(u_1, u_2, \dots, u_n) \in (E)^n$ is the unique n-dimension fuzzy vector determined by $u \in L(E^n)$, then we denote $u = (u_1, u_2, \dots, u_n)$.

Definition 2.2[8] The complete metric D_L on $(E_N^i)^n$ is defined by

$$D_L(u, v) = \sup_{0 < \alpha \leq 1} d_L([u]^\alpha, [v]^\alpha) \\ = \sup_{0 < \alpha \leq 1} \max_{1 \leq i \leq n} \{|u_{il}^\alpha - v_{il}^\alpha|, |u_{ir}^\alpha - v_{ir}^\alpha|\}$$

for any $u, v \in (E_N^i)^n$, which satisfy $d_L(u+w, v+w) = d_L(u, v)$.

Definition 2.3 Let $u, v \in C([0, T] : (E_N^i)^n)$

$$H_1(u, v) = \sup_{0 \leq t \leq T} D_L(u(t), v(t)).$$

Definition 2.4[8] The derivative $x'(t)$ of a fuzzy process $x \in (E_N^i)^n$ is defined by

$$[x'(t)]^\alpha = \prod_{i=1}^n [(x_{il}^\alpha)'(t), (x_{ir}^\alpha)'(t)]$$

provided that is equation defines a fuzzy $x'(t) \in (E_N^i)^n$.

Definition 2.5[8] The fuzzy integral $\int_b^a x(t)dt$, $a, b \in [0, T]$ is defined by

$$\left[\int_b^a x(t)dt \right]^\alpha = \prod_{i=1}^n \left[\int_b^a x_{il}^\alpha(t)dt, \int_b^a x_{ir}^\alpha(t)dt \right]$$

provided that the Lebesgue integrals on the right exist.

3. Existence and Uniqueness

In this section we consider the existence and uniqueness of the fuzzy solution for the equation(1)-(2).

We defined by

$$a = (a_1, a_2, \dots, a_n),$$

$$x = (x_1, x_2, \dots, x_n),$$

$$f = (f_1, f_2, \dots, f_n),$$

and

$$x_0 = (x_{01}, x_{02}, \dots, x_{0n})$$

then

$$a, x, f, x_0 \in (E_N^i)^n$$

Instead of (1)-(2), we consider the following fuzzy integral equations in $(E_N^i)^n$.

$$\frac{dx(t)}{dt} = a(t)x(t) + f(t, x(t)) \quad (3)$$

$$x(0) = x_0 \quad (4)$$

with fuzzy coefficient $a : [0, T] \rightarrow (E_N^i)^n$, initial value $x_0 \in (E_N^i)^n$, and given nonlinear regular fuzzy function $f : [0, T] \times (E_N^i)^n \rightarrow (E_N^i)^n$ is satisfy global Lipschitz condition.

Definition 3.1 The fuzzy process $x : I \rightarrow (E_N^i)^n$ is a fuzzy solution of the equation (3)-(4) without inhomogeneous term if and only if

$$(x_{il}^\alpha)'(t) = \min\{a_{ij}^\alpha(t)x_{ik}^\alpha(t) : j, k = l, r\}$$

$$(x_{ir}^\alpha)'(t) = \max\{a_{ij}^\alpha(t)x_{ik}^\alpha(t) : j, k = l, r\}$$

$$x_{il}^\alpha(0) = x_{0il}^\alpha, \quad x_{ir}^\alpha(0) = x_{0ir}^\alpha.$$

Theorem 3.1 For every $x_0 \in (E_N^i)^n$

$$\frac{dx(t)}{dt} = a(t)x(t), \quad x(0) = x_0$$

has a unique fuzzy solution $x \in C([0, T] : (E_N^i)^n)$.

(H1) Assume that there exists a finite constant $k > 0$ such that

$$d_L([f(s, x(s))]^\alpha, [f(s, y(s))]^\alpha) \leq kd_L([x(s)]^\alpha, [y(s)]^\alpha)$$

for all $x(s), y(s) \in (E_N^i)^n$.

(H2) That is there exists a constant $c > 0$ such that $|S_{ij}^\alpha(t)| \leq c$ for all $t \in [0, T]$.

The equation (3)-(4) is related to the following fuzzy integral equations:

$$x(t) = S(t)x_0 + \int_0^t S(t-s)f(s, x(s))ds \quad (5)$$

$$x(0) = x_0 \quad (6)$$

Theorem 3.2 Let $T > 0$, satisfies hypotheses (H1) and (H2), for every $x_0 \in (E_N^i)^n$, (5)-(6) has a unique fuzzy solution $x \in C([0, T] : (E_N^i)^n)$.

References

- [1] P. Balasubramaniam and S. Muralisankar, *Existence and uniqueness of fuzzy solution for semilinear fuzzy integrodifferential equations with nonlocal conditions*, International J. Computer & Mathematics with applications, 47 (2004), 1115–1122.
- [2] P. Diamand and P.E. Kloeden, *Metric space of Fuzzy sets*, World Scientific (1994).
- [3] O. Kaleva, *Fuzzy differential equations*, Fuzzy set and Systems 24 (1987), 301–317.
- [4] Y.C. Kwun and D. G. Park, *Optimal control problem for fuzzy differential equations*, Proceedings of the Korea-Vietnam Joint Seminar (1998), 103–114.
- [5] M. Mizmoto and K. Tanaka, *Some properties of fuzzy numbers*, Advances in Fuzzy Sets Theory and applications, North-Holland Publishing Company (1979), 153–164.
- [6] J.H. Park, J.S. Park and Y.C. Kwun, *Controllability for the semilinear fuzzy integrodifferential equations with nonlocal conditions*, Lecture Notes in Artificial Intelligence 4223 (2006), 221–230.
- [7] S. Seikkala, *On the fuzzy initial value problem*, Fuzzy Sets and Systems 24 (1987), 319–330.
- [8] G. Wang, Y. Li and C. Wen *On fuzzy n-cell number and n-dimension fuzzy vectors*, Fuzzy Sets and Systems 158 (2007), 71–84.