

## [SF-01] Kinematics of H<sub>2</sub>O Masers in Massive Star-forming Region W51M

Jaesang Cho<sup>1</sup>, Yukitoshi Kan-Ya<sup>1,2</sup>, and Yong-Ik Byun<sup>1</sup>

<sup>1</sup>*Department of Astronomy, Yonsei University*

<sup>2</sup>*Korea Astronomy and Space Science Institute*

The W51M region contains numerous strong H<sub>2</sub>O maser sources, which are associated with outflows from massive proto-stars. The purpose of our research is to study the kinematics of maser spots in W51M as a probe to the site of multiple star formation and to contribute to the understanding of massive star formation. The distance measurement of W51M with annual parallax will be also possible as a prospect. For this study, we have acquired data sets of VLBI observations at 22 GHz with Japanese VERA (VLBI Exploration in Radio Astrometry) telescopes from 2003 to 2006. We are now proceeding with the imaging analysis for astrometric solutions, which will be the measure of internal kinematics in W51M. In this paper, we report the result from single-beam imaging data of an epoch previously not analyzed. Our results on W51M kinematics will be discussed and compared with the preliminary one reported by Kan-Ya et al. (2007).

## [SF-02] Discovery of a VeLLO in the "Starless" Dense Core L328

Chang Won Lee<sup>1,2</sup>, Tyler L. Bourke<sup>2</sup>, Philip C. Myers<sup>2</sup>, Mike Dunham<sup>3</sup>, Neal Evans<sup>3</sup>, Youngung Lee<sup>1</sup>, Tracy Huard<sup>4</sup>, Jingwen Wu<sup>2</sup>, Robert Gutermuth<sup>2</sup>, Mi-Ryang Kim<sup>1</sup>, Hyun Woo Kang<sup>1</sup>

<sup>1</sup>*Korea Astronomy & Space Science Institute*

<sup>2</sup>*Harvard-Smithsonian Center for Astrophysics*

<sup>3</sup>*Department of Astronomy, The University of Texas at Austin,*

<sup>4</sup>*Department of Astronomy, The University of Maryland*

We report the discovery of a Very Low Luminosity Object (VeLLO) in the "starless" dense core L328, using the Spitzer Space Telescope and ground based observations from near-infrared to millimeter wavelengths.

The Spitzer 8 mm image indicates that L328 consists of three subcores of which the smallest one may harbor a source, L328-IRS while two other subcores remain starless.

L328-IRS is a Class 0 protostar according to its bolometric temperature (44 K) and the high fraction (~72 %) of its luminosity emitted at sub-millimeter wavelengths. Its inferred "internal luminosity" (0.04 - 0.06 L<sub>⊙</sub>) using a radiative transfer model under the most plausible assumption of its distance as 200 pc is much fainter than for a typical protostar, and even fainter than other VeLLOs studied previously.

Note, however, that its inferred luminosity may be uncertain by a factor of 2-3 if we consider two extreme values of the distance of L328-IRS (125 or 310 pc).

Low angular resolution observations of CO do not show any clear evidence of a molecular

outflow activity. But broad line widths toward L328, and Spitzer and near-infrared images showing nebulosity possibly tracing an outflow cavity, strongly suggest the existence of outflow activity. Provided that an envelope of at most ~0.1 M<sub>⊙</sub> is the only mass accretion reservoir for L328-IRS, and the star formation efficiency is close to the canonical value ~30%, and L328-IRS has not yet accreted more than ~0.05 M<sub>⊙</sub>, at the assumed distance of 200 pc, L328-IRS is destined to be a brown dwarf.