

**■ Session : 우주환경 II**  
**4월 30일(금) 13:30 - 15:30 제3발표장**

**[IV-3-1] 삼축 MI 자력계 설치 및 운용**  
 최규철<sup>1,2</sup>, 이재진<sup>1</sup>, 황정아<sup>1</sup>, 조경석<sup>1</sup>, 박영득<sup>1</sup>, 이대영<sup>2</sup>  
<sup>1</sup>한국천문연구원, <sup>2</sup>충북대학교 천문우주학과

한국천문연구원 우주과학연구본부 태양우주환경연구그룹은 일본 Tierra Tecnica사의 RFP-523C Overhauser Proton 자력계와 MISYS-09 삼축 MI 자력계를 2009년 11월에 보현산천문대 태양 망원경동에 구축하였다. 한국천문연구원은 이미 2007년 11월에 RFP-523C Fluxgate 자력계를 보현산천문대 태양망원경동에 구축하여 K 지수 산출 등의 우주환경예·경보 연구에 활용하고 있다. Fluxgate 자력계는 지자기 3축 성분의 변화량을 측정하는 장비이고 이번에 설치한 Overhauser Proton 자력계는 지자기의 총 자기장을 측정하는 장비이다. 삼축 MI 자력계는 지자기장의 wave를 측정하는 장비이다. 기존에 설치한 Fluxgate 자력계와 새로 설치한 Proton 자력계, 삼축 MI 자력계를 연계하여 운용할 경우 우주환경예 의한 지자기장 변화량의 측정 정밀도가 향상되고 지자기장을 효율적으로 관측할 수 있다. 보현산천문대에 구축한 각각의 자력계가 측정한 지자기 자료들은 S-FTP와 Socket 통신을 이용하여 대전에 있는 한국천문연구원 태양우주환경연구 그룹의 데이터 서버로 실시간으로 전송되어 저장되고 있다. 데이터 서버로 전송된 지자기 측정 자료들은 한국천문연구원 우주환경감시실에서 모니터링하고 있다.

**[IV-3-2] Morning-afternoon asymmetry of geosynchronous magnetic field variations during geomagnetic sudden commencements**  
 Jong-Sun Park, Khan-Hyuk Kim, Suk-Kyung Sung, Dong-Hun Lee  
*School of Space Research, Kyung Hee University, Kyunggi, Korea*

It has been reported that geosynchronous magnetopause crossings are more frequently observed in the prenoon sector than in the postnoon sector, indicating a dawn-dusk magnetopause asymmetry during extreme solar wind conditions. Motivated by these observations, we investigate geosynchronous magnetic field variations normalized to SYM-H when sudden commencements (SC) are observed on the ground. From a statistical analysis of the geosynchronous magnetic field responses to SC events from 1997 to 2006, we found that the normalized SC amplitude at geosynchronous orbit is larger in the morning sector than in the afternoon sector. In order to examine if this morning-afternoon asymmetry at geosynchronous orbit occurs only during disturbed geomagnetic conditions, we compared the geosynchronous magnetic field strength obtained in the morning and afternoon during undisturbed intervals ( $K_p < 3$ ). We found that the asymmetry appears under undisturbed geomagnetic conditions and it is not due to solar wind aberration. This indicates that the morning-afternoon asymmetry was not strongly affected by

changes in solar wind condition. Using solar wind data, we discuss what causes the morning-afternoon asymmetry at geosynchronous orbit.

**[IV-3-3] THEMIS Pi2 observations near dawn and dusk sectors in the inner magnetosphere**  
 Hyuck-Jin Kwon<sup>1</sup>, Khan-Hyuk Kim<sup>1</sup>, Dong-Hun Lee<sup>1</sup>, K. Takahashi<sup>2</sup>, Young-Deuk Park<sup>3</sup>, J. W. Bonnell<sup>4</sup>  
<sup>1</sup>*School of Space Research, Kyung Hee University,*  
<sup>2</sup>*APL, Johns Hopkins University, Laurel, Maryland, USA*  
<sup>3</sup>*Korea Astronomy and Space Science Institute, Korea*  
<sup>4</sup>*Space Science Laboratory, University of California, Berkeley, California, USA*

The most frequently suggested source for Pi2 pulsations in the inner magnetosphere ( $L < 4$ ) is standing fast-mode waves trapped in the plasmasphere (i.e., plasmaspheric resonances). They have been considered as the source of low-latitude Pi2 pulsations. The plasmaspheric resonance model suggests that compressional fast-mode waves can be detected at all local times inside the plasmasphere provided reflection of the wave energy is efficient. Until now, however, there are no reports about compressional Pi2s observed in the dayside inner magnetosphere. That is, there is longitudinal limit of inner magnetosphere. In February 2008, THEMIS probes were near dawn and/or dusk sides, which are the transition regions between the nightside and dayside, in the inner magnetosphere ( $L = 2-4$ ) when low-altitude Pi2s were identified at Bohyun ( $L = 1.35$ ) station in Korea. Using the THEMIS electric field data, we examined if Pi2s are excited by longitudinally localized disturbances. We found that compressional Pi2s having high coherence with a low-latitude Pi2 pulsation occur on dawnside. However, any compressional pulsations in the Pi2 frequency band were not detected on duskside. This indicates that compressional Pi2s disappear near the duskside. Our observations are discussed with spatial plasmaspheric structure and possible Pi2 mechanisms.

**[IV-3-4] Statistical properties of the fast flows accompanied by dipolarization in the near-Earth tail**  
 Hyun-Sook Kim<sup>1</sup>, Dae-Young Lee<sup>2</sup>, Byung-Ho Ahn<sup>3</sup>  
<sup>1</sup>*Astronomy & Atmospheric Science, Kyungpook National University, Korea*  
<sup>2</sup>*Astronomy & Space Science, Chungbuk National University, Korea*  
<sup>3</sup>*Earth Science Education, Kyungpook National University, Korea*

Using magnetic field and plasma moment data obtained by THEMIS satellites(A, D, and E), we selected 203 fast flow events accompanied by dipolarization in the near-Earth region(  $X(\text{GSM}) = -7 \sim -12 \text{ RE}$ ) and statistically examined