

Germination model of ‘Namdo’ Variety of Garlic with Temperature

Kyung Hwan Moon^{1*}, Hyeong-Ho Seo¹, Kyung San Choi¹, Jae-Ho Joa¹, Young-Soon Kwon²

¹*Agricultural Research Center for Climate Change, NIHHS, RDA, 1696 Odeung-dong, Jeju-si, Jeju 690-150*

²*National Institute of Horticultural & Herbal Science, RDA, 203 Cheoncheon-ro, Jangan-gu, Suwon 440-706*

A model for estimating germination rate of garlic(var. Namdo) with temperature was made by chamber experiment. We checked emergence rate after planting garlic cloves with five constant temperature treatments of 10°C, 15°C, 20°C, 25°C and 30°C. From 50% emergence days, we calculate representative germination rates for every temperature. The relationship of germination rate and temperature was elucidated by curve fitting with exponential type function. 13.23°C was the optimum temperature for garlic germination. We could calculate germination rate for every day with daily mean temperature from the temperature-germination rate relationship. Total germination percentage could estimated from accumulating daily germination rate. We also make a excel program for calculation of garlic germination rate. Application result of the germination model to field condition also suggested and limitation of the model also discussed.

* Corresponding Author: khmoon9000@gmail.com