

**[☞SE-44] Incorporation of Electromagnetic Ion cyclotron wave into
Radiation Belt environment model**

Suk-bin Kang¹, Eunjin Choi^{1,2}, Junga Hwang², Kyung-chan Kim², Jaejin Lee²,
Mei-ching Fok³, Kyoungwook Min¹, Cheongrim Choi¹, Young-Deuk Park²
¹*Korea Advanced Institute of Science and Technology (KAIST)*
²*Korea Astronomy and Space Science Institute (KASI)*
³*NASA Goddard Space Flight Center, Maryland*

Radiation Belt Environment (RBE) model has developed to understand radiation belt dynamics as it considers whistler mode hiss and chorus waves which is responsible for relativistic electron acceleration and precipitation. Recently, many studies on electron loss by pitch-angle scattering have reported that electromagnetic ion cyclotron (EMIC) wave is also responsible for main loss mechanism in dusk and equatorial region. Here, we attempt to incorporate EMIC into RBE model simulation code to understand more detailed physical dynamics in Radiation belt environment. We compare this developed model to data during storm events where both of electron loss and EMIC waves were detected

**[☞SE-45] Propagation characteristics of CMEs associated magnetic
clouds and ejecta**

Roksoon Kim¹, Nat Gopalswamy², Kyungsuk Cho¹, Yongjae Moon³,
Seiji Yashiro^{2,4}, and Youngdeuk Park¹
¹*KASI*, ²*NASA/GSFC*, ³*Kyunghee University*, ⁴*Catholic University of America*

We have investigated the characteristics of magnetic cloud (MC) and ejecta (EJ) associated coronal mass ejections (CMEs) based on the assumption that all CMEs have a flux rope structure. For this, we used 54 CMEs and their interplanetary counterparts (interplanetary CMEs: ICMEs) that constitute the list of events used by the NASA/LWS Coordinated Data Analysis Workshop (CDAW) on CME flux ropes. We considered the location, angular width, and speed as well as the direction parameter, D. The direction parameter quantifies the degree of asymmetry of the CME shape, and shows how closely the CME propagation is directed to Earth. For the 54 CDAW events, we found several properties of the CMEs as follows: (1) the average value of D for the 23 MCs (0.62) is larger than that for the 31 EJs (0.49), which indicates that the MC-associated CMEs propagate more directly to the Earth than the EJ-associated CMEs; (2) comparison between the direction parameter and the source location shows that the majority of the MC-associated

CMEs are ejected along the radial direction, while many of the EJ-associated CMEs are ejected non-radially; (3) the mean speed of MC-associated CMEs (946 km/s) is faster than that of EJ-associated CMEs (771 km/s). For seven very fast CMEs (>1500 km/s), all CMEs with large D (>0.4) are associated with MCs and the CMEs with small D are associated with EJs. From the statistical analysis of CME parameters, we found the superiority of the direction parameter. Based on these results, we suggest that the CME trajectory essentially decides the observed ICME structure.