

Medium-deep Survey (IMS), a moderately wide (120 deg<sup>2</sup>) and deep (JAB ~ 22.5 - 23 mag) near-infrared imaging survey. Combining this with the optical (ugriz) imaging data from the CFHT Legacy Survey (CFHTLS), we have identified more than 10 faint quasar candidates at  $z \sim 6$  in the IMS field by using multiple color selection criteria. From now on, we will perform spectroscopic confirmations of these faint quasar candidates with IMACS on the Magellan Baade Telescope at Las Campanas Observatory and GMOS on the Gemini South Telescope at Gemini Observatory. The confirmed quasars will be used to constrain the faint-end slope of the quasar luminosity function at  $z \sim 6$  and calculate the ratio of quasar ionizing flux to required flux for reionization of the universe. Moreover, these confirmed quasars will be followed up with near-infrared spectroscopy to determine their black hole masses and Eddington ratios to check the rapidness of their growth.

### [ㄹ GC-05] Demographics of Isolated Galaxies along the Hubble Sequence

Hong-geun Kim, Jongwon Park, Seong-woo Seo and Sukyoung K. Yi  
*Department of Astronomy, Yonsei University, Republic of Korea*

Isolated galaxies in low-density regions are significant in the sense that they are least affected by the hierarchical pattern of galaxy growth and interactions with perturbers at least for the last few Gyr. To form a comprehensive picture of the star formation history of isolated galaxies, we construct a catalog of isolated galaxies and their comparison sample in relatively denser environments. The galaxies are drawn from SDSS DR7 in the redshift range of  $0.025 < z < 0.044$ . We performed visual inspection and classified their morphology following the Hubble classification scheme. We have investigated the color-magnitude diagram and found elliptical and unbarred spiral galaxies in isolated systems are relatively fainter and bluer than those in denser regions. For the spectroscopic study, we make use of the OSSY catalog (Oh et al. 2011). Our analysis on the absorption-line properties based on the comparison with stellar population models suggests that isolated elliptical galaxies are likely to be younger and metal poorer, while isolated Sc-type galaxies seem to have older luminosity-weighted ages, than their high-density counterpart. In addition, according to the BPT diagnostics, early-type galaxies among isolated galaxies are rather evenly classified into star forming,

composite, Seyfert and LINER, whereas their comparisons are mainly populated in the LINER region. On the other hand, late-type galaxies do not show any prominent difference. We discuss the evolutionary histories of isolated galaxies in the context of the standard  $\Lambda$ CDM cosmology.

### [ㄹ GC-06] No asymmetric outflows from Sagittarius A\* during the pericenter passage of the gas cloud G2

Jong-Ho. Park<sup>1</sup>, Sascha Trippe<sup>1</sup>, Thomas Krichbaum<sup>2</sup>, Jae-Young Kim<sup>1</sup>, Motoki Kino<sup>3</sup>, Alessandra Bertarini<sup>2,4</sup>, Michael Bremer<sup>5</sup>, and Pablo de Vicente<sup>6</sup>,

<sup>1</sup>*Seoul National University,*

<sup>2</sup>*Max-Planck-Institut fur Radioastronomie,*

<sup>3</sup>*KASI,*

<sup>4</sup>*Bonn University,*

<sup>5</sup>*IRAM,*

<sup>6</sup>*Astronomical Centre of Yebes*

The gas cloud G2 falling toward Sagittarius A\* (Sgr A\*), the supermassive black hole at the center of the Milky Way, is supposed to provide valuable information on the physics of accretion flows and the environment of the black hole. We observed Sgr A\* with four European stations of the Global Millimeter Very Long Baseline Interferometry Array (GMVA) at 86 GHz on 1 October 2013 when parts of G2 had already passed the pericenter. We searched for possible transient asymmetric structure — such as jets or winds from hot accretion flows — around Sgr A\* caused by accretion of material from G2. The interferometric closure phases (which are zero if the spatial brightness distribution of the target is symmetric, and deviate from zero otherwise) remained zero within errors during the observation time. We thus conclude that Sgr A\* did not show significant asymmetric (in the observer frame) outflows in late 2013. Using simulations, we constrain the size of the outflows that we could have missed to  $\approx 2.5$  mas along the major axis,  $\approx 0.4$  mas along the minor axis of the beam, corresponding to approximately 232 and 35 Schwarzschild radii, respectively; we thus probe spatial scales on which the jets of radio galaxies are suspected to convert magnetic into kinetic energy. As probably less than 0.2 Jy of the flux from Sgr A\* can be attributed to accretion from G2, one finds an effective accretion rate  $\eta \dot{M} \lesssim 1.5 \times 10^9$  kg/s  $\approx 7.7 \times 10^{-9} M_{\text{earth}} / \text{yr}$  for material from G2. Exploiting the kinetic jet power-accretion power relation of radio galaxies, one finds that the rate of accretion of matter that ends up in jets is

limited to  $\dot{M} \lesssim 10^{17}$  kg/s  $\approx 0.5 M_{Earth} / \text{yr}$  less than about 20% of the mass of G2. Accordingly, G2 appears to be largely stable against loss of angular momentum and subsequent (partial) accretion at least on time scales  $\lesssim 1$  year. [Park et al. 2015 under review by A&A]

### [☞ GC-07] A Test of Correspondence Model with the HorizonRun 4 Simulation

Jisook Park<sup>1,2</sup>, Juhan Kim<sup>3</sup>, Changbom Park<sup>2</sup>, and Sungsoo S. Kim<sup>1,4</sup>

<sup>1</sup>*School of Space Research, Kyung Hee University*

<sup>2</sup>*School of Physics, Korea Institute for Advanced Study*

<sup>3</sup>*Center for Advanced Computation, Korea Institute for Advanced Study*

<sup>4</sup>*Department of Astronomy and Space Science, Kyung Hee University*

'The one to one correspondence model' defines the relation between a dark matter halo (DM halo) and a galaxy. A basic assumption of this model is that a more massive DM subhalo hosts a brighter galaxy. In a more improved version of the model we may be able to assign a mock galaxy with a morphological type. In this study, we are building a mock galaxy catalog using massive halo merging trees from the Horizon Run 4. We test various merging models to calculate the merging time scale of a subhalo along its merging tree. And we obtain the halo mass functions for major subhalos and satellite subhalos, separately, and compare them with the observed luminosity functions of major galaxies and satellite galaxies from the SDSS group catalog. Furthermore, we are going to make a range of mock galaxy catalogs and investigate their properties, such as spatial distributions, environmental effects, and morphologies.

### [☞ GC-08] Optical 3D Spectroscopic Survey on Gas Outflows in Type 2 AGNs

Hyun-Jin Bae<sup>1</sup>, Jong-Hak Woo<sup>2</sup>, Marios Karouzos<sup>2</sup>, Elena Gallo<sup>3</sup>, Yue Shen<sup>4</sup>, Helene Flohic<sup>5</sup>

<sup>1</sup>*Department of Astronomy, Yonsei University,*

<sup>2</sup>*Department of Physics and Astronomy, Seoul National University,*

<sup>3</sup>*Department of Astronomy, University of Michigan, USA,*

<sup>4</sup>*Carnegie Institution for Science, USA,*

<sup>5</sup>*University of the Pacific, USA*

Strong outflows from active galactic nuclei (AGNs) may play a crucial role in galaxy evolution.

Integral-field spectroscopy (IFS) is the most powerful tool to study the detailed kinematics of AGN outflows. We present the on-going optical 3D spectroscopic survey of ionized gas outflows. Type 2 AGN sample is uniquely selected from SDSS DR7 with a luminosity-limit (i.e.,  $L[\text{O III}] > 10^{41.5}$  erg/s) as well as strong kinematic signatures of ionized gas outflows ( $[\text{O III}]$  velocity shift  $> \sim 200$  km/s or  $[\text{O III}]$  velocity dispersion (FWHM)  $> 1000$  km/s), defining an extremely rare population ( $< \sim 0.5\%$ ). Thus, these AGNs with strong outflow signatures are one of the best suites for investigating AGN feedback. The IFS observations cover several kpc scales for the central region of the host galaxies, providing a detailed information of the kinematics and geometry of the gas outflows. In this contribution, we report the current status of the survey and the preliminary results on gas kinematics of 18 AGNs, based on the Magellan/IMACS-IFU and the VLT/VIMOS data.

### [☞ GC-09] Surface photometry and Structural properties of nearby dwarf galaxies.

Mira Seo, Hong Bae Ann  
*Pusan National University*

We present 2D-photometric decompositions of  $\sim 1,200$  nearby dwarf galaxies. Our representative sample is derived from 'A catalog of Visually classified galaxies in the Local Universe'(Ann, Seo and Ha APJS,,2015) of which galaxy morphological types are determined by visual inspection of color images using the Sloan Digital Sky Survey data release 7. In this catalog, dwarf galaxies were divided into 5 subtypes : dS0, dE, dSph, dEbc, dEblue with distinction of the presence of nucleation in dE, dSph, and dS0. The dSph types are less brighter than other types, and galaxies with nuclei are slightly brighter than those with no nuclei in the same types. Sersic index  $n$  have a range 1~1.5, and  $dE_{\text{un}}$  and  $dSph_{\text{un}}$  galaxies have  $n$  less than 1, and  $dSph_{\text{n}}$  galaxies have largest values. We performed two-dimensional decomposition of galaxies using GALFIT, and analyzed their structural components, and residual features which are seen in the residual image.

### [☞ GC-10] The narrow emission-line properties of radio-loud AGNs in the SDSS archive

Donghoon Son and Jong-Hak Woo  
*Astronomy Program, Department of Physics and*