

포스터 발표 소록

외부은하 / 은하단

[포 GC-01] Black Hole Activities of Red Active Galactic Nuclei

Dohyeong Kim, Myungshin Im, Jong-Hak Woo
Astronomy Program, Department of Physics and Astronomy, Seoul National University

We investigate black hole (BH) activities of 16 red active galactic nuclei (AGNs). The 16 red AGNs selected by red colors in optical through near-infrared (NIR) and radio detection. In order to derive BH activities of the red AGNs, we use P β line with NIR spectra obtained by the SpeX on the IRTF. The P β line suffers from dust extinction less than UV/optical BH mass estimators. We compared Eddington ratios of the red AGNs and "normal" AGNs, and the Eddington ratios of red AGNs are significantly higher than those of "normal" AGNs. The result is consistent with a scenario that red AGNs are the intermediate population between star forming galaxies and "normal" AGNs, and BHs of red AGNs are very active and grow rapidly in such a stage.

[포 GC-02] Spectroscopic Studies of Wolf-Rayet galaxies

Myunghak Kim and Hyunjin Shim
Department of Earth Science Education, Kyungpook National University

Wolf-Rayet galaxies, galaxies which show spectral features produced by Wolf-Rayet stars, are thought to be the place of recent massive star formation since the Wolf-Rayet phase covers relatively short timescale in the life of massive O and B type stars. Studying these galaxies provides a unique chance to understand how massive star formation occurs in a galaxy within a short timescale. In this work, we present the intermediate resolution optical spectra of 12 Wolf-Rayet galaxies obtained using longslit spectrograph on Bohyunsan Optical Astronomy Observatory. We derived the emission line ratios for a number of star-forming knots in each Wolf-Rayet galaxy. Star formation properties in these galaxies are discussed.

[포 GC-03] Mid-IR Luminosity Functions of Local Galaxies in the North Ecliptic Pole Field.

Seong Jin Kim¹, Hyung Mok Lee², Woong-Seob Jeong¹, and the NEP team members
¹*Korea Astronomy and Space Science Institute,*
²*Seoul National University*

We present the mid-infrared (MIR) luminosity function (LF) of local ($z < 0.3$) star forming (SF) galaxies based on the AKARI's NEP-Wide Survey data. We utilized a combination of the NEP-Wide point source catalogue containing a large number (114,000) of infrared (IR) sources distributed over the wide (5.4 sq. deg) field and spectroscopic redshift (z) data for 1790 selected targets obtained by optical follow-up surveys with MMT/Hectospec and WIYN/Hydra. The AKARI's continuous 2 ~ 24 μ m wavelength coverage and the spectroscopic redshifts for sample galaxies enable us to derive accurate spectral energy distributions (SEDs) in the mid-infrared.

We carried out SED-fit analysis and employed 1/V_{max} method to derive the mid-IR (e.g., 8 μ m, 12 μ m, and 15 μ m rest-frame) luminosity functions. Our results for local galaxies from the NEP region generally consistent with various previous works for other fields over wide luminosity ranges. The comparison with the results of the NEP-Deep data implies the luminosity evolution from higher redshifts towards the present epoch. We attempted to fit our derived LFs to the double power-laws and present the resulting power indices. We also examined the correlation between mid-IR luminosity and total IR luminosity.

[포 GC-04] IMS High- z Quasar Survey - Faint $z \sim 6$ Quasar Candidates in IMS Fields

Yongjung Kim, Myungshin Im, Yiseul Jeon and IMS team
CEOU / Dept. of Physics and Astronomy, Seoul National University

Over the last decade, more than 50 quasars have been discovered at redshift about 6 when reionization of the universe occurred. However, most of them are luminous quasars ($z_{AB} < 21$ mag), implying that such a biased quasar sample, which cannot represent the entire population of quasars at $z \sim 6$, is not enough to understand the properties of quasars in the early universe. Recently, we have been performing the Infrared

Medium-deep Survey (IMS), a moderately wide (120 deg²) and deep (JAB ~ 22.5 - 23 mag) near-infrared imaging survey. Combining this with the optical (ugriz) imaging data from the CFHT Legacy Survey (CFHTLS), we have identified more than 10 faint quasar candidates at $z \sim 6$ in the IMS field by using multiple color selection criteria. From now on, we will perform spectroscopic confirmations of these faint quasar candidates with IMACS on the Magellan Baade Telescope at Las Campanas Observatory and GMOS on the Gemini South Telescope at Gemini Observatory. The confirmed quasars will be used to constrain the faint-end slope of the quasar luminosity function at $z \sim 6$ and calculate the ratio of quasar ionizing flux to required flux for reionization of the universe. Moreover, these confirmed quasars will be followed up with near-infrared spectroscopy to determine their black hole masses and Eddington ratios to check the rapidness of their growth.

[ㄷ GC-05] Demographics of Isolated Galaxies along the Hubble Sequence

Hong-geun Kim, Jongwon Park, Seong-woo Seo and Sukyoung K. Yi
Department of Astronomy, Yonsei University, Republic of Korea

Isolated galaxies in low-density regions are significant in the sense that they are least affected by the hierarchical pattern of galaxy growth and interactions with perturbers at least for the last few Gyr. To form a comprehensive picture of the star formation history of isolated galaxies, we construct a catalog of isolated galaxies and their comparison sample in relatively denser environments. The galaxies are drawn from SDSS DR7 in the redshift range of $0.025 < z < 0.044$. We performed visual inspection and classified their morphology following the Hubble classification scheme. We have investigated the color-magnitude diagram and found elliptical and unbarred spiral galaxies in isolated systems are relatively fainter and bluer than those in denser regions. For the spectroscopic study, we make use of the OSSY catalog (Oh et al. 2011). Our analysis on the absorption-line properties based on the comparison with stellar population models suggests that isolated elliptical galaxies are likely to be younger and metal poorer, while isolated Sc-type galaxies seem to have older luminosity-weighted ages, than their high-density counterpart. In addition, according to the BPT diagnostics, early-type galaxies among isolated galaxies are rather evenly classified into star forming,

composite, Seyfert and LINER, whereas their comparisons are mainly populated in the LINER region. On the other hand, late-type galaxies do not show any prominent difference. We discuss the evolutionary histories of isolated galaxies in the context of the standard Λ CDM cosmology.

[ㄷ GC-06] No asymmetric outflows from Sagittarius A* during the pericenter passage of the gas cloud G2

Jong-Ho. Park¹, Sascha Trippe¹, Thomas Krichbaum², Jae-Young Kim¹, Motoki Kino³, Alessandra Bertarini^{2,4}, Michael Bremer⁵, and Pablo de Vicente⁶,

¹*Seoul National University,*

²*Max-Planck-Institut fur Radioastronomie,*

³*KASI,*

⁴*Bonn University,*

⁵*IRAM,*

⁶*Astronomical Centre of Yebes*

The gas cloud G2 falling toward Sagittarius A* (Sgr A*), the supermassive black hole at the center of the Milky Way, is supposed to provide valuable information on the physics of accretion flows and the environment of the black hole. We observed Sgr A* with four European stations of the Global Millimeter Very Long Baseline Interferometry Array (GMVA) at 86 GHz on 1 October 2013 when parts of G2 had already passed the pericenter. We searched for possible transient asymmetric structure — such as jets or winds from hot accretion flows — around Sgr A* caused by accretion of material from G2. The interferometric closure phases (which are zero if the spatial brightness distribution of the target is symmetric, and deviate from zero otherwise) remained zero within errors during the observation time. We thus conclude that Sgr A* did not show significant asymmetric (in the observer frame) outflows in late 2013. Using simulations, we constrain the size of the outflows that we could have missed to ≈ 2.5 mas along the major axis, ≈ 0.4 mas along the minor axis of the beam, corresponding to approximately 232 and 35 Schwarzschild radii, respectively; we thus probe spatial scales on which the jets of radio galaxies are suspected to convert magnetic into kinetic energy. As probably less than 0.2 Jy of the flux from Sgr A* can be attributed to accretion from G2, one finds an effective accretion rate $\eta \dot{M} \lesssim 1.5 \times 10^9$ kg/s $\approx 7.7 \times 10^{-9} M_{\text{earth}} / \text{yr}$ for material from G2. Exploiting the kinetic jet power-accretion power relation of radio galaxies, one finds that the rate of accretion of matter that ends up in jets is