

components, respectively, as well as the orbital eccentricity distribution. There are notable gradients in the V velocity over [Fe/H] in both populations: -23 km s⁻¹ dex⁻¹ for the thin disk and +44 km s⁻¹ dex⁻¹ for the thick disk. The velocity dispersion of the thick disk decrease with increasing [Fe/H], while the velocity

Gungwon Kang³, Chunglee Kim², Whansun Kim⁴, John J. Oh⁴, Sang Hoon Oh⁴, Chan Park³, Edwin J. Son⁴, Ho Jung Paik⁵

¹*Korea Research Institute of Standards and Science*, ²*Korea Astronomy & Space Science Institute*, ³*Korea Institute of Science and Technology Information*, ⁴*National Institute for Mathematical Sciences*, ⁵*U. of Maryland (USA)*

New Frontier of Gravitational Wave Research

[ㄱ GW-01] Superconducting Low-frequency Gravitational-wave Telescope (SLGT): pilot study status report

Chunglee Kim¹, Sang-Hyeon Ahn¹, Yeong-Bok Bae¹, Gungwon Kang², Whansun Kim³, John J. Oh³, Sang Hoon Oh³, Chan Park², Edwin J. Son³, Yong Ho Lee⁴, Ho Jung Paik⁵

¹*Korea Astronomy & Space Science Institute*, ²*Korea Institute of Science and Technology Information*, ³*National Institute for Mathematical Sciences*, ⁴*Korea Research Institute of Standards and Science*, ⁵*U. of Maryland (USA)*

The discovery of GW150914, black hole - black hole merger via gravitational waves (GWs) opened a new window to observe the Universe. GW frequencies from heavenly bodies and early Universe are expected to span between sub-nHz up to kHz. At present, GW detectors on Earth (LIGO, Virgo, KAGRA, LIGO-India) aims frequency ranges between 10-2000 Hz. The space-borne GW detector and Pulsar Timing Array targets mHz and nHz sources. Starting in March 2017, the KKN (KASI-KISTI-NIMS) collaboration launched a pilot study of SLGT (Superconducting Low-frequency Gravitational-wave Telescope). This project is funded by NST (Korea Institute of Science and Technology). The main detection bands expected for SLGT ranges between 0.1-10Hz, which is complementary of LIGO-type detectors and LISA for multi-band GW observation. We will present an overview of the SLGT project and report the status of the NST pilot study. We will also present prospective of GW astronomy with SLGT.

[초 GW-02] Development of Superconducting Low-frequency Gravitational-wave Telescope (SLGT): Technical Challenge and Feasibility

Yong Ho Lee¹, Sang-Hyeon Ahn², Yeong-Bok Bae²,

Recent success of gravitational wave (GW) detection by LIGO opened a new window to expand our understanding of the Universe. In addition to LIGO, several other developments are going on or under planning. However, each of these detectors has a specific sensitive frequency range. There is a missing frequency band, 0.1-10 Hz, where detectors loose sensitivity significantly due to Newtonian noise on the Earth. We introduce a plan to develop a Superconducting Low-frequency Gravitational-wave Telescope (SLGT), which can observe massive black holes in 0.1-10 Hz. The SLGT system consists of magnetically levitated six test masses, superconducting quantum interference devices (SQUIDs), rigid support frame, cooling system, vibration isolation, and signal acquisition. By taking the advantage of nearly quantum-limited low-noise SQUIDs and capacitor bridge transducers, SLGT's detection sensitivity can be improved to allow astrophysical observation of black holes in cosmological distances. We present preliminary design study and expected sensitivity, and its technical feasibility.

[ㄱ GW-03] Optical/NIR Follow-up Observation of GW Sources

Myungshin Im¹, Hyung Mok Lee¹, Changsu Choi¹, Joonho Kim¹, Lim Gu¹, Hyun-Il Sung², Yeong-Beom Jeon², Seung-Li Kim², Chung-Uk Lee², Soojong Pak³ Shuhrat Eghamberdiev⁴

¹*Astronomy Program/CEO, Dept. of Physics & Astronomy, Seoul National University*
²*Korea Astronomy & Space Science Institute*
³*School of Space Research and Institute of Natural Sciences, Kyunghee University*
⁴*Ulugh Beg Astronomical Institute, Uzbekistan*

Identification of gravitational wave (GW) sources in electromagnetic (EM) wave observations is important because it enables us to understand the property of the GW-emitting sources/mechanisms much better than the GW detection. For that reason, a large number of astronomers are working on observations to identify the position and the nature of GW sources. We give a short