

[7 IM-02] High-resolution ALMA Study of the Proto-Brown-Dwarf Candidate L328-IRS

Chang Won Lee^{1,2}, Gwanjeong Kim^{1,2,3}, Philip C. Myers⁴, Masao Saito⁵, Shinyoung Kim^{1,2}, Woojin Kwon^{1,2}, A-Ran Lyo¹, Archana Soam¹, & Mi-Ryang Kim¹

¹*Korea Astronomy & Space Science Institute,*

²*Korea University of Science & Technology*

³*Nobeyama Radio Observatory, NAOJ*

⁴*Harvard-Smithsonian Center for Astrophysics*

⁵*NAOJ*

We present our observational attempts to precisely measure the central mass of a proto-brown dwarf candidate, L328-IRS, in order to investigate whether L328-IRS is in the substellar mass regime.

Observations were made for the central region of L328-IRS with the dust continuum and CO isotopologue line emission at ALMA band 6, discovering the detailed outflow activities and a deconvolved disk structure of a size of $\sim 87 \text{ AU} \times \sim 37 \text{ AU}$. We investigated the rotational velocities as a function of the disk radius, finding that its motions between 130 AU and 60 AU are partially fitted with a Keplerian orbit by a stellar object of $\sim 0.30 M_{\odot}$, while the motions within 60 AU do not follow any Keplerian orbit at all. This makes it difficult to lead a reliable estimation of the mass of L328-IRS.

Nonetheless, our ALMA observations were useful enough to well constrain the inclination angle of the outflow cavity of L328-IRS as $\sim 66^{\circ}$ degree, enabling us to better determine

the mass accretion rate of $\sim 8.9 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$. From assumptions that the internal luminosity of L328-IRS is mostly due to this mass accretion process in the disk, or that L328-IRS has mostly accumulated the mass through this constant accretion rate during its outflow activity, its mass was estimated to be $\sim 0.012 - 0.023 M_{\odot}$, suggesting L328-IRS to be a substellar object.

However, we leave our identification of L328-IRS as a proto-brown dwarf to be tentative because of various uncertainties especially regarding the mass accretion rate.

[7 IM-03] Accretion Flow and Raman-scattered O VI and C II Features in the Symbiotic Nova RR Telescopii

Jeong-Eun Heo¹, Hee-Won Lee¹, Rodolfo Angeloni², Tali Palma³ and Francesco Di Mille⁴

¹*Sejong University, Korea,* ²*Universidad de La*

Serena, Chile, ³*Observatorio Astronómica, Argentina,* ⁴*Las Campanas Observatory, Chile*

RR Tel is an interacting binary system in which a hot white dwarf (WD) accretes matter from a Mira variable via gravitational capture of the stellar wind. We present a high-resolution optical spectrum of RR Tel obtained with MIKE at Magellan-Clay telescope, Chile. We find broad emission features at 6825, 7082, 7023, and 7053 Å, which are formed through Raman scattering of far-UV O VI $\lambda\lambda$ 1032 and 1038 Å, C II $\lambda\lambda$ 1036 and 1037 Å with atomic hydrogen. Raman O VI 6825 and 7082 features are characterized by double-peaked profiles indicative of an accretion flow with a characteristic speed $\sim 30 \text{ km/s}$, whereas the Raman C II features exhibit a single Gaussian profile with FWHM $\sim 10 \text{ Å}$. Monte Carlo simulations for Raman O VI and C II are performed by assuming that the emission nebula around the WD consists of the inner O VI disk with a representative scale of 1 AU and the outer part with C II sphere. The best fit for Raman profiles is obtained with an asymmetric matter distribution of the O VI disk, the mass loss rate of the cool companion $\dot{M} \sim 2 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$ and the wind terminal velocity $v \sim 10 \text{ km/s}$. We also find O VI doublet at 3811 and 3834 Å, which are blended with other emission lines. Our profile decomposition shows that the O VI $\lambda\lambda$ 3811, 3834 doublet have a single Gaussian profile with a width $\sim 25 \text{ km/s}$. A comparison of the restored fluxes of C II $\lambda\lambda$ 1036 and 1037 from Raman C II features with the observed C II λ 1335 leads to an estimate of a lower bound of $N(\text{CII}) > 9.87 \times 10^{13} \text{ cm}^{-2}$ toward RR Tel, which appears consistent with the presumed distance $D \sim 2.6 \text{ kpc}$.

[7 IM-04] Distance measurements for double red clump in the Milky Way bulge using Gaia DR2

Dongwook Lim, Seungsoo Hong, Young-Wook Lee
Center for Galaxy Evolution Research & Department of Astronomy, Yonsei University

The presence of double red clump (RC) in the Milky Way bulge is widely accepted as evidence for a giant X-shaped structure originated from the bar instability. We suggested, however, a drastically different interpretation based on the multiple stellar populations phenomenon as is observed in globular clusters. Our discovery of a significant difference in CN-band between two RCs strengthens our scenario. On the other hand, recent Gaia survey provides trigonometric parallax