

the H₂ emission lines for six SNRs are well consistent ($\pm 5 \text{ km s}^{-1}$) with those obtained in previous radio observations, while for the other five SNRs (G18.1-0.1, G18.9-1.1, Kes 69, 3C 396, W49B), they are significantly different. We discuss the velocity discrepancies in these five SNRs. In G9.9-0.8, the H₂ emission shows non-thermal line ratios and very narrow line width ($\sim 4 \text{ km s}^{-1}$), and we discuss its origin.

태양/태양계

[구 SS-01] Observation of the Rebound Shock Waves and the EUV Brightening of a Light Bridge Jet

Heesu Yang

Korea Astronomy and Space Science Institute

H α jets of cool chromospheric plasma are protruding into the solar corona 10-100 Mm above the photosphere. The driving mechanisms of H α jets have been widely studied for decades. However, the detailed process is still elusive. We observed shock signatures moving along a dark jet using 1.6 meter Goode Solar Telescope at Big Bear Solar Observatory. The first shock front of the jet shows sharp --- when it moves upward, while fuzzy and granulated when it moves downward. The jet itself extends upward when the second shock front of the jet reaches the top of the jet. We find abrupt EUV brightenings when the second shock front collides with the edge of the jet. The third front and the fourth front quasi-periodically. These phenomena might be the signs of the rebound shock waves triggered by p-mode wave leakages at the bottom of the jets. Our observation suggests that the jet can be triggered by the rebound shock waves generated by the p-mode waves leaked at the bottom of the jets.

[구 SS-02] Development of a diagnostic coronagraph on the ISS: progress report

Yeon-Han Kim¹, Seonghwan Choi¹, Su-Chan Bong¹, Kyungsuk Cho^{1,2}, Young-Deuk Park¹, Jeffrey Newmark³, Nat. Gopalswamy³, Seiji Yashiro³, Nelson Reginald³

¹*Korea Astronomy and Space Science Institute, Korea*

²*University of Science and Technology, Korea*

³*NASA Goddard Space Flight Center, USA*

The Korea Astronomy and Space Science

Institute (KASI) has been collaborating with the NASA Goddard Space Flight Center (GSFC), to install a diagnostic coronagraph on the International Space Station (ISS). The coronagraph is designed to obtain simultaneous measurements of electron density, temperature, and velocity using multiple filters in the 3-10 Rs range. In 2019, we developed a new coronagraph and launched it on a stratospheric balloon (BITSE) from Fort Sumner, New Mexico in USA. As the next step, the coronagraph will be further developed, installed and operated on the ISS (CODEX) in 2023 to understand the physical conditions in the solar wind acceleration region, and enable and validate the next generation space weather models. In this presentation, we will report recent progress and introduce future plan.

[구 SS-03] Investigation of sunspot substructure using chromospheric bright patches in a merging sunspot

Kyuhyoun Cho

Seoul National University

Sunspot substructure is an important subject to explain their stability and energy transport. Previous studies suggested two substructure models, monolithic and spaghetti model, but no clear evidence has been found supporting a particular model. To obtain the clue of the sunspot substructure the IRIS Mg II 2796Å slit-jaw images (SJI) were examined. The Mg II images formed in the chromosphere show bright patches inside umbrae which are regarded as an observational signature of upward propagating slow magnetohydrodynamic (MHD) waves. The slow MHD waves are expected to be generated by convective motion below the photosphere. By tracking the motion of the bright patches it is possible to estimate the locations of oscillation centers that correspond to the occurrence position of the convections. I investigated the spatial distribution of the oscillation center in a merging sunspot and found it is randomly distributed. It implies that the occurrence rate of the convective motion inside the sunspot is not much different from that of between the two sunspots, and supports the spaghetti model as the sunspot substructure.

[구 SS-04] Inference of Chromospheric Plasma Parameters on the Sun from Strong Absorption Lines

Jongchul Chae, Maria S. Madjarska, Hannah Kwak, Kyuhyoun Cho

Department of Physics and Astronomy, Seoul National University (서울대)

The solar chromosphere can be observed well through strong absorption lines. We infer the physical parameters of chromospheric plasmas from these lines using a multilayer spectral inversion. This is a new technique of spectral inversion. We assume that the atmosphere consists of a finite number of layers. In each layer the absorption profile is constant and the source function is allowed to vary with optical depth. Specifically, we consider a three-layer model of radiative transfer where the lowest layer is identified with the photosphere and the two upper layers are identified with the chromosphere. This three-layer model is fully specified by 13 parameters. Four parameters can be fixed to prescribed values, and one parameter can be determined from the analysis of a satellite photospheric line. The remaining eight parameters are determined from a constrained least-squares fitting. We applied the multilayer spectral inversion to the spectral data of the H α and the Ca II 854.21 nm lines taken in a quiet region by the Fast Imaging Solar Spectrograph (FISS) of the Goode Solar Telescope (GST). We find that our model successfully fits most of the observed profiles and produces regular maps of the model parameters. We conclude that our multilayer inversion is useful to infer chromospheric plasma parameters on the Sun.

[7 SS-05] Detection of Opposite Magnetic Polarity in a Light Bridge : Its Emergence and Cancellation in association with LB Fan-shaped Jets

Eun-Kyung Lim¹, Heesu Yang¹, Vasyl Yurchyshyn², Jongchul Chae³, Donguk Song⁴, Maria S. Madjarska⁵

¹KASI, ²BBSO, ³SNU, ⁴NAOJ, ⁵MPIS

Light bridges (LBs) are relatively bright structures that divide sunspot umbrae into two or more parts. Chromospheric LBs are known to be associated with various activities including fan-shaped jet-like ejections and brightenings. Although magnetic reconnection is frequently suggested to be responsible for such activities, not many studies presented firm evidence to support the scenario. We carry out magnetic field measurements and imaging spectroscopy of a LB where fan-shaped jet-like ejections occur with co-spatial brightenings at their footpoints. We study their fine photospheric structures and magnetic field changes using TiO images, Near-Infrared Imaging Spectropolarimeter data, and H α data taken by the 1.6 m Goode Solar Telescope. As a result, we detect magnetic flux

emergence in the LB that is of opposite polarity to that of the sunspot. The new flux cancels with the pre-existing flux at a rate of 5.6×10^{18} Mx hr⁻¹. Both recurrent jet-like ejections and their footpoint brightenings are initiated at the vicinity of the magnetic cancellation, and show apparent horizontal extension along the LB at a projected speed of 4.3 km s⁻¹ to form the fan-shaped appearance. Based on these observations, we suggest that the fan-shaped ejections may have resulted due to slipping reconnection between the new flux emerging in the LB and the ambient sunspot field.

[7 SS-06] Surface exposure age of (25143) Itokawa estimated from the number of mottles on the boulder

Sunho Jin, Masateru Ishiguro
Seoul National University

Various processes, such as space weathering and granular convection, are occurring on asteroids' surfaces. Estimation of the surface exposure timescale is essential for understanding these processes. The Hayabusa mission target asteroid, (25143) Itokawa (Sq-type) is the only asteroid whose age is estimated from remote sensing observations as well as sample analyses in laboratories. There is, however, an unignorable discrepancy between the timescale derived from these different techniques. The ages estimated based on the solar flare track density and the weathered rim thickness of regolith samples range between 102 and 104 years [1][2]. On the contrary, the ages estimated from the crater size distributions and the spectra cover from 106 to 107 years [3][4].

It is important to notice that there is a common drawback of both age estimation methods. Since the evidence of regolith migration is found on the surface of Itokawa [5], the surficial particles would be rejuvenated by granular convection. At the same time, it is expected that the erasure of craters by regolith migration would affect the crater size distribution.

We propose a new technique to estimate surface exposure age, focusing on the bright mottles on the large boulders. Our technique is less prone to the granular convection. These mottles are expected to be formed by impacts of mm to cm-sized interplanetary particles. Together with the well-known flux model of interplanetary dust particles (e.g., Grün, 1985 [6]), we have investigated the timescale to form such mottles before they become dark materials again by the space weathering. In this work, we used three AMICA (Asteroid Multi-band Imaging Camera) v-band