

麻醉劑의 心臟藥理學的 研究

第3報 Fructose 가 Halothane 低下遊離心房의 收縮反應에 미치는 效果

慶熙大學校 醫科大學 藥理學教室

高 啓 昌 · 鄭 址 昌 · 韓 大 燮

Cardiac Pharmacology of Anesthetics

3. Contractile Response of Halothane-Depressed Isolated Atria to Fructose

Kye Chang Ko, Jee Chang Jung and Dae Sup Han

*Department of Pharmacology Kyung Hee University School of Medicine
Seoul, Korea*

INTRODUCTION

This investigation is a continuation of studies dealing with the mechanism of the depressant action of inhalation anesthetics on cardiac function¹⁻³. It has been reported that anoxia and halothane produce similar decreases in contractility and potassium content in the perfused rat heart. That anoxia produced an increase in coronary flow rate not seen with halothane and produced irreversible damage to the contractile mechanism, again not seen with halothane, suggested that different biochemical changes were occurring with the two variable⁵.

Recently, we found that pyruvate partially restored the contractility of rat atria depressed 50% with approximately 6 mg/100 ml halothane, and that lactate and acetate also partially restored the halothane-depressed atria, despite the fact that additional glucose had no significant effect on the depressed contractility¹⁻³. From these findings we concluded that at

least part of the negative inotropic action of halothane is the result of inhibition of glucose uptake or utilization in the glycolytic pathway of the heart. The site of blockade by halothane must precede the conversion of pyruvate to acetyl CoA.

The present studies represent an attempt to localize further the site of halothane action in the glycolytic sequence by using the metabolizable substrate, fructose. We found a few references with respect to the utilization of fructose by the heart, and those indicated that fructose was a poorly-utilized substrate compared with glucose^{6,7}. Therefore, we first demonstrated dose-response curves in the substrate-depleted heart to determine if, and at what concentrations, fructose could serve as a source of fuel for the contractile process. Next, we attempted to determine whether fructose was metabolized via the phosphofructokinase step and, finally, we observed the effect of fructose on the halothane-depressed atria. From the results we conclude: 1) fructose can serve as a source of fuel for the contraction of isolated

rat atria: 2) metabolism of fructose occurs via the phosphofructokinase step: 3) fructose partially restores the contractility of atria depressed by halothane. Thus, the site of halothane blockade must be either the uptake of glucose or its utilization prior to the phosphofructokinase step.

METHODS

Male Sprague-Dawley rats weighing 180 to 200 g. having ad lib. access to food and water, were employed. Atria were removed from decapitated rats suspended in a modified Krebs-Ringer bicarbonate glucose medium of the following composition¹⁻³⁾ (mM): NaCl, 120; KCl, 4.8; CaCl₂, 1.22; MgSO₄·7H₂O, 1.33; KH₂PO₄, 1.2; NaHCO₃, 25.3; glucose, 5.55. The medium was gassed with 95% O₂-5%CO₂ at pH 7.4 and 30°C.

A constant resting tension of 750 mg was maintained throughout the experiment. The developed tension was recorded with a Statham strain gauge, and the atria were electrically stimulated at a rate of 200 pulses/min. An equilibration period of 60 min in the above medium was allowed before readings were taken. The experimental values of contractility (peak tension) were compared with those of the control records obtained at zero time (following equilibration) and expressed as per cent change in developed tension. Halothane was administered to the medium by means of the anesthetistat previously described by Paradise and Griffith^{4,5)}. Halothane concentration in the medium was determined at 10 to 30 min intervals with a gas chromatograph throughout the experimental period⁵⁾.

In some experiments (fig. 1) the medium was changed to substrate-free following the

one-hour equilibration period.

In the experiment with bicarbonate-free medium (fig. 2), the procedures were conducted by means of techniques previously described by Ko et al.⁹⁾

RESULTS

Effect of Fructose on Substrate-Depleted Atria

The effect of fructose on the functional properties of atria had to be established to study its action on halothane-depressed atria. Experiments were designed using substrate-depleted medium to determine the contractile behavior of atria in the absence of exogenous substrate,




Fig. 1. Effect of fructose on substrate-depleted atria.

In this and subsequent figures zero time is that time following a 60-minute equilibration of the atria in the normal Krebs-Ringer glucose medium. By substrate-depleted atria is meant exposure to substrate-free medium, but otherwise, normal medium. Fructose was added 30 minutes after exposure to substrate-free medium. Vertical bars indicate \pm one standard error of the mean. Each curve represents six experiments.

to provide control data with which the response to fructose might be compared. Results are summarized in figure 1.

Developed tension of the atria progressively decreased in the substrate-free medium after the one-hour equilibration period with Krebs Ringer bicarbonate glucose medium.

After 30 min in the substrate-free medium, fructose was added at a concentration of 10, 20, 30 or 40 mM (fig. 1). The addition of fructose resulted in marked recovery of the force of contraction: the maximally effective concentration of the fructose was 30 mM. The same concentration of the nonmetabolized sugar, sucrose, however, had no effect on the substrate-depleted atria, indicating that the action of fructose at this high concentration is a result of its metabolism.

Effects of Fructose, Glucose and Pyruvate on Atria depressed by Bicarbonate-Free Medium

Having established the availability of fructose as an energy source for contraction, it was important to determine the pathway by which fructose is utilized. Shaw and Stadie^{10,11} demonstrated the dependence of the phosphofructokinase reaction in the isolated rat diaphragm on the presence of bicarbonate in the medium. In the absence of bicarbonate neither radioactive glucose nor fructose in the bathing medium could be incorporated into radioactive fructose diphosphate, although earlier products of the metabolism of glucose were found. In the presence of bicarbonate both glucose and fructose could be converted to fructose diphosphate. If bicarbonate were also necessary for the activity




Fig. 2. Effect of substrates on atria in bicarbonate-free medium.

of phosphofructokinase in rat atria, and if fructose were metabolized via this enzyme, we would expect: 1) a fall in contractility of atria incubated in bicarbonate-free glucose medium and 2) resoration of contractility with pyruvate (a substrate not metabolized via phosphofructokinase) but not with added glucose or fructose. Figure 2 demonstrated these findings exactly. Thus, it appears that fructose is metabolized via the phosphofructokinase enzyme.

Effect of Fructose on Halothane-Depressed Atria

Addition of 30 mM fructose 30 minutes after start of administration of halothane resulted in a prompt and sustained increase in force of contraction despite the continued administration of halothane (fig. 3). This effect was

similar to that seen with pyruvate, acetate and lactate but not glucose, on halothane-depressed atria¹⁻³). This antagonism of halothane depression by fructose but not glucose, along with the evidence pointing to the utilization of fructose via the phosphofructokinase step, suggests that the mechanism of the negative inotropic action of halothane in these atria is blockade of the uptake or utilization of glucose prior to the phosphofructokinase step.

Effect of Fructose on Normal Atria

Addition of 30 mM fructose to atria bathed in the normal Krebs-Ringer bicarbonate glucose medium resulted in no demonstrable change in cardiac contractility. These results confirm similar observations by Gimeno et al.⁷) and emphasize the importance of halothane for the positive inotropic effect of fructose.




Fig. 3. Effect of fructose on halothane-depressed atria.

DISCUSSION

Fructose, when used in high concentration (30 mM) has been shown to serve as an excellent substrate for the maintenance of contractility by the isolated rat atria. Opie et al.⁶ found fructose (5 mM) to be taken up and metabolized to CO₂ less than 1/5 as rapidly as glucose (5 mM). Gimeno et al.⁷ demonstrated that fructose, at concentrations of 5.5 or 11 mM, is utilized for contractility but not nearly as efficiently as the corresponding concentration of glucose. Thus, the uptake of fructose or its conversion to fructose-6-phosphate may be rate-limiting, higher concentration of fructose than glucose being necessary for similar effects.

The lack of a positive inotropic effect of fructose in bicarbonate free medium is an agreement with the data of Shaw and Stadie, obtained in a study of the rat diaphragm^{10,11}. They showed a failure of conversion of labelled fructose to fructose diphosphate in bicarbonate-free medium, along with other data indicating the importance of bicarbonate for progression of the phosphofructokinase reaction. Thus, fructose apparently is utilized via phosphofructokinase in the diaphragm.

A number of studies in the rat heart are consistent with a lack of phosphofructokinase activity in bicarbonate-free medium. In this medium glucose is relatively ineffective in maintaining the contractile activity of rat ventricle strips¹²⁻¹⁴. Rice and Berman^{15,16} demonstrated that the oxidation of glucose by heart strips incubated in bicarbonate-free medium is lower than the oxidation of pyruvate or acetate. In contrast, they have observed that in medium containing bicarbonate glucose maintains contractile activity¹³, and Hood and Saunders have reported that glucose is rapidly

oxidized in this medium¹⁷.

The positive inotropic effect of fructose on halothane-depressed, but not on normal, atria, along with the previous report¹³ showing a lack of positive inotropic effect of additional glucose in halothane-depressed atria, suggest that the negative inotropic effect of halothane is at least partly the result of an interference with glucose uptake or metabolism prior to the phosphofructokinase step. Figure 4 is a schematic representation of the glycolytic pathway.




Fig. 4. Fructose and halothane effect on the heart. Schematic representation of glycolysis, showing points at which various substrates enter the scene. A, B and C represent possible sites of halothane blockade. Lactate, acetate, pyruvate, but not glucose, have been shown to overcome halothane-induced cardiac depression (1-3). In this report fructose also was shown to overcome depression by halothane but not by bicarbonate-free medium which inhibits phosphofructokinase step.

The possible sites of halothane blockade are A) uptake of glucose into the heart, B) conversion of glucose-6-phosphate by the enzyme hexokinase, and C) conversion of glucose-6-phosphate to fructose-6-phosphate by phosphohexose isomerase. Site B is not very likely since, in addition to glucose, hexokinase catalyzes the conversion of fructose to fructose-6-phosphate, a reaction seemingly unimpeded since fructose is apparently well utilized by the halothane-depressed atria. Site A has interesting implications since, at least in the erythrocyte, fructose and glucose appear to be taken up by different mechanisms: at least, the kinetics for uptake are different¹⁸⁾. It would be of interest to study the uptake of the nonmetabolizable sugar, 3-O-methyl glucose, in the presence of halothane. This sugar taken into the heart by the same mechanism as glucose, with which it competes for uptake. This will be the subject of future investigations.

A report by Hech et al.¹⁹⁾ indicates that halothane (5 vol per cent) is without effect on anaerobic glycolysis of rat brain. These studies, however, were done on homogenates, where uptake is not a factor. If glycolysis of homogenized brain and intact atria are similar, this would seem a further indication that uptake is a likely mechanism of halothane blockade.

REFERENCES

- 1) Ko, K. C. and Paradise R.R.: *The effect of Substrates on Contractility of Rat Atria Depressed with Halothane*, *Anesthesiology* 31: 532, 1969.
- 2) 高啓昌：麻醉劑의 心臟藥理學的 研究. 第1報 全身麻醉劑 Halothane 의 心臟代謝抑制作用에 關한 基礎的 考察, 大韓藥理學雜誌 10(1):21, 1974.
- 3) 高啓昌·鄭址昌·韓大燮：麻醉劑의 心臟藥理學的 研究 第2報. 各種代謝基質에 對한 Halothane 低下遊離心房의 收縮反應. 大韓藥理學雜誌 10(1): 55, 1974.
- 4) Paradise, R.R., and Griffith, L.K.: *Influence of halothane, chloroform and methoxyflurane on potassium content of rat atria*, *Anesthesiology* 26:195, 1965.
- 5) Paradise, R.R., and Griffith, L.K.: *Electrolyte content of perfused rat ventricles exposed to halothane or anoxia*, *J. Pharmacol. Exp. Ther.*, 154:181, 1966.
- 6) Opie, L.H., Shipp, J.D., Evans, J.R., and Leoboef, B.: *Metabolism of glucose-U-C¹⁴ in perfused rat heart*, *Amer. J. Physiol.*, 203:839, 1962.
- 7) Gimeno, A.L., Lacuara, J.L., Gimeno, M.F., and Savino, E.A.: *Effects of monosaccharides, acetate, butyrate, lactate and pyruvate on the developed tension of isolated rat atria*, *Proc. Soc. Biol. Med.*, 130:1033, 1969.
- 8) Paradise, R.R., and Griffith, L.K.: *Control of concentration of volatile agents in open in vitro systems*, *Anesthesiology* 27:687, 1966.
- 9) Ko, K.C., Gimeno, A.L., and Berman, D.A.: *Effects of buffers on developed tension, membrane potentials, and ATP levels of atria*, *Amer. J. Physiol.*, 216:853, 1969.
- 10) Shaw, W.N., and Stadie, W.C.: *Coexistence of insulin-responsive and insulin-non-responsive glycolytic systems in rat diaphragm*, *J. Biol. Chem.*, 227:115, 1957.
- 11) Shaw, W.N., and Stadie, W.C.: *Two identical Embden-Myerhof enzyme systems in normal rat diaphragm differing in cytological location and response to insulin*, *J. Biol. Chem.*, 234:2491, 1959.
- 12) Covin, J.M., and Berman, D.A.: *Metabolic aspects of the positive inotropic action of fluoride on rat ventricle*, *J. Pharmacol. Exp. Ther.*, 125:137, 1959.
- 13) Berman, D.A., and Saunders, P.R.: *Energy sources for contraction of the rat ventricle in phosphate media*, *Circ. Res.*, 3:559, 1955.
- 14) Covin, J.M., and Berman, D.A.: *Studies on the mechanism of the positive inotropic action of malonate*, *J. Pharmacol. Exp. Ther.*, 117:443,

- 1956.
- 15) Rice, L.I., and Berman, D.A.: *Oxidation glucose-1-C¹⁴, glucose-6-C¹⁴ and pyruvate-2-C¹⁴ by contracting rat ventricle strips in the presence and absence of arsenate*, *J. Pharmacol. Exp. Ther.*, 127:11, 1959.
- 16) Rice, L.I., and Berman, D.A.: *Malonate and fluoride effects on metabolism and contraction of electrically stimulated heart strips*, *Amer. J. Physiol.*, 200:727, 1961.
- 17) Hood, J.E., and Saunders, P.R.: *Oxidation of carbon¹⁴-labeled glucose, pyruvate and succinate by isolated contracting myocardium*, *Amer. J. Physiol.*, 190:525, 1956.
- 18) Wilbrandt, W., and Rosenberg, T.: *The concept of carrier transport and its derivatives in pharmacology*, *Pharmacol. Rev.* 13:109, 1961.
- 19) Hoech, G.P., Jr., Matteo, R.S., and Fink, B.R.: *Effect of halothane on oxygen consumption of rat brain, liver and heart and anaerobic glycolysis of rat brain*, *Anesthesiology* 27:770, 1966.