


an orientation does not only provide for a closest approach between the allylic hydrogen and the oxygen atom which is not involved in C-O bonding, but also for a maximum overlap of the p -orbitals of the developing new π_{c-c} bond.

With regard to VCP-DPh, the conformation of 'axially positioned' allylic hydrogen atom is hindered by the two phenyl groups attached to C_1 atom of cyclopropane ring and the π -electron density of the vinyl group is not enough to produce photoene-product because of the absence of electron-donating groups attached to the vinyl group.




In most cases, 1,2-cycloaddition of singlet oxygen to olefins requires the presence of electron-donating groups in the alkene, such as $-NR_2$, $-OR$, and $-SR$. photooxygenation of alkyl-substituted olefins generally results in the formation of allylic hydroperoxides. However, if the ene reaction is precluded by the substitution pattern or for steric reasons, 1,2-dioxetanes or their cleavage products may be formed.

The enhanced reactivity of π -electron-rich olefins in 1,2-cycloaddition with singlet oxygen is in accord with the formulation of this species as an electrophilic reagent—a conclusion in harmony with its electronic structure. Bond formation thus requires transfer of electron density from the highest occupied molecular orbital (HOMO) of the olefin to the lowest unoccupied molecular orbital (LUMO) of singlet oxygen.

In the case of VCP-DPh, if the vinyl group is conjugated

with the two phenyl groups *via* π -character of the cyclopropane ring, the vinyl group is activated and undergoes 1,2-cycloaddition with singlet oxygen to yield 1,2-dioxetane. However, two phenyl groups attached to cyclopropane ring are not conjugated with the vinyl group and cyclopropane ring does not transmit the conjugation. Therefore, the vinyl group of VCP-DPh is not π -electron-rich and does not undergo 1,2-cycloaddition with singlet oxygen to produce 1,2-dioxetanes.

Cosensitized photooxygenation of VCP-DPh ($1 \times 10^{-2} M$) in oxygen-saturated acetonitrile solution with biphenyl ($1 \times 10^{-2} M$) and 9,10-dicyanoanthracene ($1 \times 10^{-4} M$) yielded 3,3-diphenyl-5-vinyl-1,2-dioxolane (64.8 % yield), benzophenone (35.2 % yield), and 1,3-butadiene.⁷



References

- (1) M. Lukina, *Russ. Chem. Rev.*, **31**, 419 (1962).
- (2) J. Pete, *Bull. Soc. Chim. France*, **357** (1967).
- (3) H. M. Frey and R. Walsh, *Chem. Rev.*, **69**, 103 (1969).
- (4) M. Charton in "The Chemistry of Alkenes," J. Zabicky Ed., Interscience Publishers, London, England, Vol. II, Chapter 10, 1970.
- (5) J. M. Stewart and G. K. Pagenkopf, *J. Org. Chem.*, **34**, 7 (1969).
- (6) S. C. Shim and S. W. Suh, *J. Korean Chem. Soc.*, **23**, 115 (1979).
- (7) The reaction was carried out at 10°C under oxygen with a Hanovia 450 W medium-pressure mercury arc lamp by using a $CuSO_4$ -filter solution. 1,3-Butadiene was detected in brominated form, 1,2,3,4-tetrabromobutane.

Novel Dechlorination of Arylsulfonyl Chlorides with Superoxide Anion (O_2^-)


Bong Chul Chung and Yong Hae Kim¹

Department of Chemistry, Korea Advanced Institute of Science and Technology, P.O. Box 150 Chongyangni Seoul 131, Korea (Received October 6, 1984)

The discovery of superoxide dismutase (SOD) by Fridovich and McCord in 1969¹ has considerably stimulated the investigation of superoxide anion radical (O_2^-), not only in biochemistry but also in organic chemistry recently. Although commercially available potassium superoxide (KO_2), sodium superoxide (NaO_2), and ammonium superoxide (NH_4O_2), as sources of superoxide had been prepared, both its instability in protic solvents and its limited solubility in organic solvents such as dimethylsulfoxide (DMSO) and acetonitrile had prevented studies of the chemistry of superoxide ion. How-

ever, the observation of Valentine and Curtis in which KO_2 can be appreciably dissolved in aprotic solvents by complexation with crown ethers² has quickly promoted the use of this reagent for many reactions with simple organic substrates. Numerous reports dealing with reactions of various organic substrates with "naked" superoxide anion have appeared within only past five years. However, only a few sporadic works on the reaction of organic sulfur compounds with O_2^- appeared until oxidations of organic sulfur compounds with O_2^- in the presence of 18-crown-6-ether were initiated.³⁻⁶

The substrates of organic sulfur compounds are either those which have sulfur-sulfur linkage such as disulfide, thioisulfonic-S-ester, thioisulfonic-S-ester, or thiolate and sulfinate, all of which are expected to give peroxy-sulfur compounds such as peroxy-sulfenate (1), -sulfinate (2), and -sulfonate (3) as initial intermediates in the reactions with O_2^- .




Earlier, Berger⁷ postulated the first intermediary peroxy-sulfenate (1) and peroxy-sulfinate (2) in the autoxidation of thiols to the sulfinic and sulfonic acids in strong alkaline media without any evidence for the formation of peroxy-sulfur intermediates. Though the two reactions are presumed to involve common intermediates of peroxy-sulfur species, these peroxy-sulfur intermediates which may have potential oxidizing abilities have neither been isolated nor detected.

In our previous communication⁸ on the epoxidations of olefins with a peroxy-sulfur intermediate generated from superoxide anion (O_2^-) and nitrobenzenesulfonyl chlorides, the formation of the peroxy-sulfonate intermediate was postulated to be 3, which is stable enough at low temperature. During the study of the peroxy-sulfenate intermediate, we have found that arylsulfenyl chlorides reacted with O_2^- in tetrahydrofuran or acetonitrile at low temperature to give the corresponding diaryldisulfides or arylsulfonates depend upon the substrates used. All the reactions were carried out without using crown ether.¹¹ The products were identified by the comparison of their IR, ¹H NMR spectra, and mp. with those of the authentic samples. The results are summarized in Table 1.

In a typical experiment, KO_2 (74 mg, 1.0 mmol) in dry THF (1.0 ml) was crushed with a glass rod. 2,4-Dinitrobenzenesulfonyl chloride (117 mg, 0.5 mmol) was dissolved in dry THF (2.0 ml). A solution of sulfenyl chloride was added into a heterogeneous solution of KO_2 , and then stirred vigorously at ca. 0°C under argon atmosphere. After 1.0 h, the reaction mixture was centrifuged for three times to separate the clean solution, which was concentrated to recover the starting material. Solid part was washed with H_2O and ethyl ether three times respectively and dried in vacuo to give di-*p*-nitrobenzene disulfide (99.5 mg, 85 %, mp, 278–280°C, decomp., ref.⁹ mp. 280°C, dec.).

2,4-Dinitrobenzenesulfonyl chloride, which has two strong electron withdrawing nitro groups, reacted with O_2^- to form the corresponding disulfide (85 %) as the main product together with trace amount of the corresponding sulfonate. While, less electron withdrawing benzenesulfonyl chloride yielded the benzenesulfonate potassium salt (91 %) as the main products.

The formation of peroxy-sulfenate (1) is indicated by the epoxidation of chalcone to the epoxide (10 %) when the former is used as a trapping agent for the activated oxygen of 1 which is expected to have an oxidizing ability.¹²



Scheme 1

TABLE 1: Reaction of Arylsulfenyl Chlorides with KO_2

Run	4, Ar	Solvent	$KO_2/4^b$	Reactn. Time(h)	Yield(%) ^c	
					5	6
1	$C_6H_5^{9a}$	THF	2	6.0	91	trace
2	$C_6H_5^{9a}$	THF	1.3	1.0	90	—c
3	<i>p</i> -Me- $C_6H_4^{9b}$	THF	2	6.0	85	10
4	<i>p</i> -Br- $C_6H_4^{9c}$	THF	2	6.5	71	21
5	β -Naphthyl ^{9c}	THF	2	5.5	50	45
6	<i>o</i> -NO ₂ - $C_6H_4^d$	THF	2	1.5	29	62
7	<i>o,p</i> -di-NO ₂ - $C_6H_3^d$	THF	2	1.0	trace	85
8	<i>o,p</i> -di-NO ₂ - $C_6H_3^d$	CH ₃ CN	2	1.5	—c	84

^a isolated yields, ^b molar ratio; KO_2 : substrate, ^c not checked
^d purchased from Aldrich Chemical Co.

The reactions appear to be initiated *via* a nucleophilic attack of O_2^- on the sulfenyl sulfur atom to form a peroxy-sulfenate radical, which may be readily converted to 1 by one electron transfer from O_2^- . The intermediate 1 may be further oxidized to 5. In the case of dinitrobenzene peroxy-sulfenate, the thiyl radical formed by the cleavage of sulfur-oxygen bonds of 1 are stabilized by the two nitro groups and readily couple to 6.

There have been a few reductive synthesis of disulfides from aryl sulfenyl chlorides using carbonyl complexes of metals¹³ and potassium iodide¹⁴ in protic solvents. The methods for the preparation of arylsulfonic or arylsulfinic acids by the oxidations of thiols^{14–19} or sulfenyl²⁰ and sulfonyl chlorides²¹ need strong oxidation conditions such as boiling nitric acid¹⁵, potassium permanganate¹⁶, Caro's acid¹⁷, alkaline autoxidation¹⁸ or basic hydrogen peroxide¹⁹. Our new reactions, however, can be carried out in aprotic solvents such as tetrahydrofuran and acetonitrile at lower temperature than 0°C.

Acknowledgment. This work was supported by KAIST grant.

References

- (1) J. M. McCord and I. Fridovich, *J. Biol. Chem.*, **244**, 6049 (1969).
- (2) J. S. Valentine and A. B. Curtis, *J. Amer. Chem. Soc.*, **97**,

- 224 (1975).
- (3) E. W. Neumann, *J. Chim. Phys.*, **2**, 31 (1934).
- (4) (a) T. Takata, Y. H. Kim and S. Oae, *Tetrahedron Lett.*, 821 (1979); (b) S. Oae, T. Takata and Y. H. Kim, *Tetrahedron*, **37**, 37 (1981).
- (5) (a) A. LeBerre and Y. Berguer, *Bull. Soc. Chim. Fr.*, 2306 (1966); (b) J. San Fillippo, Jr., L. J. Romano, C.-I. Chern, and J. S. Valentine, *J. Org. Chem.*, **40**, 1678 (1975).
- (6) S. Oae and T. Takata, *Tetrahedron Lett.*, **21**, 3689 (1980).
- (7) H. Berger, *Recl. Trav. Chem. Pays-Bas*, **82**, 773 (1963).
- (8) Y. H. Kim and B. C. Chung, *J. Org. Chem.*, **48**, 1562 (1983).
- (9) K. Pilgrim, *Tetrahedron*, **23(7)**, 3007 (1967).
- (10) (a) D. N. Harpp, B. T. Friedlander and R. A. Smith, *Synthesis*, 181 (1979); (b) K. R. Brower and I. B. Douglass, *J. Amer. Chem. Soc.*, **73**, 5787 (1951); (c) M. B. Sparke, J. L. Cameron and N. Kharasch, *ibid.*, **75**, 4907 (1953).
- (11) Most of known reactions with superoxide ion require to use crown ether² to promote the solubility of KO₂. However, surprisingly, our reactions underwent smoothly without using crown ether within 6 h at 0°C.
- (12) Since an intermediate, **1** is unstable, the epoxidation was carried out at -20°C in the reaction system of o, p-dinitrobenzenesulfonyl chloride, potassium superoxide and chalcone.
- (13) E. Lindner and G. Vitzthum, *Angew. Chem., Int. Ed. Engl.*, **8**, 518 (1969).
- (14) N. Kharasch and M. H. Wald, *Anal. Chem.*, **27**, 996 (1955).
- (15) W. Pendergast, *J. C. S. Perkin* **1**, 2240 (1975).
- (16) T. H. Chao and W. B. Hardy, *Chem. and Ind. (London)*, **81**, (1965).
- (17) R. J. Kenedy and A. M. Stock, *J. Org. Chem.*, **25**, 1901 (1960).
- (18) T. J. Wallace and A. Schriesheim, *Tetrahedron Lett.*, 1131 (1963).
- (19) W. Knobloch and K. Reintelen, *Arch. Pharm.*, **291**, 180 (1958).
- (20) N. Kunieda, K. Sakai and S. Oae, *Bull. Chem. Soc. Japan*, **41**, 3015 (1968).
- (21) W. Mieg and F. Wieners, *U. S. Pat.*, 2,215,083 (1941) (*Chem. Abst.*, **35**, 756, 1941).