

## FIXED POINT THEOREMS ON COMPACT CONVEX SETS IN TOPOLOGICAL VECTOR SPACES, II

SEHIE PARK

In our previous work [7], we established fixed point theorems for multimaps  $F : K \rightarrow 2^E$ , where  $K$  is a nonempty compact convex subset of a topological vector space  $E$  having sufficiently many linear functionals (that is,  $E^*$  separates points of  $E$ ). On the other hand, in a recent work [5], C. - W. Ha obtained multimap versions of fixed point theorems of Ky Fan [3, Theorems 1 and 3], which are comparable to the results in [7].

In the present paper, we give improvements and generalizations of some known results, mainly in [7] and [5]. In fact, Theorems 1 and 2 are improved versions of [7, Theorems 3 and 4]. Theorem 3 is a strengthened form of [5, Theorem 3] and equivalent to Reich [11, Theorem 2]. Theorem 4 generalizes [10, Theorem 3.1] and [5, Theorem 4]. Finally, in Theorem 5, we state sufficient conditions in order that a self-multimap  $F : K \rightarrow 2^K$  have a fixed point. Consequently, as in [7], various generalizations of the Brouwer fixed point theorem are improved in this paper.

A t. v. s. stands for a Hausdorff topological vector space and an l. c. s. for a locally convex t. v. s. The notion of an upper semicontinuous (u. s. c.), a lower semicontinuous (l. s. c.), or an upper hemicontinuous (u. h. c.) multimap is standard, see [1], [7].

For a t. v. s.  $E$  and a  $K \subset E$ , the inward and outward sets of  $K$  at  $x \in K$ ,  $I_K(x)$  and  $O_K(x)$ , resp., are defined as follows:

$$\begin{aligned} I_K(x) &:= \{x + r(u - x) \in E : u \in K, r > 0\}, \\ O_K(x) &:= \{x - r(u - x) \in E : u \in K, r > 0\}. \end{aligned}$$

The closures of  $I_K(x)$  and  $O_K(x)$  are denoted by  $\bar{I}_K(x)$  and  $\bar{O}_K(x)$ , resp. In the sequel,  $W(x)$  denotes either  $\bar{I}_K(x)$  or  $\bar{O}_K(x)$ .

---

Received October 26, 1988.

Supported by a grant from the Korea Science and Engineering Foundation, 1987.

Let  $cc(E)$  denote the family of nonempty closed convex subsets of  $E$  and  $kc(E)$  the family of nonempty compact convex subsets.

For a t. v. s.  $E$ , let  $E^*$  denote its topological dual.

We begin with the following improved version of [7, Theorem 3].

**THEOREM 1.** *Let  $K$  be a nonempty compact convex subset of a t. v. s.  $E$  and  $F$  a continuous (i. e., u. s. c. and l. s. c.) multimap defined on  $K$  such that either*

- (A)  $E^*$  separates points of  $E$  and  $F : K \rightarrow kc(E)$ , or
- (B)  $E$  is locally convex and  $F : K \rightarrow cc(E)$ .

*Then either  $F$  has a fixed point, or there exist a point  $v \in K$  and a continuous seminorm  $p$  on  $E$  such that*

$$0 < p(v - Fv) \leq p(w - Fv) \text{ for all } w \in W(v),$$

*where  $p(w - Fv) = \inf\{p(w - z) : z \in Fv\}$ .*

*Proof.* Use standard separation theorems [12], and just follow the proof of [7, Theorem 3].

Note that Reich [11, Theorem 3] follows from Theorem 1(B). Now we improve [7, Theorem 4] with the same proof.

**THEOREM 2.** *Under the hypothesis of Theorem 1,  $F$  has a fixed point if*

- (1) *for each  $x \in Bd K \setminus Fx$ , there exists a number  $\lambda$  (real or complex, depending on whether the vector space  $E$  is real or complex) such that*

$$|\lambda| < 1 \text{ and } (\lambda x + (1 - \lambda)Fx) \cap W(x) \neq \phi.$$

The following improved version of Ha [5, Theorem 3] can be compared with Theorem 1(B).

**THEOREM 3.** *Let  $K$  be a nonempty compact convex subset of a l. c. s.  $E$ , and  $F : K \rightarrow kc(E)$  an u. s. c. multimap. Then either  $F$  has a fixed point, or there exist  $v \in K$ ,  $u_0 \in Fv$ , and a continuous seminorm  $p$  on  $E$  such that*

$$0 < p(v - u_0) \leq p(w - u_0) \text{ for all } w \in W(v).$$

*Proof.* Suppose that  $F$  has no fixed point. Then by [5, Theorem 3], there exist  $v \in K$ ,  $u_0 \in Fv$ , and a continuous seminorm  $p$  on  $E$  such that

$$0 < p(v - u_0) \leq p(w - u_0) \text{ for all } w \in K,$$

and hence, by the method in the proof of [7, Theorem 1], the inequality holds for all  $w \in \bar{I}_K(v)$ .

For the outward case, the map  $F' : K \rightarrow kc(E)$  defined by  $F'x = 2x - Fx$  for each  $x \in K$  is u. s. c. Therefore, by the above argument, there exist  $v \in K$ ,  $u_1 \in F'v$ , and a continuous seminorm  $p$  on  $E$  such that

$$0 < p(v - u_1) \leq p(w' - u_1) \text{ for all } w' \in I_K(v).$$

For  $w \in O_K(v)$ , let  $w' = 2v - w$  and  $u_1 = 2v - u_0$  where  $u_0 \in Fv$ . Then we have

$$0 < p(v - u_0) \leq p(w - u_0) \text{ for all } w \in O_K(v)$$

and hence, for all  $w \in \bar{O}_K(v)$ . This proves Theorem 3 for the case  $W(v) = \bar{O}_K(v)$ .

Note that Theorem 3 is equivalent to Reich [11, Theorem 2] which contains some known results in [3], [10], [2], and the Tychonoff fixed point theorem.

In certain circumstance, Theorem 3 is more useful than Theorem 1. Consider the following example due to Gwinner [4, p. 575]: Let  $K = [0, 1] \times \{0\} \subset \mathbf{R}^2 \cong E$  with the ordinary norm. Define  $F : K \rightarrow 2^E$  by

$$\begin{aligned} F(c, 0) &= \text{co}\{(1, 1), (1, 2)\} \text{ if } c \in [0, 1), \\ F(1, 0) &= \text{co}\{(0, 0), (1, 1), (1, 2)\}. \end{aligned}$$

Note that  $F$  is u. s. c., but not l. s. c., and Theorem 1 is not applicable. However, Theorem 3 holds for this example by choosing  $v = (1, 0)$  and  $u_0 = (1, 1)$ .

As a direct consequence of Theorem 3, we have the following.

**THEOREM 4.** *Let  $K$  be a nonempty compact convex subset of a l. c. s.  $E$ , and  $F : K \rightarrow kc(E)$  an u. s. c. multimap. Then  $F$  has a fixed point if*

- (2) *for each  $x \in \text{Bd } K \setminus Fx$  and  $u \in Fx$ , there exists a number  $\lambda$  (as in (1)) such that*

$$|\lambda| < 1 \text{ and } \lambda x + (1 - \lambda)u \in W(x).$$

*Proof.* Suppose that  $F$  has no fixed point. By Theorem 3, there exist  $v \in K$ ,  $u_0 \in Fv$ , and a continuous seminorm  $p$  on  $E$  such that

$$0 < p(v - u_0) \leq p(w - u_0) \text{ for all } w \in W(v).$$

Since  $v \notin Fv$ , if  $v \in \text{Int } K$ , then there exists a  $\lambda$  (say  $\lambda = 1/2$ ) such that  $|\lambda| < 1$  and  $\lambda v + (1 - \lambda)u_0 \in W(v) = E$ . Therefore, there always exist a  $\lambda$  such that  $|\lambda| < 1$  and  $w := \lambda v + (1 - \lambda)u_0 \in W(v)$  whether  $v \in \text{Int } K$

or  $v \in \text{Bd } K$ . Then we have

$$0 < p(v - u_0) \leq p(w - u_0) = |\lambda| p(v - u_0),$$

which contradicts  $|\lambda| < 1$ .

Note that (2)  $\Rightarrow$  (1).

For a real t. v. s.  $E$ , (2) is equivalent to  $Fx \subset W(x)$  for all  $x \in K$ , and hence there exist more general results than Theorem 4 (see [7, Theorems 6 and 7]). (Note here that the first part of the proof of [7, Theorem 6] showing the existence of  $h \in E^*$  such that  $(h, v) < 0$  for all  $v \in u - Fu$  is incorrectly stated. Instead just use the separation theorem for a t. v. s.  $E$  on which  $E^*$  separates points (e. g., [12, p. 70]). Furthermore, in a recent work [8], we noted that [7, Theorem 6] holds for an u. h. c. map instead of an u. s. c. map.)

However, for a complex l. c. s.  $E$ , Theorem 4 generalizes Reich [10, Theorem 3.1; 7, Theorem 5] and Ha [5, Theorem 4]. We note that these two results are the same. In fact, Reich adopted one of the following boundary conditions:

(3) for each  $x \in \text{Bd } K \setminus Fx$ ,

$$Fx \subset IF_K(x) = \{x + c(y - x) : y \in K, \text{Re}(c) > 1/2\}.$$

(3)' for each  $x \in \text{Bd } K \setminus Fx$ ,

$$Fx \subset OF_K(x) = \{x + c(y - x) : y \in K, \text{Re}(c) < -1/2\}.$$

On the other hand, Ha used the following in Theorem 4 instead of (2):

(4) for each  $x \in K$  and  $u \in Fx$ , there exists a number  $\lambda$  (as in (1)) such that

$$|\lambda| < 1 \text{ and } \lambda x + (1 - \lambda)u \in K.$$

Note that (4)  $\Rightarrow$  (2) since  $K \subset I_K(x)$  and that  $z \in IF_K(x)$  iff there is a number  $\lambda$  (as above) such that  $|\lambda| < 1$  and  $\lambda x + (1 - \lambda)z \in K$  [9]. Therefore, (3)  $\Leftrightarrow$  (4)  $\Rightarrow$  (2). Reich [9] noted that (3) can not be replaced by  $Fx \cap IF_K(x) \neq \emptyset$ . For a real t. v. s.  $E$ , (4) is equivalent to  $Fx \subset I_X(x)$  for all  $x \in K$ .

Finally, for a selfmap  $F : K \rightarrow 2^K$ , we have the following improved version of [6, Theorem 1].

**THEOREM 5.** *Let  $K$  be a nonempty compact convex subset of a t. v. s.  $E$  having sufficiently many linear functionals, and  $F : K \rightarrow cc(K)$ . Then  $F$  has a fixed point if one of the following holds:*

(i)  $F$  is continuous.

(ii)  $E$  is real and  $F$  is u. h. c.

(iii)  $E$  is locally convex and  $F$  is u. s. c.

*Proof.* (i) follows from Theorem 2, (ii) is a consequence of the new version of [7, Theorem 6] in [8], and (iii) follows from Theorem 4.

### References

1. J.-P. Aubin and I. Ekeland, *Applied Nonlinear Analysis*, John Wiley & Sons, New York, 1984.
2. F. E. Browder, *On a sharpened form of the Schauder fixed point theorem*, Proc. Nat. Acad. Sci. USA **74**(1977), 4749-4751.
3. Ky Fan, *Extensions of two fixed point theorems of F. E. Browder*, Math. Z. **112**(1969), 234-240.
4. J. Gwinner, *On fixed points and variational inequalities — a circular tour*, Nonlinear Analysis, TMA **5**(1981), 565-583.
5. C.-W. Ha, *On a minimax inequality of Ky Fan*, Proc. Amer. Math. Soc. **99**(1987), 680-682.
6. Sehie Park, *Common fixed points of maps on topological vector spaces having sufficiently many linear functionals*, J. Korean Math. Soc. **24**(1987), 59-63.
7. Sehie Park, *Fixed point theorems on compact convex sets in topological vector spaces*, Fixed Point Theory and Its Applications(Proc. of a Conf. held at the I. C. M., Aug. 4-6, 1986), Contemp. Math. **72**(1988), 183-191.
8. Sehie Park, *Fixed point and coincidence theorems for u. h. c. multifunctions on paracompact convex sets*, to appear.
9. S. Reich, *Fixed points in locally convex spaces*, Math. Z. **125**(1972), 17-31.
10. S. Reich, *Approximate selections, best approximations, fixed points, and invariant sets*, J. Math. Anal. Appl. **62**(1978), 104-113.
11. S. Reich, *Fixed point theorems for set-valued mappings*, J. Math. Anal. Appl. **69**(1979), 353-358.
12. W. Rudin, *Functional Analysis*, McGraw-Hill, New York, 1973.

Seoul National University

and

The Mathematical Sciences Research Institute of Korea

Seoul 151-742, Korea