

with the chemical shift of the residual protons of the solvent used as internal standard. ^{13}C NMR chemical shifts are reported in ppm by using the natural abundance of ^{13}C of the solvent as an internal standard. Mass spectra were provided by the University of Illinois mass spectrometry facility. YMC RP and Silica gel (230-400 mesh) were used for flash column chromatography. YMC ODS 30×10 mm column was used for hplc separation.

Collection and Isolation. The sponge was collected Oct., 1992 from Manado bay north along the coast of Sulawesi up to Bunaken Island in Indonesia. The sponge was identified by Dr. Peter Karuso, University of Macquarie. The freeze-dried sponge (50.5 g) was extracted for 2 days with 400 mL of CH_2Cl_2 -iPrOH (1:1), and the residue (3.02 g) was partitioned between MeOH/ H_2O (9:1) and *n*-hexane. The MeOH/ H_2O layer yielded 2.03 g of red solid. The MeOH/ H_2O layer was subjected to RP- C_{18} flash column chromatography and Si gel flash column chromatography sequentially using MeOH- H_2O (75:25) and MeOH- CH_2Cl_2 (8:92), yielding 40 mg of mixture of bastadins. The mixture was separated by C_{18} reversed-phase hplc [ODS column, MeOH/ H_2O /MeCN (1:1:1), 2 mL/min] to give 5 mg of **1** as a pure white solid and 19 mg of **2** as shown in scheme 1.

HMBC condition for **1**: concentration of sample, 5 mg/DMSO- d_6 , 0.6 mL, 256 of blocks \times 2048 data matrix with 64 of scan per t_1 increment, 3.0 delay period for long-range couplings, 100° shifted since bell squared filtering for t_1 and 5100 MHz line broadening for it.

Bastadin (1). Hrfabms m/z $[\text{M}+\text{H}]^+$ 956.8606 (calcd for $\text{C}_{34}\text{H}_{26}\text{N}_4\text{O}_{9.75}\text{Br}_2$, 956.8627); hreims m/z 417.8925 (calcd for $\text{C}_{16}\text{H}_8\text{N}_2\text{O}_2$, 417.8953); ir ν_{max} (Nujol) 3600-3100, 1660, 1640, 1490, 1470, 1285, 1220 cm^{-1} ; UV ν_{max} (MeOH) ($\log\epsilon$) 278 (4.1) nm; ^1H and ^{13}C nmr see Table 1.

Bastadin 3(2)/⁴. ^1H NMR (in MeOH- d_4): δ 7.36 (2H, d, $J=2.2$ Hz), 7.28 (2H, d, $J=2.2$ Hz), 7.12 (2H, d, $J=2.2$ Hz), 6.89 (2H, dd, $J=8.3$, 2.2 Hz), 6.72 (2H, d, $J=8.3$ Hz), 3.80 (4H, s), 3.37 (4H, t, $J=6.9$ Hz), 2.66 (4H, t, $J=7.3$ Hz); ^{13}C NMR, δ 165.91, 153.92, 153.74, 153.61, 134.16, 133.29, 133.01, 132.27, 130.31, 130.07, 128.80, 117.21, 114.09, 110.67, 42.09, 35.19, 28.83; Irradiation at δ 3.80 (H-1) induced nuclear Overhauser enhancement (nOe) at δ 7.28 and 7.12. Irradiation at δ 2.62 (H-6) induced nOe at δ 3.37, 7.36 and 6.89. Irradiation at δ 3.37 (H-5) induced nOe at δ 2.62, 7.36 and 6.89.

Acknowledgment. We thank Toshio Ichiba and Mark Hamman for collecting the sponge; Mike Severns and Loky Herlambang of the Nusantara Diving Center for logistic support; Dr. Peter Karuso for sponge taxonomy; Wesley Yoshida for nmr assistance; Drs. Kenneth Rinehart and Ryuichi Sakai, University of Illinois at Urbana-Champaign, for mass spectral data.

References

1. Pordesimo, E. O.; Schmitz, F. J. *J. Org. Chem.* **1990**, *55*, 4704.
2. Miao, S.; Andersen, R. J.; Allen, T. M. *J. Nat. Prod.* **1990**, *53*, 1441.
3. Kazlauskas, R.; Lidgard, R. O.; Murphy, P. T.; Wells, R. *J. Tetrahedron Lett.* **1980**, *21*, 2277.
4. Kazlauskas, R.; Lidgard, R. O.; Murphy, P. T.; Wells, R. J.; Blount, J. F. *Aust. J. Chem.* **1981**, *34*, 765.
5. Butler, M. S.; Lim, T. K.; Capon, R. J.; Hammond, L. S. *Aust. J. Chem.* **1991**, *44*, 287.
6. Carney, J. R.; Scheuer, P. J.; Kelly-Borges, M. *J. Nat. Prod.* **1993**, *56*, 153.
7. Dexter, A. F.; Garson, M. J.; Hemling, M. E. *J. Nat. Prod.* **1993**, *56*, 782.
8. Gulavita, N. K.; Wright, A. E.; McCarthy, P. J.; Pomponi, S. A.; Kelly-Borges, M.; Chin, M.; Sills, M. A. *J. Nat. Prod.* **1993**, *56*, 1613.

Structure of Bis(N,N-dimethyl-2-thiophenemethylammonium)Tetrachlorocobaltate(II)

Mi-A Kwon, Young Jin Kim, Nak Suk Cho,
and Sung-Nak Choi*

Department of Chemistry, Pusan National University,
Pusan 609-735, Korea

Received April 18, 1995

2-(dimethylaminomethyl)thiophene (DMAT) ligand usually provides an [NS] donor set to form a delocalized five membered chelate ring with metal(II) ions and the structural characteristics of (DMAT) CuX_2 and (DMAT) NiX_2 where $\text{X}^- = \text{Cl}^-$ or Br^- have been characterized and reported (1). However, when cobalt(II)chloride reacts with DMAT in similar condition in ethanol, it does not produce a 1:1 complex of type (DMAT) CoCl_2 . Instead, bis(N,N-dimethyl-2-thiophenemethylammonium)tetra-chlorocobaltate(II), (DMTMAH) $_2\text{CoCl}_4$ is formed. We isolated the single crystals of this cobalt (II) complex and conducted X-ray crystallographic studies on it.

Experimental

The (DMTMAH) $_2\text{CoCl}_4$ was prepared by the direct reaction of DMAT (0.3 g, 2.0 mmol) with dried $\text{CoCl}_2\cdot 6\text{H}_2\text{O}$ (0.4 g, 1.7 mmol) in ethanol (50 mL) and triethylorthoformate (10 mL). The mixture was stirred for two hours at warm (yield, 54%). Dark blue single crystals were obtained by slow recrystallization in acetonitrile. A flat crystal of size $0.80\times 0.60\times 0.20$ mm was used for intensity measurements on an Enraf-Nonius CAD-4 diffractometer using the ω - θ scan technique; lattice parameters from setting angles of 23 reflections in the range $8<\theta<10^\circ$; scan width $(0.8+0.350\tan\theta)$ and with variable scan rate 1 to 7°min^{-1} ; $2\theta_{\text{max}}=45.0^\circ$; $0<h<10$, $0<k<18$, $-14<l<14$. Intensity data for 3175 reflections were collected; 2858 unique observed reflections; 2621 included reflections with $F_o^2>1.5$ (F_o^2). Lorentz and polarization corrections were applied. The linear absorption coefficient is 14.6 cm^{-1} for Mo K α radiation. An empirical absorption correction based on a series of ψ -scans was applied to the data. Relative transmission coefficients ranged from 0.937 to 0.999. A secondary extinction correction was applied (2).

Table 1. Crystal Data, Data Collection, and Refinement of the Structure for (DTMAH)₂CoCl₄

chem formula	C ₁₄ H ₂₂ N ₂ S ₂ Cl ₄ Co
fw	483.22
F (000)	988
crystal dimensions (mm)	0.80×0.60×0.20
peak width at half-height (°)	0.80
Mo Kα radiation	λ=0.71073
temperature (°C)	23±1
monoclinic space group	P2 ₁ /c (# 14)
a=9.340(3) Å, b=17.667(2) Å, c=13.523(4) Å	
β, deg	101.00(1)
V (Å ³)	2190.4
Z	4
ρ _{calc} (g/cm ³)	1.47
μ (cm ⁻¹)	14.6
scan type	θ-ω
scan width, deg	0.8+0.350 tanθ
no. of refl. measured	3175 total, 2858 unique
reflections included	2621 with F _o ² >1.5σ (F _o ²)
R=(Σ F _o -F _c)/Σ F _o	0.037
Rw=(Σ F _o -F _c w ^{1/2})/Σ F _o w ^{1/2}	0.038

The final coefficient, refined in least-squares, was 0.0000007. The structure was solved by direct methods using 258 reflections (E>1.20). A total of one atom were located from an E-map prepared from the phase set with probability statistics: absolute figure of merit=1.19, residual=1.10, and psi zero=10.750. The remaining atoms were located in succeeding difference Fourier syntheses. Hydrogen atoms were located and added to the structure factor calculations but their positions were not refined. The structure was refined in full-matrix least-squares calculations where the function minimized was Σw(|F_o-|F_c|)² and the weight w is defined as 4F_o²/σ²(F_o²). Scattering factors were taken from Cromer and Waber (3). Anomalous dispersion effects were included in F_c (4); the values for Δf' and Δf'' were those of Cromer (5). The final cycle of refinement included 213 variable parameters; max.shift/e.s.d.=0.02. Convergence R=0.037, wR=0.038. The standard deviation of an observation of unit weight was 15.22. (Δρ)_{max}=0.65 e/Å³ with an estimated error based on ΔF (6) of 0.08. All calculations were performed on a VAX computer using MolEN (7).

Discussion

The single crystal structure of (DMTMAH)₂CoCl₄ was solved and the crystallographic data are summarized in Table 1. The atomic coordinates are listed in Table 2. The bond distances and angles are given in Table 3. The structure consists of two discrete DMTMAH⁺ cations and CoCl₄²⁻ anion. Two DMTMAH⁺ cations are surrounding the CoCl₄²⁻ anion from the opposite directions. Figure 1 illustrates the structure of the (DMTMAH)₂CoCl₄.

The cobalt(II) ions are 4-coordinated with chloride ions in normal tetrahedral geometry. The average Cl-Co-Cl angle is 108.5°. Commonly Co-Cl bond length of CoCl₄²⁻ in DMF

Table 2. Atomic coordinates and equivalent isotropic thermal parameters (Å²) with e.s.d.'s in parentheses

	U _{eq} =(U ₁₁ +U ₂₂ +U ₃₃)/3.			U _{eq}
	x	y	z	
Co	1.00049(4)	0.14455(2)	0.25620(2)	0.0405(1)
Cl(1)	0.89937(7)	0.11595(4)	0.39484(5)	0.0512(3)
Cl(2)	0.86608(7)	0.23704(4)	0.16020(4)	0.0487(3)
Cl(3)	1.22428(8)	0.19497(5)	0.31206(5)	0.0598(4)
Cl(4)	1.00432(8)	0.03930(4)	0.16226(5)	0.0572(3)
S(1)	1.5547(1)	0.12334(8)	0.5676(1)	0.0754(6)
S(1')	0.5176(1)	0.13325(7)	0.10565(8)	0.0749(6)
N(7)	1.1708(2)	0.1248(1)	0.5742(2)	0.0441(9)
N(7')	0.7478(2)	0.1775(1)	-0.0607(2)	0.0462(8)
C(2)	1.6785(4)	0.0759(3)	0.6435(3)	0.0793(2)
C(2')	0.3400(5)	0.1145(2)	0.0761(4)	0.1127(2)
C(3)	1.6328(4)	0.0173(2)	0.6883(3)	0.0713(2)
C(3')	0.2961(4)	0.0880(2)	-0.0131(5)	0.0893(4)
C(4)	1.4716(2)	-0.0018(1)	0.6673(2)	0.0661(9)
C(4')	0.3987(3)	0.0805(1)	-0.0737(2)	0.0710(1)
C(5)	1.4174(3)	0.0638(1)	0.5857(2)	0.0457(1)
C(5')	0.5449(3)	0.1047(2)	-0.0077(2)	0.0507(1)
C(6)	1.2653(3)	0.0699(2)	0.5293(2)	0.0463(1)
C(6')	0.6898(3)	0.1016(2)	-0.0378(2)	0.0503(1)
C(8)	1.2262(4)	0.2035(2)	0.5819(2)	0.0597(1)
C(8')	0.6452(4)	0.2206(2)	-0.1387(3)	0.0637(2)
C(9)	1.1384(4)	0.0984(2)	0.6720(3)	0.0740(2)
C(9')	0.8936(3)	0.1680(2)	-0.0891(2)	0.0627(2)

Table 3. Bond lengths (Å) and bond angles(°)

Co-Cl(1)	2.3106(8)	N(7')-C(6')	1.500(4)
Co-Cl(2)	2.3041(7)	N(7')-C(8')	1.492(4)
Co-Cl(3)	2.2650(8)	N(7')-C(9')	1.494(4)
Co-Cl(4)	2.2560(8)	C(2)-C(3)	1.312(6)
S(1)-C(2)	1.625(4)	C(2')-C(3')	1.286(8)
S(1)-C(5)	1.712(3)	C(3)-C(4)	1.516(4)
S(1')-C(2')	1.664(4)	C(3')-C(4')	1.381(7)
S(1')-C(5')	1.678(3)	C(4)-C(5)	1.613(3)
N(7)-C(6)	1.514(4)	C(4')-C(5')	1.543(4)
N(7)-C(8)	1.481(4)	C(5)-C(6)	1.482(4)
N(7)-C(9)	1.486(4)	C(5')-C(6')	1.486(4)
Cl(1)-Co-Cl(2)	110.43(3)	S(1)-C(2)-C(3)	116.3(3)
Cl(1)-Co-Cl(3)	108.03(3)	S(1')-C(2')-C(3')	115.0(4)
Cl(1)-Co-Cl(4)	109.24(3)	C(2)-C(3)-C(4)	119.2(3)
Cl(2)-Co-Cl(3)	106.14(3)	C(2')-C(3')-C(4')	117.3(4)
Cl(2)-Co-Cl(4)	109.43(3)	C(3)-C(4)-C(5)	98.0(2)
Cl(3)-Co-Cl(4)	113.52(3)	C(3')-C(4')-C(5')	105.5(3)
C(2)-S(1)-C(5)	93.1(2)	S(1)-C(5)-C(4)	113.2(2)
C(2')-S(1')-C(5')	92.3(2)	S(1)-C(5)-C(6)	123.5(2)
C(6)-N(7)-C(8)	113.9(2)	C(4)-C(5)-C(6)	123.3(2)
C(6)-N(7)-C(9)	112.9(2)	S(1')-C(5')-C(4')	109.8(2)
C(8)-N(7)-C(9)	111.1(2)	S(1')-C(5')-C(6')	124.5(2)
C(6')-N(7')-C(8')	113.2(2)	C(4')-C(5')-C(6')	125.7(2)
C(6')-N(7')-C(9')	109.6(2)	N(7)-C(6)-C(5)	114.6(2)
C(8')-N(7')-C(9')	111.7(2)	N(7')-C(6')-C(5')	114.0(2)




Figure 1. The molecular structure of bis(N,N-dimethyl-2-thio-phenemethylammonium)tetrachlorocobaltate(II). Drawing of single molecule showing 50% probability ellipsoids. (Cl---H-N hydrogen bonds are shown in thin lines with bond distances Cl₁---H and Cl₂---H being 2.40 and 2.38 Å, respectively)

and water are reported to be 2.29Å(8). But in (DMTMAH)₂CoCl₄, we found two different type of bond lengths. The four Co-Cl observed bond lengths are 2.3106(8), 2.3041(7), 2.2650(8), and 2.2560(8) Å, respectively. Two chlorine atoms involved in longer bond lengths have been noted to participate in the hydrogen bonding with DTMAH⁺ unit, *i.e.* Co-Cl---H-N. The effect of the intramolecular hydrogen bonding results in the increase in the Co-Cl bond lengths. In most amine compounds such as (CH₃)₃NHCl, (CH₃)₂NH₂Cl, and (NH₃OH)Cl, the N-H---Cl bond lengths are reported to be 3.10±0.1 Å (9). The N-H---Cl bond lengths estimated for

this compounds are 3.16 Å and the N-H---Cl bond angles are 160° for (DMTMAH)₂CoCl₄, which are typical structural characteristics found in amine hydrogen-halide system.

Acknowledgment. This work was supported by KOSEF (Korean Science and Engineering Foundation, Grant 921-0200-002-2).

Supplementary Material Available. Detailed descriptions of X-ray works, listings of anisotropic thermal parameters, hydrogen positional parameters, and observed and calculated structure factors are available from one of the authors (S.N. Choi) upon request.

References

1. Kim, Y. I.; Choi, S. N.; Ro, C. U. *Bull. Kor. Chem. Soc.* **1994**, *15*, 549.
2. Zaghariassen, W. H. *Acta Cryst.* **1963**, *16*, 1139.
3. Cromer, D. T.; Waber, J. T. *International Tables for X-ray Crystallography*; The Kynoch Press: Birmingham, England, 1974; Vol IV, Table 2. 2B.
4. Ibers, J. A.; Hamilton, W. C. *Acta Cryst.* **1964**, *17*, 781.
5. Cromer, D. T. *International Tables for X-ray Crystallography*; The Kynoch Press: Birmingham, England, 1974; Vol IV, Table 2.3.1.
6. Cruickshank, D. W. J. *Acta Cryst.* **1949**, *2*, 154.
7. MolEN. *An Interactive Structure Solution Procedure*; Enraf-Nonius, Delft, The Netherlands, 1990.
8. Ozutsumi, K.; Tohji, K.; Udagawa, Y.; Ishiguro, S. *Bull. Chem. Soc. Jpn.* **1991**, *64*, 1528.
9. Wells, A. F. *Structural Inorganic Chemistry*; Oxford; 1984.

ERRATUM

BKCS VOL. 16, NO. 6 Contents and Author Index on. P 489.

Jung Ho Park → Jeong Ho Park